

Sheaves of maximal intersection and multiplicities of stable log maps

JINWON CHOI

MICHEL VAN GARREL

SHELDON KATZ

NOBUYOSHI TAKAHASHI

A great number of theoretical results are known about log Gromov-Witten invariants [24, 13, 1], but few calculations are worked out. In this paper we restrict to surfaces and to genus 0 stable log maps of maximal tangency. We ask how various natural components of the moduli space contribute to the log Gromov-Witten invariants. The first such calculation [23, Proposition 6.1] by Gross-Pandharipande-Siebert deals with multiple covers over rigid curves in the log Calabi-Yau setting. As a natural continuation, in this paper we compute the contributions of non-rigid curves in the log Calabi-Yau setting and that of the union of two rigid curves in general position. For the former, we construct and study a moduli space of “logarithmic” 1-dimensional sheaves. For the latter, we explicitly describe the components of the moduli space and work out the logarithmic deformation theory in full, which we then compare with the deformation theory of the analogous relative stable maps.

14N35, 14A21, 14B10, 14D15, 14D20, 14J26

Contents

1	Introduction	2
1.1	Idealized geometries	2
1.2	Overview of methods	3
1.3	Log BPS numbers	5
1.4	Analogy with K3 surfaces	7
1.5	Contribution of curves with two image components	8
1.6	Applications of Theorem 1.14 and future directions	9
2	Nonsingularity of the relative compactified Picard scheme	10
3	Basic stable log maps	17
4	Proof of Theorem 1.14	19
Bibliography		34

1 Introduction

Let X be a smooth surface, let D be an effective divisor on X and denote by D_{sm} the smooth part of D . An \mathbb{A}^1 -curve on (X, D) is a proper irreducible curve C on X such that the normalization of $C \setminus D$ is isomorphic to \mathbb{A}^1 . We calculate the contributions of the following curves to the genus 0 log Gromov-Witten invariants of maximal tangency:

- (A) Corollary 1.12: Under the assumption that X is a projective rational surface, $K_X + D \sim 0$ and $P \in D_{\text{sm}}$, an \mathbb{A}^1 -curve C which is smooth at $P = C \cap D$.
- (B) Theorem 1.14: The sum of two immersed \mathbb{A}^1 -curves Z_1, Z_2 with $(K_X + D) \cdot Z_i = 0$, intersecting at $P \in D_{\text{sm}}$ in a general way.

In the proof of (A), we study a moduli space of ‘‘logarithmic’’ 1-dimensional sheaves, which shows an intriguing analogy with the case of K3 surfaces. For (B), a rather concrete description of a space of stable log maps is given.

1.1 Idealized geometries

Gromov-Witten invariants were devised as a virtual count of curves in projective or compact symplectic manifolds. However, their relationship with actual counts of curves, even when understood, is often quite subtle.

Let Y be a Calabi-Yau (CY) threefold and consider its genus 0 Gromov-Witten (GW) invariants $N_\beta(Y)$ for $\beta \in H_2(Y, \mathbb{Z})$. They are rational numbers in general, and BPS numbers $n_\beta(Y)$ were proposed as underlying \mathbb{Z} -valued invariants. They were originally defined via the recursive relationship

$$(1-1) \quad N_\beta(Y) = \sum_{k|\beta} \frac{1}{k^3} n_{\beta/k}(Y).$$

Still, even in the case of a compact Y , typically the BPS numbers $n_\beta(Y)$ are enumerative only in low degrees, in the sense that they agree with the count of rational curves. For example, if Y is a general quintic threefold, then $n_d(Y)$ equals the number of rational curves of degree d only when $d \leq 9$. For larger degrees, the story is more subtle.

Equation (1–1) is derived by postulating that Y symplectically deforms to an *idealized geometry* \tilde{Y} where all rational curves are infinitesimally rigid, i.e. have normal bundle $\mathcal{O}(-1) \oplus \mathcal{O}(-1)$. If such a curve is in class β , then its contribution to $N_{l\beta}(\tilde{Y})$ for $l \in \mathbb{N}$ is given by $1/l^3$ [4, 29, 34, 46], leading to the above formula. By deformation invariance one would then conclude that $n_\beta(Y)$ is a count of rational curves in \tilde{Y} . While the existence of such an idealized geometry is unknown, it is remarkable that the so defined $n_\beta(Y)$ are integers [25].

Assume instead that Y is the local Calabi-Yau threefold given as the total space $\text{Tot } \mathcal{O}_X(K_X)$ of the canonical bundle over a del Pezzo surface X . Then the enumerative interpretation of the local BPS numbers $n_\beta(K_X) := n_\beta(Y)$ is even more mysterious. Not only is their relationship to counts of rational curves in Y previously not known, they also are alternating negative with interesting divisibility properties [17, Conjecture 1.2]. As an illustration, the BPS numbers for local \mathbb{P}^2 in degrees d up to 6 are 3, -6, 27, -192, 1695, -17064, all of which are divisible by $3d$, a conjecture which was proven in [7] based on [22, 8].

An interpretation of $n_\beta(K_X)$, which also makes it clear why they are integral, was given using moduli spaces of sheaves. Denote by $M_{\beta,1}(X)$ the (smooth) moduli space of $(-K_X)$ -stable 1-dimensional sheaves of class β and of holomorphic Euler characteristic 1 on X , and let $w := -K_X \cdot \beta$. By [27, 11, 43, 44, 45], the genus 0 local BPS invariants can be identified with the topological Euler characteristics of $M_{\beta,1}(X)$:

$$n_\beta(K_X) = (-1)^{w-1} e(M_{\beta,1}(X)).$$

Another interpretation comes from log geometry. Based on the predictions of [42], in [17, 16] we started a program to show that $(-1)^{w-1} n_\beta(K_X)/w$ is a count of log curves in the surface X , namely that it equals the log BPS invariants

of Definition 1.7. Let D be a smooth anticanonical curve on X . Denote by $\overline{\mathcal{M}}_\beta(X, D)$ the moduli space of genus 0 basic stable log maps [24, 13, 1] in X of class β and of maximal tangency with D , see Section 3. From this space one defines the log Gromov-Witten invariants $\mathcal{N}_\beta(X, D)$, which virtually count \mathbb{A}^1 -curves, i.e. curves C such that the normalizations of $C \setminus D$ are isomorphic to \mathbb{A}^1 .

While there are a great number of theoretical results about log Gromov-Witten invariants, there are very few worked out examples. One of the aims of this paper is to remedy to that shortcoming. Our two main results will apply to a broad range of computations. We were informed of one such application in the upcoming work [5].

The stable log maps can meet D in a finite number of points and for such a point $P \in D$, one can consider $\mathcal{N}_\beta^P(X, D)$, the log GW invariant at P . We say the triple (X, D, P) is an *idealized log CY geometry for β* if $\mathcal{N}_\beta^P(X, D)$ equals the number of \mathbb{A}^1 -curves of class β at P . The advantage of the log setting is that for generic P and general D , the expectation is that (X, D, P) is idealized. The disadvantage is that there always are points P where (X, D, P) is not idealized, so looking at idealized geometries only captures a part of the moduli space of stable log maps.

At this point, one may define the log BPS numbers as the number of \mathbb{A}^1 -curves in an idealized log CY geometry, which conjecturally is equivalent to Definition 1.7. The next step then is to understand how \mathbb{A}^1 -curves contribute to $\mathcal{N}_\beta(X, D)$ in non-idealized log geometries. Unlike the CY case, there are countably many ways in which \mathbb{A}^1 -curves contribute to (higher degree) log GW invariants. The case of multiple covers over rigid integral curves was treated in [23, Proposition 6.1]. In this paper we treat the next two cases: of non-rigid \mathbb{A}^1 -curves and of two rigid distinct \mathbb{A}^1 -curves glued together. We expect that combining these 3 cases will lead to a solution of the general case.

In non-idealized geometries, passing from \mathbb{A}^1 -curves to virtual counts is related to surprisingly interesting geometry. For example, in [16, Proposition 1.16] we proved that the contribution of multiple covers over rigid \mathbb{A}^1 -curves to the log BPS numbers is given as the Donaldson-Thomas invariants of loop quivers. In the first part of this paper, we introduce a certain moduli space \mathcal{MMI}_β of sheaves of maximal intersection, which can be regarded as a logarithmic analogue of $M_{\beta,1}(S)$. Using \mathcal{MMI}_β , we show that \mathbb{A}^1 -curves in (X, D) share the same properties as rational curves in K3 surfaces.

Let us start with a maximally tangent stable log map of the simplest type, namely $f : \mathbb{P}^1 \rightarrow C \subset X$ with f immersed and C an integral rational curve maximally tangent to D . Such a curve contributes 1 to $\mathcal{N}_\beta(X, D)$. In other words, the naive multiplicity of the \mathbb{A}^1 -curve $C \setminus D$ is the correct multiplicity. Let us consider possible degenerations of f . For example, by deforming D , two log maps with image curve nodal cubics might collapse to one log map with image curve a cuspidal cubic. Then the \mathbb{A}^1 -curve $C \setminus D$ is not immersed and it contributes 2 to $\mathcal{N}_\beta(X, D)$. More generally, we show (Corollary 1.12) that an integral rational curve C maximally tangent to D and smooth at D contributes its natural stable map multiplicity to $\mathcal{N}_\beta(X, D)$, i.e. the log structure introduces no new infinitesimal deformations.

In the second part of this paper, we give an in-depth description of the log deformation theory of stable log maps obtained by gluing two \mathbb{A}^1 -curves. If C_1 and C_2 are distinct immersed integral rational curves maximally tangent to D at the same point, smooth at that point and intersect there in a general way, then they will contribute to $\mathcal{N}_{[C_1]+[C_2]}(X, D)$, a contribution we calculate in Theorem 1.14. We compare it to the case of relative stable maps and find that the log structures more finely distinguish between the possible maps.

1.2 Overview of methods

The moduli space $\overline{\mathcal{M}}_\beta(X, D)$ admits a finite forgetful morphism to the moduli space of stable maps [47], and it is natural to ask about the interplay between infinitesimal deformations of the underlying stable maps and infinitesimal deformations of the log structures. While this is difficult to answer in general, we get explicit solutions in terms of topological data for certain components of dimension 0. In this paper, we compute the contributions of such 0-dimensional components of $\overline{\mathcal{M}}_\beta(X, D)$ to the associated log Gromov-Witten invariants and log BPS numbers.

The 0-dimensional components of $\overline{M}_\beta(X, D)$ we consider here fall into two categories. The simplest components are built from \mathbb{A}^1 -curves C of class β . In the first part of this paper, we deal with such curves.

The arguments are modelled on the case of K3 surfaces. Let S be a K3 surface, γ a curve class on S and C a rational curve of class γ on S . Then the multiplicity of C is $e(\overline{\text{Pic}}^0(C))$, the Euler characteristic of the compactified Jacobian $\overline{\text{Pic}}^0(C)$ of C , by [18] (see also [6]). Let us elaborate a little. For a rational curve C we consider the moduli space $M_{0,0}(C, [C])$ of genus 0 stable maps to C , which is a thickened point corresponding to the normalization map $n : \mathbb{P}^1 \rightarrow C$. Let $l(C) := l(M_{0,0}(C, [C]))$ be its length. If C has only planar singularities, then $l(C)$ is equal to $m(C)$, the degree of the genus 0 locus in the versal deformation space of C ([18, Theorem 1]), which in turn is equal to $e(\overline{\text{Pic}}^0(C))$ by [18, Theorem 2].

As a curve on a surface S , the natural multiplicity would be $l(M_{0,0}(S, \gamma), n)$, the length of the moduli space of genus 0 stable maps to S at n , and it is equal to $l(C)$ if S is a K3 surface ([18, Theorem 2]). The key fact used in the proof is the smoothness of the relative compactified Jacobian over the complete linear system (at the points over C), or equivalently, of the moduli space $M_{\gamma,1}(S)$ of stable 1-dimensional sheaves of class γ , proven in [35].

Remark 1.1 If C is a rational curve with planar singularities and \tilde{C} is its minimal unibranch partial normalization, then $m(C) = m(\tilde{C})$. In particular, if C is immersed, i.e. the differential of n is nowhere vanishing, then there are no infinitesimal deformations and $m(C) = 1$ ([6, Proposition 3.3], [18, § 1]). In general, $m(C)$ is a product over the singularities of C of factors depending on the analytic type ([18, § 1], [6, Proposition 3.8], [39]). See [6, § 4] for explicit calculations.

Now we return to the case of an \mathbb{A}^1 -curve C in (X, D) . Assume that C is smooth at $P = C \cap D$ and denote by $n : \mathbb{P}^1 \rightarrow C$ the normalization map. Then one of our main results, Corollary 1.12, states that n contributes $l(C)$ to the log Gromov-Witten invariant. This follows from two facts: (i) the infinitesimal deformations of n as a log map can be identified with the infinitesimal deformations of the underlying stable map preserving the maximal tangency condition, and (ii) such infinitesimal deformations of n factor scheme-theoretically through C .

For the proof of (ii), we introduce a certain moduli space \mathcal{MMI}_β of sheaves of maximal intersection, which can be regarded as a logarithmic analogue of $M_{\gamma,1}(S)$. Just as in the case of K3 surfaces, we show the smoothness of \mathcal{MMI}_β (Theorem 1.11), from which we deduce that infinitesimal deformations of n as a log map factor through C . This might give a glimpse into a logarithmic version of sheaf-theoretic methods in curve counting, in analogy to the interpretation of genus 0 BPS numbers as Donaldson-Thomas invariants. Also, the relation between $M_{\beta,1}(X)$ and \mathcal{MMI}_β will be the subject of further investigation.

Before explaining how these results relate to the local BPS numbers, let us introduce a little more general setting. Let X be a smooth projective surface. In the first part of this paper, we will often require X to be *regular*, by which we mean that its irregularity $h^1(\mathcal{O}_X)$ vanishes. We denote by D an effective divisor on X . We will sometimes require additional conditions on X and D .

Definition 1.2 Assume X is regular, let $P \in D$ and let $\beta \in H_2(X, \mathbb{Z})$ be a curve class. Consider the linear system $|\mathcal{O}_X(\beta, P)|$ of curves of class β that meet D maximally at P (see Definition 1.9) and let $L \subseteq |\mathcal{O}_X(\beta, P)|$. We consider the following condition on L :

Condition (\star) Every rational curve $C \in L$ that is unibranch at P is in fact smooth at P .

In the setting of Theorem 1.3 below, we expect Condition (\star) to hold for general choices of (X, D) .

From Corollary 1.13 we derive an enumerative meaning of the local BPS numbers $n_\beta(K_X)$ subject to Conjecture 1.8 ([16, Conjecture 1.3]). This is a BPS version of the log-local principle pursued in [21, 19, 9, 10, 36]. Notice that Conjecture 1.8 is proven for $X = \mathbb{P}^2$ in [7] based on [22, 8], and for any del Pezzo surface and classes of arithmetic genus ≤ 2 in [17, 16].

Theorem 1.3 (Log-local principle for BPS numbers) *Let X be a del Pezzo surface, let D be a smooth anticanonical curve on X and let $P \in D$ be β -primitive (Definition 1.5). Then there is a finite number of rational curves in $|\mathcal{O}_X(\beta, P)|$. Assume that:*

- Conjecture 1.8 holds for X . (For example when $X = \mathbb{P}^2$ by [8, 7], or for arithmetic genus ≤ 2 by [17, 16].)
- Condition (\star) holds for $|\mathcal{O}_X(\beta, P)|$.

Then

$$(1-2) \quad n_\beta(K_X) = (-1)^{\beta \cdot D - 1} (\beta \cdot D) \sum_{\substack{C \in |\mathcal{O}_X(\beta, P)| \\ \text{rational and} \\ \text{unibranch at } P}} l(C).$$

Note that $l(C) = e(\overline{\text{Pic}}^0(C))$ by [18].

Note that each component of $\overline{M}_\beta(X, D)$ with tangency at a β -primitive P comes from an (irreducible) \mathbb{A}^1 -curve, explaining the terms $l(C)$ in Equation (1–2).

The other category of zero-dimensional components consists of stable log maps $C \rightarrow X$ with image consisting of two distinct rational curves Z_1 and Z_2 , each maximally tangent to D at P . This situation occurs very often for a non- β -primitive point P . For example, if there are two \mathbb{A}^1 -curves Z_1 and Z_2 in the class β which meet D at the same point P , then their sum contributes to $\overline{M}_{2\beta}(X, D)$.

In this case, C consists of three components, two mapped to Z_1 and Z_2 and one mapped to P . It is not straightforward to see what log structure C should have, unlike the case treated in the first part. Moreover, it turns out that the moduli spaces of stable log maps and relative stable maps are not isomorphic in the neighborhood of such a map, although they are guaranteed to give the same numerical invariants by [2].

As the second main result of this paper, we calculate in Theorem 1.14 the number of such log maps and the contributions of each to the log Gromov-Witten invariants in terms of intersection data, subject to genericity conditions. The proof involves a rather concrete (and long) calculation on infinitesimal families of log maps. We explicitly separate the infinitesimal deformations coming from the underlying stable maps from the ones coming from the log structure and explicitly describe both. This result sheds light on the interplay of the log structures with the underlying stable maps.

Now let us give a little more detailed explanation on what we are going to deal with.

1.3 Log BPS numbers

Let X be a smooth projective surface and let $\beta \in H_2(X, \mathbb{Z})$ be a curve class. We write $w = \beta \cdot D$ and assume that $w > 0$. If X is regular, then there is a unique $L \in \text{Pic}(X)$ such that $c_1(L)$ is Poincaré dual to β . By $\beta|_D$ we mean $L|_D \in \text{Pic}^w(D)$. For X regular, set

$$(1-3) \quad D(\beta) := \{P \in D_{\text{sm}} : \beta|_D = \mathcal{O}_D(wP) \text{ in } \text{Pic}^w(D)\}.$$

Remark 1.4 (1) If D is an elliptic curve, $D(\beta)$ is a torsor for $\text{Pic}^0(D)[w] \simeq \mathbb{Z}/w\mathbb{Z} \times \mathbb{Z}/w\mathbb{Z}$ (cf. [16, Lemma 2.14]).

(2) If $D = \bigcup_{i=1}^k D_i$ is the decomposition into irreducible components and $D(\beta) \neq \emptyset$, then $D(\beta) \subset D_{i_0}$ for a unique i_0 and $\beta|_{D_i} \sim 0$ for $i \neq i_0$.

Definition 1.5 Let $P \in D(\beta)$. Then P is β -primitive if there is no decomposition into non-zero pseudo-effective classes $\beta = \beta' + \beta''$, with $\beta' \cdot D > 0$ and such that $P \in D(\beta')$.

Proposition 1.6 (Proposition 4.11 in [16]) Assume that X is a del Pezzo surface and that D is a smooth anticanonical curve on X . If the pair (X, D) is general, then there is a β -primitive point $P \in D(\beta)$.

If (X, D) is a log smooth pair, denote by $\mathcal{N}_\beta(X, D)$ the genus 0 log Gromov-Witten invariant of maximal tangency and class β of (X, D) , whose definition we review in Section 3. If X is regular, D is smooth and $p_a(D) > 0$, then $D(\beta)$ is a finite set and the moduli space decomposes into a disjoint union according to $P \in D(\beta)$:

$$\overline{\mathbf{M}}_\beta(X, D) = \bigsqcup_{P \in D(\beta)} \overline{\mathbf{M}}_\beta^P(X, D).$$

Hence we can define the contribution $\mathcal{N}_\beta^P(X, D)$ from each P so that

$$\mathcal{N}_\beta(X, D) = \sum_{P \in D(\beta)} \mathcal{N}_\beta^P(X, D)$$

holds.

We often take D to be anticanonical. Note that a regular surface with a nonzero anticanonical curve is rational by Castelnuovo's criterion.

Definition 1.7 Assume that X is rational and D is smooth anticanonical, and let $P \in D(\beta)$. The *log BPS number at P* , m_β^P , is defined recursively via

$$\mathcal{N}_\beta^P(X, D) = \sum_{k|\beta} \frac{(-1)^{(k-1)w/k}}{k^2} m_{\beta/k}^P,$$

where $m_{\beta'}^P := 0$ if $P \notin D(\beta')$.

Conjecture 1.8 (Conjecture 1.3 in [16]) For all $P, P' \in D(\beta)$,

$$m_\beta^P = m_{\beta'}^{P'}.$$

Equivalently, for $P \in D(\beta)$,

$$n_\beta(K_X) = (-1)^{\beta \cdot D - 1} (\beta \cdot D) m_\beta^P.$$

What makes this highly nontrivial is that $\overline{\mathbf{M}}_\beta^P(X, D)$ can be quite different according to the local geometry of D near P (see [16, §§6.1] and §§1.6). Conjecture 1.8 was proven for \mathbb{P}^2 in [8, 7] using [22].

Definition 1.9 Assume that X is regular and $P \in D_{\text{sm}}$. We denote by $|\beta|$ the linear system of curves of class β and set

$$|\mathcal{O}_X(\beta, P)| := \{C \in |\beta| : C|_D \supseteq wP \text{ as subschemes of } D\},$$

as well as its open subsets

$$|\mathcal{O}_X(\beta, P)|^\circ := \{C \in |\beta| : C|_D = wP \text{ as subschemes of } D\}$$

and

$$|\mathcal{O}_X(\beta, P)|^{\circ\circ} := \{C \in |\beta| : C|_D = wP \text{ as subschemes of } D \text{ and } C \text{ is integral}\}.$$

Moreover, we write

$$p_a(\beta) := \frac{1}{2} \beta(\beta + K_X) + 1$$

for the arithmetic genus of members of $|\beta|$.

Remark 1.10 For a rational X with smooth anticanonical D , notice that the set of rational curves in $|\mathcal{O}_X(\beta, P)|^\circ$ is identified with the set of rational curves in $|\mathcal{O}_X(\beta, P)|$. For a regular surface X , a curve D on X and a β -primitive point $P \in D(\beta)$, note that every member of $|\mathcal{O}_X(\beta, P)|^\circ$ is an integral curve, i.e. $|\mathcal{O}_X(\beta, P)|^\circ = |\mathcal{O}_X(\beta, P)|^{\circ\circ}$.

In Section 2 we construct certain moduli spaces, denoted \mathcal{MMI}_β and \mathcal{MMI}_β^P , associated to any smooth surface X and a curve D on it. These moduli spaces parametrize certain sheaves supported on integral curves and having “maximal intersection” with D . For \mathcal{MMI}_β^P with $P \in D_{\text{sm}}$, the additional condition is imposed that the tangency is at P . If X is regular and the Abel map of D is immersive at P (e.g. if D is integral with $p_a(D) > 0$ or D is anticanonical in rational X ; see Lemma 2.8(3)), then, by Lemma 2.8(2), \mathcal{MMI}_β decomposes scheme-theoretically as a disjoint union

$$\mathcal{MMI}_\beta = \coprod_{P \in D(\beta)} \mathcal{MMI}_\beta^P.$$

Theorem 1.11 (=Theorem 2.12) *Let X be a smooth projective rational surface, D an anticanonical curve on X and $P \in D_{\text{sm}}$. Then \mathcal{MMI}_β and \mathcal{MMI}_β^P are nonsingular of dimension $2p_a(\beta) = \beta^2 - w + 2$.*

Consequently, the relative compactified Picard scheme over $|\mathcal{O}_X(\beta, P)|^\circ$ is nonsingular at a point $[F]$ over $[C]$ if F is an invertible \mathcal{O}_C -module near P (or, equivalently, $F|_D \cong \mathcal{O}_C|_D$).

We use this theorem to calculate the contribution of an \mathbb{A}^1 -curve to the log Gromov-Witten invariant. The simplest components in the moduli space of genus 0 basic stable log maps consist of (possibly thickened) points. We will mainly be concerned with the case $(K_X + D) \cdot \beta = 0$, since otherwise the virtual dimension is nonzero. Then, one such case arises from an irreducible \mathbb{A}^1 -curve. The following result, proven in [16] subject to Theorem 1.11 (and Lemma 2.8(2)), calculates the contribution of such a point to $\mathcal{N}_\beta(X, D)$.

Corollary 1.12 (Proposition 1.7(3) in [16]) *Let X be a smooth projective rational surface and D an anticanonical curve. Let C be an irreducible rational curve of class β maximally tangent to D at $P \in D(\beta)$. Denote the normalization map by $n : \mathbb{P}^1 \rightarrow C$ and assume that C is smooth at P . Then n contributes $l(C) := l(M_{0,0}(C, [C]))$ to $\mathcal{N}_\beta(X, D)$.*

Consequently, we have:

Corollary 1.13 (Proposition 1.7(5) in [16]) *Let X be a del Pezzo surface and D a smooth anticanonical curve on X , and let $P \in D(\beta)$ be β -primitive. Assume that $|\mathcal{O}_X(\beta, P)|^\circ$ satisfies Condition (\star) . Then*

$$m_\beta^P = l(\overline{M}_\beta^P(X, D)) = \sum_{\substack{C \in |\mathcal{O}_X(\beta, P)|^\circ \\ \text{rational and} \\ \text{unibranch at } P}} l(C).$$

In Corollary 1.13, Condition (\star) is needed to ensure that the compactified Picard variety of C is contained in \mathcal{MMI}_β .

1.4 Analogy with K3 surfaces

Assume now that X is a del Pezzo surface and that D is smooth anticanonical. Let $\beta \in H_2(X, \mathbb{Z})$ be a curve class, set $h = p_a(\beta)$ and choose $P \in D(\beta)$ to be β -primitive. Then the linear system $|\mathcal{O}_X(\beta, P)|$ is of dimension h [16, Proposition 4.15].

Denote by S a K3 surface. To S are associated genus 0 BPS numbers $r_{0,h}$, see [37]. Remarkably [28], $r_{0,h}$ only depends on the arithmetic genus h of a curve class on S . Choose a complete linear system L of curves of arithmetic genus h . Then L is h -dimensional as is $|\mathcal{O}_X(\beta, P)|$. Under the assumption that all curves in L are integral, $r_{0,h}$ is given as the sum of $l(C) = e(\overline{\text{Pic}}^0(C))$ for C a rational curve in L ([48, 6, 14, 18]). This is in perfect analogy to Corollary 1.13.

By [16, Lemma 4.10], saying that P is β -primitive amounts to P being of maximal order for a suitably chosen zero element. Keeping track of the order of P is analogous for S to pass to reduced classes [37, Section 5.3.1]. While the Gromov-Witten theory of S is trivial due to non-algebraic deformations, it still admits non-zero reduced invariants when the deformation space is restricted to algebraic deformations. Assume that there is a type II degeneration of S into two components, one of them being (X, D) , the other one (X', D') , so that D is identified with D' . Consider the normal bundles N and N' of X , resp. X' , in the family. A necessary smoothability condition is that $N|_D = N|_{D'}$.

Let now C and C' be rational curves in (X, D) and (X', D') meeting $D = D'$ in the same point P of maximal tangency. For $C \cup_P C'$ to be algebraically deformable to the general fiber, the orders of tangency $C \cdot D$ and $C' \cdot D'$ should be the same. In addition, with respect to a common zero, P should have the same order in $D([D])$ as in $D'([C'])$. For example, P could be both $[C]$ - and $[C']$ -primitive. Without matching the order, there can not be an algebraic deformation.

In [16], we calculated m_β^P , for β -primitive P and $p_a(\beta) \leq 2$. We found (Theorems 1.8 and 1.9) in these cases that m_β^P depends only on the intersection number $e(S) - \eta$ for η the number of line classes l with $\beta \cdot l = 0$. Similarly, $r_{0,h}$ only depends on the intersection number $\beta \cdot \beta = 2h - 2$ [28].

The analogy carries over to SYZ fibrations. For K3 surfaces, [33, 32] proves that counts of Maslov index 0 disks with boundary on a SYZ-fiber correspond to tropical curves in the base. In the case of \mathbb{P}^2 , the analogous correspondence [22] is between log BPS numbers and tropical curves in the scattering diagram.

1.5 Contribution of curves with two image components

In the second part of this paper, Sections 3 and 4, we consider another type of zero-dimensional component, where the images of the stable log maps consist of two maximally tangent rational curves.

Theorem 1.14 *Let (X, D) be a pair consisting of a smooth surface and an effective divisor. Denote by $\overline{M}_\beta = \overline{M}_\beta(X, D)$ the moduli stack of maximally tangent genus 0 basic stable log maps of class β to the log scheme associated to (X, D) .*

Let Z_1 and Z_2 be proper integral curves on X satisfying the following:

- (1) Z_i is a rational curve of class β_i maximally tangent to D ,
- (2) $(K_X + D) \cdot \beta_i = 0$,
- (3) $Z_1 \cap D$ and $Z_2 \cap D$ consist of the same point $P \in D_{\text{sm}}$, and
- (4) The normalization maps $f_i : \mathbb{P}^1 \rightarrow Z_i$ are immersive and $(Z_1 \cdot Z_2)_P = \min\{d_1, d_2\}$, where $d_i = D \cdot Z_i$.

Write $d_1 = de_1, d_2 = de_2$ with $\gcd(e_1, e_2) = 1$. Then there are d stable log maps in $\overline{M}_{\beta_1+\beta_2}$ whose images are $Z_1 \cup Z_2$, and they are isolated with multiplicity $\min\{e_1, e_2\}$.

When X is projective and (X, D) is log smooth, then these curves contribute $\min\{d_1, d_2\}$ to the log Gromov-Witten invariant $\mathcal{N}_{\beta_1+\beta_2}(X, D)$.

Remark 1.15 In Theorem 1.14, the condition that $(Z_1 \cdot Z_2)_P = \min\{d_1, d_2\}$ means that Z_1 and Z_2 are assumed to intersect generically at P . We expect that this condition is satisfied for general D when $Z_1 \neq Z_2$.

If $d_1 \neq d_2$, then as a consequence of the immersivity of f_i and Proposition 4.1(3), Z_1 and Z_2 are smooth at P and the condition $(Z_1 \cdot Z_2)_P = \min\{d_1, d_2\}$ holds. In the case $d_1 = d_2 = d$, in analytic coordinates x, y near P with $D = (y = 0)$, we can write $Z_i = (y = a_i x^d + \dots)$. Then $(Z_1 \cdot Z_2)_P = \min\{d_1, d_2\}$ translates into $a_1 \neq a_2$.

An example where this condition is obviously not satisfied is the case $Z_1 = Z_2$. In this case, the space of log maps with image cycle $Z_1 + Z_2$, as well as its contribution to the log Gromov-Witten invariant, is quite different ([23, Proposition 6.1]).

It is illuminating to compare Theorem 1.14 with the analogous result [41] for the relative stable maps of [30, 31]. Whereas there is only one relative stable map with multiplicity $\min\{d_1, d_2\}$, there are d log maps each with multiplicity $\min\{e_1, e_2\}$, making the same contribution as expected by [2]. This illustrates that there can be several ways of associating a log map to a relative map.

Theorem 1.14 is illustrated by Example 4.19, which the reader may consider as the running example for the second part of this paper. We fully work out the same example in the language of relative stable maps [30, 31] in Example 4.20 following [41].

1.6 Applications of Theorem 1.14 and future directions

The components that occur in $\overline{\mathcal{M}}_\beta(X, D)$ are classified by [16, Corollary 2.10] (see Proposition 4.1). Outside of the calculations of this paper, the only other components whose contribution to log Gromov-Witten invariants is known are multiple covers over rigid maximally tangent rational curves [23, Proposition 6.10].

Knowledge of the contributions of some components would allow for new enumerative calculations. As an illustration, consider $X = \mathbb{P}^2$ and the log BPS numbers m_{5H}^P of degree $5H \in H_2(\mathbb{P}^2, \mathbb{Z})$ of Definition 1.7 for $P \in D(5H)$. According to Conjecture 1.8 (proven in [8, 7]), m_{5H}^P is constant whether P is a flex point or a $5H$ -primitive point. For the latter, $m_{5H}^P = 113$ is an actual count of rational curves.

Take now P to be a flex point. Denote by k_5 the number of degree 5 rational curves maximally tangent to D at P . Cf. [16, Section 6] and [40], provided D is general, in lower degree there are 1 flex line, 2 nodal cubics and 8 nodal quartics. Taking the description of [16, Section 6], we have that

$$\begin{aligned} m_{5H}^P &= 113 = \text{DT}_5^{(2)} + 8 \cdot 3 \cdot \min\{4, 1\} + 2 \cdot \text{Contr}(3, 1^2) + k_5 \\ &= 5 + 24 + 2 \cdot \text{Contr}(3, 1^2) + k_5. \end{aligned}$$

Here $\text{DT}_5^{(2)}$ is the 5th 2-loop quiver invariant, which is the contribution of $5 : 1$ multiple covers over the flex line to m_{5H}^P [16, Proposition 6.4]. The term $3 \cdot \min\{4, 1\}$ is the contribution according to Theorem 1.14 of the 3 stable log maps with image (fixed quartic) \cup (flex line). $\text{Contr}(3, 1^2)$ is the unknown contribution of the component corresponding to stable log maps $\mathbb{P}^1 \cup \mathbb{P}^1 \cup \mathbb{P}^1 \rightarrow \mathbb{P}^2$ with the central component collapsed, the first component an immersion into one of the nodal cubics and the third component a $2 : 1$ cover over the flex line.

So knowing $\text{Contr}(3, 1^2)$ one would be able to calculate k_5 . In fact, local BPS numbers are calculated through local mirror symmetry [15] and so are the m_{dH}^P via Conjecture 1.8 (proven in [8, 7]). Hence, knowing the contributions of each component of $\overline{\mathcal{M}}_\beta(X, D)$ corresponding to stable log maps with reducible image would recursively allow to calculate k_d , the number of rational degree d curves maximally tangent to a flex point, for all d . Moreover, the same analysis holds for counts of maximally tangent rational curves at any other point $P \in D(dH)$.

Acknowledgements

We wish to thank Mark Gross and Dhruv Ranganathan for a discussion on multiplicities of stable log maps that motivated parts of this work. We wish to thank Alexei Oblomkov for discussions on multiplicities of stable maps to surfaces.

JC is supported by the Korea NRF grant NRF-2018R1C1B6005600. MvG is supported by the EC REA MSCA-IF-746554. SK is supported in part by NSF grant DMS-1502170 and NSF grant DMS-1802242. NT is supported by JSPS KAKENHI Grant Number JP17K05204. This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 746554.

2 Nonsingularity of the relative compactified Picard scheme

We start with a couple of lemmas that might belong to common knowledge.

Lemma 2.1 *If X is a regular surface, a connected component of $\text{Hilb}(X)$ containing a curve C is nonsingular and coincides with $|C|$.*

Proof Let Λ be a complete linear system and C a member of Λ . The first order deformations of C in $\text{Hilb}(X)$ are given by

$$\text{Hom}_{\mathcal{O}_X}(\mathcal{I}_C, \mathcal{O}_C) \simeq \text{Hom}_{\mathcal{O}_X}(\mathcal{O}_X(-C), \mathcal{O}_C) \simeq H^0(\mathcal{O}_C(C)).$$

From the short exact sequence $0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_X(C) \rightarrow \mathcal{O}_C(C) \rightarrow 0$ we obtain the exact sequence

$$0 \rightarrow H^0(\mathcal{O}_X) \rightarrow H^0(\mathcal{O}_X(C)) \rightarrow H^0(\mathcal{O}_C(C)) \rightarrow H^1(\mathcal{O}_X) = 0.$$

The space $H^0(\mathcal{O}_X(C))/H^0(\mathcal{O}_X)$ can be regarded as the tangent space of Λ at C , and this exact sequence shows that the natural map $T_C \Lambda \rightarrow T_C \text{Hilb}(X)$ is an isomorphism. Since Λ is projective and nonsingular (by definition) and is embedded into $\text{Hilb}(X)$, it can be identified with a connected component of $\text{Hilb}(X)$. \square

For a curve class β , let $M_\beta(X)$ denote the moduli space of stable 1-dimensional sheaves of class β on X with respect to a certain polarization. In the following, we will mainly consider sheaves F that can be regarded as torsion-free sheaves of rank 1 on integral curves of class β . Such a sheaf F defines a point $[F] \in M_\beta(X)$ for any polarization.

Lemma 2.2 *Let X be a smooth surface, β a curve class with $K_X \cdot \beta < 0$ and C an integral curve of class β . If F is a torsion-free rank 1 sheaf on C , then $M_\beta(X)$ is nonsingular of dimension $\beta^2 + 1$ at $[F]$.*

Proof The first order deformations of F in $M_\beta(X)$ are described by $\text{Ext}_{\mathcal{O}_X}^1(F, F)$ and one calculates

$$\sum_{i=0}^2 (-1)^i \dim \text{Ext}_{\mathcal{O}_X}^i(F, F) = -c_1(F)^2 = -\beta^2.$$

By stability of F (or, rather, by the arguments for the proof of stability), $\text{Hom}_{\mathcal{O}_X}(F, F) = \mathbb{C}$. Moreover, $\text{Ext}_{\mathcal{O}_X}^2(F, F)$ is dual to $\text{Hom}_{\mathcal{O}_X}(F \otimes_{\mathcal{O}_X} \mathcal{O}_X(-K_X), F)$, and the latter is 0 by the inequality $\chi(F \otimes_{\mathcal{O}_X} \mathcal{O}_X(-K_X)) - \chi(F) = -K_X \cdot C > 0$ and the stability of F . \square

Lemma 2.3 *Let C be an integral curve on a smooth surface X and F a torsion-free sheaf of rank 1 on C . Then locally near $[F]$, the Chow morphism $M_\beta(X) \rightarrow \text{Hilb}(X)$ lifts to an isomorphism of a neighborhood of $[F]$ in $M_\beta(X)$ to an open set in $\overline{\text{Pic}}(\mathcal{C}/\mathcal{H})$, where \mathcal{H} is the connected component of $\text{Hilb}(X)$ containing $[C]$ and \mathcal{C} is the universal subscheme over \mathcal{H} .*

Proof First, take a deformation of F in $M_\beta(X)$: Let T be a scheme over \mathbb{C} , $0 \in T$ a point, and \mathcal{F} a coherent sheaf on $X \times T$ which is flat over T , such that $\mathcal{F}_0 \cong F$ and \mathcal{F}_t is stable for any geometric point t of T . We may replace T by a neighborhood of 0 (actually, it suffices to take T to be a neighborhood of $[F]$ in $M_\beta(X)$), and we have the Chow morphism $\varphi : T \rightarrow \mathcal{H}$.

Let us show that, after shrinking T if necessary, the ideal of \mathcal{C}_T in $X \times T$ annihilates \mathcal{F} . For this purpose, we recall the definition of the Fitting ideal of \mathcal{F} . Let R denote the local ring of $X \times T$ at a point over 0 and M an R -module corresponding to \mathcal{F} . If we take a surjection $R^n \rightarrow M$, using the fact that F is pure of dimension 1 and \mathcal{F} is flat over T , we can show that the kernel is also free of rank n , i.e. we have an exact sequence

$$0 \rightarrow R^n \xrightarrow{\phi} R^n \rightarrow M \rightarrow 0.$$

The Fitting ideal is locally generated by $\det \phi$, and is globally well-defined, independent of the local resolution of \mathcal{F} . It is immediate to see that the Fitting ideal defines the flat family of subschemes \mathcal{C}_T corresponding to the Chow morphism $T \rightarrow \text{Hilb}(X)$. Now $\det \phi$ certainly annihilates M , since on R^n it can be written as the composition of ϕ and its adjoint. Hence \mathcal{F} can be regarded as a family of sheaves on \mathcal{C}_T .

Note that \mathcal{F}_t is a torsion-free sheaf on $(\mathcal{C}_T)_t$ for any geometric point t of T , since a torsion subsheaf would destabilize \mathcal{F}_t . By shrinking T , we may assume that $(\mathcal{C}_T)_t$ is integral for any t , and then \mathcal{F}_t is of rank 1 since its first Chern class is β . We therefore obtain a morphism $T \rightarrow \overline{\text{Pic}}(\mathcal{C}/\mathcal{H})$.

Conversely, if \mathcal{G} is a family of rank 1 torsion-free modules on a family of integral curves \mathcal{C}_T for some $T \rightarrow \mathcal{H}$, then it can be considered as a family of stable sheaves on X over T .

These correspondences are inverse to each other, and isomorphisms between families also coincide. Thus we have a local isomorphism of the moduli spaces. \square

We will use the following theorem on relative compactified Picard schemes.

Theorem 2.4 ([3]) *Let \mathcal{C}/S be a projective family of integral curves of arithmetic genus p_a that can be embedded into a smooth projective family of surfaces over S . Then its relative compactified Picard scheme is flat over S and the geometric fibers are integral locally complete intersections of dimension p_a .*

We return to the setting of a smooth surface X and a curve D on X . For a curve class β , let us denote $D.\beta$ by w , which we assume to be positive. For a scheme T over \mathbb{C} , we consider the following condition (*) on a coherent sheaf \mathcal{F} on $X \times T$:

- (a) \mathcal{F} is flat over T , and for each geometric point t of T , \mathcal{F}_t is a torsion-free sheaf of rank 1 on an integral curve C_t of class β , not contained in D .
- (b) There exists a section $\sigma : T \rightarrow D_{\text{sm}} \times T \hookrightarrow X \times T$ with $\mathcal{F}|_{D \times T} \cong \mathcal{O}_{w \cdot \sigma(T)}$ as $\mathcal{O}_{D \times T}$ -modules, where $w \cdot \sigma(T)$ is the closed subscheme of $D \times T$ defined by the w -th power of the ideal sheaf of $\sigma(T) \subset D \times T$.

We will later see that σ is unique.

Lemma 2.5 *For a sheaf \mathcal{F} satisfying condition (*), the following also holds:*

- *In a neighborhood of each point of $\sigma(T)$, the sheaf \mathcal{F} is isomorphic to the structure sheaf of the family of curves induced by the Chow morphism.*

Also, the conditions $\text{rank}_{C_t} \mathcal{F}_t = 1$ and $C_t \not\subseteq D$ follow from the rest of the conditions.

Proof From $\mathcal{F}|_{D \times T} \cong \mathcal{O}_{w \cdot \sigma(T)}$ and Nakayama's Lemma, \mathcal{F} is generated by 1 element near any point of $\sigma(T)$, giving rise to a surjective homomorphism $\mathcal{O}_{X \times T} \rightarrow \mathcal{F}$ locally. The kernel contains the ideal of the associated family of curves, and from the torsion-freeness, they coincide. \square

Definition 2.6 (1) We define a moduli functor \mathcal{MMI}_β (for “modules with maximal intersection”) on the category of schemes over \mathbb{C} as the sheafification of the presheaf

$$T \mapsto \{\mathcal{F} \mid \mathcal{F} \text{ is a sheaf on } X \times T \text{ satisfying the condition } (*)\} / \cong,$$

where \cong denotes isomorphisms of coherent sheaves on $X \times T$.

(2) For $P \in D_{\text{sm}}$, we define \mathcal{MMI}_β^P as the subfunctor of \mathcal{MMI}_β parameterizing families where σ can be locally taken to be the constant section $T \cong P \times T \hookrightarrow D_{\text{sm}} \times T$.

(3) We define a moduli functor \mathcal{MI}_β of integral curves on X of class β with maximal intersection with D : For a scheme T over \mathbb{C} , an element of $\mathcal{MI}_\beta(T)$ is a closed subscheme \mathcal{Z} of $X \times T$, flat over T , with fibers integral curves of class β such that the intersection of \mathcal{Z} and $D \times T$ is $w \cdot \sigma(T)$ for a section $\sigma : T \rightarrow D_{\text{sm}} \times T$.

(4) For $P \in D_{\text{sm}}$, we define \mathcal{MI}_β^P as the subfunctor of \mathcal{MI}_β parameterizing families where σ can be taken to be the constant section with value P .

Lemma 2.7 (1) The functor \mathcal{MMI}_β is represented by a locally closed subscheme of $M_\beta(X)$, and \mathcal{MMI}_β^P is represented by a closed subscheme of \mathcal{MMI}_β .

(2) The functor \mathcal{MI}_β is represented by a locally closed subscheme of $\text{Hilb}(X)$, and \mathcal{MI}_β^P is represented by a closed subscheme of \mathcal{MI}_β .

(3) We may also regard \mathcal{MMI}_β (resp. \mathcal{MMI}_β^P) as an open subscheme of $M_\beta(X) \times_{\text{Hilb}(X)} \mathcal{MI}_\beta$ (resp. $M_\beta(X) \times_{\text{Hilb}(X)} \mathcal{MI}_\beta^P$), or of the relative compactified Picard scheme over \mathcal{MI}_β (resp. \mathcal{MI}_β^P).

(4) There exist unique morphisms $\mathcal{MMI}_\beta \rightarrow D$ and $\mathcal{MI}_\beta \rightarrow D$ representing sections σ such that $\mathcal{F}|_{D \times T} \cong \mathcal{O}_{w \cdot \sigma(T)}$ locally over T and $\mathcal{Z}|_{D \times T} = w \cdot \sigma(T)$. These morphisms commute with the Chow morphism.

The spaces \mathcal{MMI}_β^P and \mathcal{MI}_β^P are the scheme theoretic inverse images of P by these morphisms.

Proof First we prove (1) and (4) for \mathcal{MMI}_β . Proofs of (2) and (4) for \mathcal{MI}_β are similar.

In $M_\beta(X)$, the condition (a) in (*) is an open condition. The condition that \mathcal{F} is generated by one section near D is also open, so these conditions define an open subscheme M° of $M_\beta(X)$.

Let \mathcal{F} be a family in M° over T . In a neighborhood of $\sigma(T)$, \mathcal{F} is isomorphic to the structure sheaf of a flat family of curves, which is a family of principal divisors. Hence, locally over T , the restriction $\mathcal{F}|_{D \times T}$ is isomorphic to the structure sheaf of a flat family of 0-dimensional subschemes of D of length w . This family can also be described as the one defined by the annihilator of $\mathcal{F}|_{D \times T}$, hence is determined by \mathcal{F} .

Thus we have a morphism $M^\circ \rightarrow \text{Hilb}^w(D)$, the latter being isomorphic to the symmetric w -th power $D^{(w)}$ of D . We claim that subschemes of the form wP are represented by the diagonal set $\Delta \subset D^{(w)}$ with the reduced induced structure. To show this, we can work with the formal neighborhood of P since we are concerned with infinitesimal deformations of 0-dimensional subschemes. In a formal coordinate x on D , the Hilbert scheme can be described as the spectrum of $\mathbb{C}[[a_1, \dots, a_w]]$ with the universal subscheme $x^w - a_1x^{w-1} + \dots + (-1)^w a_w = 0$. The diagonal set Δ , as a reduced closed subscheme, is given by $a_i = \binom{w}{i} (a_1/w)^i$ ($i = 2, \dots, w$). Consider a family over a complete local ring R , corresponding to $\varphi : \text{Spec } R \rightarrow \text{Spec } \mathbb{C}[[a_1, \dots, a_w]]$ given by $a_i = r_i$. If it satisfies the condition (b), with σ corresponding to $x \mapsto r \in R$, then $x^w - r_1x^{w-1} + \dots + (-1)^w r_w$ is equal to $(x - r)^w$ (this follows from the fact that the coefficients of x^0, x^1, \dots, x^{w-1} are the coordinates of the representing space, or more concretely, by writing $x^w - r_1x^{w-1} + \dots + (-1)^w r_w = (\text{unit})(x - r)^w$ and using Weierstrass preparation theorem). This means that $r_i = \binom{w}{i} r^i$ ($i = 1, \dots, w$) and φ factors through Δ .

Thus \mathcal{MMI}_β is the scheme theoretic inverse image of Δ .

In the calculation above, r is determined by (r_1, \dots, r_w) : Specifically, $r = r_1/w$. This shows the existence and uniqueness of $\mathcal{MMI}_\beta \rightarrow D$ as in (4), and $\mathcal{MMI}_\beta^P = \mathcal{MMI}_\beta \times_D P$ follows from the definition of \mathcal{MMI}_β^P .

(3) follows from the description above of families in \mathcal{MMI}_β as families in M° whose support curves have maximal intersection with D . \square

Lemma 2.8 Assume that X is a regular surface.

- (1) The space \mathcal{MI}_β^P can be identified with $|\mathcal{O}_X(\beta, P)|^\circ$ considered as a nonsingular variety.
- (2) Assume furthermore that the Abel map of D is immersive at P and C belongs to \mathcal{MI}_β^P . Then, in a neighborhood of $[C]$, the morphism $\mathcal{MI}_\beta \rightarrow D$ representing the intersection point is scheme-theoretically the constant map with value P .

Consequently, if the Abel map of D is immersive at each $P \in D(\beta)$,

$$\mathcal{MI}_\beta = \coprod_{P \in D(\beta)} \mathcal{MI}_\beta^P$$

and

$$\mathcal{MMI}_\beta = \coprod_{P \in D(\beta)} \mathcal{MMI}_\beta^P$$

scheme theoretically, and \mathcal{MI}_β can be identified with $\coprod_{P \in D(\beta)} |\mathcal{O}_X(\beta, P)|^\circ$ and \mathcal{MMI}_β can be considered as an open subscheme of the relative compactified Picard scheme over $\coprod_{P \in D(\beta)} |\mathcal{O}_X(\beta, P)|^\circ$.

- (3) If $h^0(\mathcal{O}_D) = 1$ and $P \in D_{\text{sm}}$, the following are equivalent:

- (a) The Abel map of D is immersive at P .
- (b) $h^0(D, \mathcal{O}_D(P)) = 1$.
- (c) ω_D has a global section nonzero at P .

In particular, the Abel map is immersive at any smooth point if D is integral with $p_a(D) > 0$ or $K_X + D \sim 0$.

For more about the immersivity of the Abel map of reduced Gorenstein curves we refer to [12].

Proof (1) Set-theoretically, this is obvious. Let $[C]$ be a point of \mathcal{MI}_β^P . By Lemma 2.1, the component of $\text{Hilb}(X)$ containing C can be identified with $[C]$. Taking a basis $\varphi_0, \dots, \varphi_d$ of $H^0(\mathcal{O}_X(C))$ with φ_0 corresponding to C , $[C]$ has natural local coordinates (s_1, \dots, s_d) near the point $[C]$ and \mathcal{MI}_β^P is defined by the vanishing of $\varphi_0 + s_1\varphi_1 + \dots + s_d\varphi_d$ on wP as a section of $\mathcal{O}_{wP}(C)$. This gives linear equations on s_1, \dots, s_d , and \mathcal{MI}_β^P can be scheme-theoretically identified with $|\mathcal{O}_X(\beta, P)|^\circ$.

(2) Let $[C]$ be a point of \mathcal{MI}_β with $P = C \cap D$ and \mathcal{Z} the family corresponding to a small neighborhood T of $[C]$. By Lemma 2.1, there is a morphism $T \rightarrow \mathbb{P}(H^0(\mathcal{O}_X(C)))$ for which \mathcal{Z} is the pullback of the universal curve \mathcal{C} . Taking the pullback of the universal curve by $U := H^0(\mathcal{O}_X(C)) \setminus \{0\} \rightarrow \mathbb{P}(H^0(\mathcal{O}_X(C)))$, we have a universal section of $\mathcal{O}_{X \times U}(C \times U)$ defining the family of curves \mathcal{C}_U and hence an isomorphism $\mathcal{O}_{X \times U}(\mathcal{C}_U - (C \times U)) \cong \mathcal{O}_{X \times U}$. By taking the pullback by a local lift $T \rightarrow U$ and restricting, we have $\mathcal{O}_{D \times T}(w \cdot \sigma(T) - w(P \times T)) \cong \mathcal{O}_{D \times T}$, where σ is as in (b) of (*).

Let $u : D_{\text{sm}} \rightarrow \text{Pic}(D)$ be the “Abel morphism”, defined roughly by $Q \mapsto \mathcal{O}_D(Q)$, and $[w] : \text{Pic}(D) \rightarrow \text{Pic}(D)$ the multiplication-by- w morphism. Then the above isomorphism shows that $[w] \circ u \circ \sigma$ is a constant map, where we regard σ as a morphism $T \rightarrow D$.

From the assumption that u is immersive at P and the étaleness of $[w]$, we see that σ is the constant map with value P . Thus the family $\mathcal{Z} \rightarrow T$ belongs to \mathcal{MI}_β^P . Since $\mathcal{MMI}_\beta \rightarrow D$ factors through \mathcal{MI}_β , the assertion on \mathcal{MMI}_β also holds.

- (3) The exact sequence $0 \rightarrow \mathcal{O}_D \rightarrow \mathcal{O}_D(P) \rightarrow \mathcal{O}_P(P) \rightarrow 0$ induces

$$0 \rightarrow H^0(\mathcal{O}_D) \xrightarrow{f} H^0(\mathcal{O}_D(P)) \rightarrow T_P D \xrightarrow{d_P u} T_{[\mathcal{O}_D(P)]} \text{Pic}(D),$$

and so (a) and (b) are equivalent.

From the exact sequence $0 \rightarrow \omega_D(-P) \rightarrow \omega_D \rightarrow \omega_D|_P \rightarrow 0$ we have a long exact sequence

$$H^0(\omega_D) \rightarrow H^0(\omega_D|_P) \rightarrow H^1(\omega_D(-P)) \xrightarrow{g} H^1(\omega_D),$$

and g is the Serre dual to f . Thus (c) is also equivalent.

If D is integral and $p_a(D) > 0$, $|P|$ consists of one point since otherwise it would give an isomorphism $D \cong \mathbb{P}^1$, so (b) holds.

If $K_X + D \sim 0$, then (c) is obvious. For an anticanonical curve $D(\neq 0)$ on a rational surface X the fact that $h^0(\mathcal{O}_D) = 1$ (and $h^1(\mathcal{O}_D) = 1$) is standard: This follows from the long exact sequence associated to $0 \rightarrow \mathcal{O}_X(-D) \cong \mathcal{O}_X(K_X) \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_D \rightarrow 0$, using Serre duality. \square

Recall that, if a class in $\text{Ext}_{\mathcal{O}_X}^1(F, F)$ is represented by an extension $0 \rightarrow F \xrightarrow{\alpha} \tilde{F} \xrightarrow{\beta} F \rightarrow 0$, then the corresponding deformation over $\mathbb{C}[\varepsilon]/\varepsilon^2$ is given by \tilde{F} with the action of ε on \tilde{F} defined as $\alpha \circ \beta$.

Lemma 2.9 *Let $P \in D_{\text{sm}}$ be a point, x a local parameter on D at P , and $F = \mathcal{O}_{D,P}/(x^w)$. Then a first order deformation of F as a coherent sheaf on D is given by $(\mathcal{O}_{D,P} \otimes \mathbb{C}[\varepsilon])/(x^w - g(x)\varepsilon, \varepsilon^2)$ for a unique polynomial $g \in \mathbb{C}[X]$ of degree $\leq w-1$. The corresponding extension is isomorphic to*

$$0 \rightarrow \mathcal{O}_{D,P}/(x^w) \xrightarrow{\alpha} (\mathcal{O}_{D,P} \otimes \mathbb{C}[\varepsilon])/((x^w - g(x)\varepsilon, \varepsilon^2)) \xrightarrow{\beta} \mathcal{O}_{D,P}/(x^w) \rightarrow 0,$$

where $\alpha(\tilde{f}) = \overline{f\varepsilon}$ and β is the reduction modulo ε .

Proof As in the proof of Lemma 2.7, a small deformation of F is equivalent to the deformation of the supporting scheme, and we may replace $\mathcal{O}_{D,P}$ by $\mathbb{C}[[x]]$. Then the assertion follows from the description of Hilbert schemes of points on a smooth curve as symmetric powers. \square

Remark 2.10 In the following, we describe the tangent spaces of \mathcal{MMI}_β and \mathcal{MMI}_β^P as the images of the natural maps $\text{Ext}_{\mathcal{O}_X}^1(F, F(-(w-1)P)) \rightarrow \text{Ext}_{\mathcal{O}_X}^1(F, F)$ and $\text{Ext}_{\mathcal{O}_X}^1(F, F(-D)) \rightarrow \text{Ext}_{\mathcal{O}_X}^1(F, F)$, respectively. In the former, we allow the intersection point to move along D . In the case of our main concern, the intersection point does not move in D by Lemma 2.8, and the tangent spaces coincide. We use $\text{Ext}_{\mathcal{O}_X}^1(F, F(-D))$ to prove our main result here, but a similar proof using $\text{Ext}_{\mathcal{O}_X}^1(F, F(-(w-1)P))$ is also possible.

Lemma 2.11 *Let $[F]$ be a point of \mathcal{MMI}_β^P for $P \in D_{\text{sm}}$.*

- (1) *The tangent space of \mathcal{MMI}_β^P at $[F]$ is naturally isomorphic to the image of the natural map $\text{Ext}_{\mathcal{O}_X}^1(F, F(-D)) \rightarrow \text{Ext}_{\mathcal{O}_X}^1(F, F)$.*
- (2) *Define $F(-(w-1)P) := \text{Ker}(F \rightarrow F|_{(w-1)P})$, where $(w-1)P$ is the closed subscheme of D defined by $(\mathcal{I}_{P \subset D})^{w-1}$.*

Then the tangent space of \mathcal{MMI}_β at $[F]$ is naturally isomorphic to the image of the natural map

$$\text{Ext}_{\mathcal{O}_X}^1(F, F(-(w-1)P)) \rightarrow \text{Ext}_{\mathcal{O}_X}^1(F, F).$$

Note that, if the supporting curve C of F is smooth at P , $F(-(w-1)P)$ can also be described as $F \otimes_{\mathcal{O}_C} \mathcal{O}_C(-(w-1)P)$.

Proof We begin by observing that $\text{Tor}_1^{\mathcal{O}_X}(F, \mathcal{O}_D) = 0$. This follows from the exact sequence

$$0 \rightarrow \text{Tor}_1^{\mathcal{O}_X}(F, \mathcal{O}_D) \rightarrow F(-D) \rightarrow F,$$

since the final map is immediately seen to be injective by the local form of F near D . We will use this vanishing without comment in the remainder of the proof to conclude that short exact sequences ending in F remain exact after restriction to D .

- (1) Take a tangent vector of \mathcal{MMI}_β^P at $[F]$ and let $0 \rightarrow F \rightarrow \tilde{F} \rightarrow F \rightarrow 0$ be the corresponding extension.

The restriction $0 \rightarrow F|_D \rightarrow \tilde{F}|_D \rightarrow F|_D \rightarrow 0$ is a split extension, so let $\tilde{G} \subset \tilde{F}|_D$ be the image of a splitting. Then we have a commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc}
& & 0 & & & & \\
& & \downarrow & & & & \\
& & \tilde{G} & \xrightarrow{\sim} & F|_D & & \\
& & \downarrow & & \parallel & & \\
0 \longrightarrow & F|_D & \longrightarrow & \tilde{F}|_D & \longrightarrow & F|_D & \longrightarrow 0 \\
\parallel & \downarrow & & \downarrow & & \parallel & \\
F|_D & \xrightarrow{\sim} & (\tilde{F}|_D)/\tilde{G} & & & & \\
\downarrow & & & & & & \\
& & 0. & & & &
\end{array}$$

The sheaf $F(-D)$ is the kernel of $F \rightarrow F|_D$, so if we write \tilde{F}' for the inverse image of \tilde{G} in \tilde{F} , we have a commutative diagram with exact rows:

$$\begin{array}{ccccccc}
0 \longrightarrow & F(-D) & \longrightarrow & \tilde{F}' & \longrightarrow & F & \longrightarrow 0 \\
\downarrow & & \downarrow & & \downarrow & & \\
0 \longrightarrow & F & \longrightarrow & \tilde{F} & \longrightarrow & F & \longrightarrow 0,
\end{array}$$

where the top row is the kernel of the natural surjection from the bottom row to the bottom row of the preceding commutative diagram. Thus our extension comes from a class in $\text{Ext}_{\mathcal{O}_X}^1(F, F(-D))$.

Conversely, if we are given an element of $\text{Ext}_{\mathcal{O}_X}^1(F, F(-D))$, let $0 \rightarrow F(-D) \rightarrow \tilde{F}' \rightarrow F \rightarrow 0$ be the corresponding extension. By push-out, we obtain a sheaf \tilde{F} and a commutative diagram with exact rows as above, where the lower row represents the induced class in $\text{Ext}_{\mathcal{O}_X}^1(F, F)$. By restricting to D , we have a commutative diagram with exact rows:

$$\begin{array}{ccccccc}
0 \longrightarrow & F(-D)|_D & \longrightarrow & \tilde{F}'|_D & \longrightarrow & F|_D & \longrightarrow 0 \\
i \downarrow & & j \downarrow & & \downarrow & & \\
0 \longrightarrow & F|_D & \longrightarrow & \tilde{F}|_D & \longrightarrow & F|_D & \longrightarrow 0.
\end{array}$$

Here i is 0, and therefore the induced map $\text{Im}(j) \rightarrow F|_D$ is an isomorphism. Thus the lower row is split, and \tilde{F} gives a tangent vector to \mathcal{MMI}_β^P .

(2) Let $0 \rightarrow F \rightarrow \tilde{F} \rightarrow F \rightarrow 0$ be an extension corresponding to a tangent vector of \mathcal{MMI}_β at $[F]$. By restriction, we have $0 \rightarrow F|_D \rightarrow \tilde{F}|_D \rightarrow F|_D \rightarrow 0$ satisfying $F|_D \cong \mathcal{O}_{wP}$.

We take a local parameter x of D at P , and identify finite-length modules over $\mathcal{O}_{D,P}$ with those on $\mathbb{C}[[x]]$.

The section σ in the definition of \mathcal{MMI}_β can be written as $x = a\varepsilon$, $a \in \mathbb{C}$. Then, with $c = wa$, we have $\tilde{F}|_D \cong \mathbb{C}[[x, \varepsilon]]/(x^w - cx^{w-1}\varepsilon)$. Here the map $F|_D \rightarrow \tilde{F}|_D$ is the map induced from the map $e : \tilde{F}|_D \rightarrow \tilde{F}|_D ; s \mapsto \varepsilon s$, where $(\tilde{F}|_D)/\ker(e)$ is identified with $F|_D$. Explicitly, it is given by $f(x) \pmod{(x^w)} \mapsto f(x)\varepsilon \pmod{(x^w - cx^{w-1}\varepsilon, \varepsilon^2)}$.

Write $G := \mathcal{I}_P^{w-1} \cdot (F|_D) \cong (x^{w-1})/(x^w)$ and let \tilde{G} be the \mathbb{C} -subspace of $\mathbb{C}[[x, \varepsilon]]/(x^w - cx^{w-1}\varepsilon, \varepsilon^2)$ with a basis $\bar{1}, \bar{x}, \dots, \bar{x}^{w-1}, \bar{x}^{w-1}\varepsilon$. Then the latter is also a $\mathbb{C}[[x]]$ -submodule, since $x \cdot \bar{x}^{w-1} = \bar{x}^w = cx^{w-1}\varepsilon$ and

$x \cdot \overline{x^{w-1}\varepsilon} = \overline{x^w\varepsilon} = \overline{cx^{w-1}\varepsilon^2} = 0$, and they fit in a commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \downarrow & & \downarrow & & \\
0 & \longrightarrow & G & \longrightarrow & \tilde{G} & \longrightarrow & F|_D \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \parallel \\
0 & \longrightarrow & F|_D & \longrightarrow & \tilde{F}|_D & \longrightarrow & F|_D \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \\
& & F|_{(w-1)P} & \xrightarrow{\sim} & (\tilde{F}|_D)/\tilde{G} & & \\
& & \downarrow & & \downarrow & & \\
& & 0 & & 0. & &
\end{array}$$

The inverse image of G in F is exactly $F(-(w-1)P)$. If we write \tilde{F}' for the inverse image of \tilde{G} in \tilde{F} , we have a commutative diagram with exact rows:

$$\begin{array}{ccccccc}
0 & \longrightarrow & F(-(w-1)P) & \longrightarrow & \tilde{F}' & \longrightarrow & F \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & F & \longrightarrow & \tilde{F} & \longrightarrow & F \longrightarrow 0,
\end{array}$$

and \tilde{F} comes from $\text{Ext}_{\mathcal{O}_X}^1(F, F(-(w-1)P))$.

Conversely, given an element of $\text{Ext}_{\mathcal{O}_X}^1(F, F(-(w-1)P))$, let $0 \rightarrow F(-(w-1)P) \rightarrow \tilde{F}' \rightarrow F \rightarrow 0$ be the corresponding extension. We obtain a sheaf \tilde{F} by push-out, and a commutative diagram with exact rows as above, the lower row representing the induced class in $\text{Ext}_{\mathcal{O}_X}^1(F, F)$. Restriction to D gives a commutative diagram with exact rows:

$$\begin{array}{ccccccc}
0 & \longrightarrow & F(-(w-1)P)|_D & \longrightarrow & \tilde{F}'|_D & \longrightarrow & F|_D \longrightarrow 0 \\
& & i \downarrow & & j \downarrow & & \downarrow \\
0 & \longrightarrow & F|_D & \longrightarrow & \tilde{F}|_D & \longrightarrow & F|_D \longrightarrow 0.
\end{array}$$

This induces an exact sequence $0 \rightarrow \text{Im}(i) \rightarrow \text{Im}(j) \rightarrow F|_D \rightarrow 0$, and we see by a local calculation that $\text{Im}(i)$ is of length 1 and is annihilated by x .

Let us identify the lower row with the exact sequence in the Lemma 2.9. We have to show that $g(x) = cx^{w-1}$ for some $c \in \mathbb{C}$. Since $\text{Im}(j)$ maps surjectively to $F|_D$, it contains an element of the form $\overline{1+h(x)\varepsilon}$, and we have $x^w \cdot \overline{1+h(x)\varepsilon} = \overline{g(x)\varepsilon} \in \text{Im}(j)$. Since it is mapped to $0 \in F|_D$, we have $\overline{g(x)\varepsilon} \in \text{Im}(\alpha \circ i)$. By the remark in the previous paragraph, we have $\overline{g(x)\varepsilon} = 0$ in $\mathbb{C}[[x, \varepsilon]]/(x^w - g(x)\varepsilon)$. If we write $g(x) = \sum_{i=0}^{w-1} c_i x^i$, this amounts to $\sum_{i=0}^{w-2} c_i x^{i+1}\varepsilon = 0$. Since $\overline{1}, \overline{x}, \dots, \overline{x^{w-1}}, \overline{\varepsilon}, \overline{x\varepsilon}, \dots, \overline{x^{w-1}\varepsilon}$ form a \mathbb{C} -basis of $\mathbb{C}[[x, \varepsilon]]/(x^w - g(x)\varepsilon, \varepsilon^2)$, we have $c_0 = \dots = c_{w-2} = 0$, which shows our assertion. \square

Theorem 2.12 (=Theorem 1.11) *Let X be a smooth projective rational surface, D an anticanonical curve on X and $P \in D_{\text{sm}}$. Then \mathcal{MMI}_β and \mathcal{MMI}_β^P are nonsingular of dimension $2p_a(\beta) = \beta^2 - w + 2$.*

Consequently, the relative compactified Picard scheme over $|\mathcal{O}_X(\beta, P)|^\circ$ is nonsingular at a point $[F]$ over $[C]$ if F is an invertible \mathcal{O}_C -module near P (or, equivalently, $F|_D \cong \mathcal{O}_C|_D$).

Proof Let $[F]$ be a point of \mathcal{MMI}_β^P with support curve C . Since $|\beta|$ contains C , which is integral by the definition of \mathcal{MMI}_β^P , the dimension of \mathcal{MI}_β^P is $p_a(C)$ by the remark after [16, Definition 4.16]. (Note that [16, Proposition

4.15] holds for a rational surface S , $E \in |-K_S|$ possibly reducible or non-reduced, β a curve class containing C with $h^0(\mathcal{O}_C) = 1$ and $P \in E(\beta)$.) By Theorem 2.4 and Lemma 2.7(3), \mathcal{MMI}_{β}^P is of dimension $2p_a(C)$ at $[F]$. Thus it suffices to show that the dimension of the tangent space at $[F]$ is $2p_a(C)$.

To see this, we note that the tangent space is the image of $\text{Ext}_{\mathcal{O}_X}^1(F, F(-D)) \rightarrow \text{Ext}_{\mathcal{O}_X}^1(F, F)$ by the previous lemma. Consider the natural exact sequence

$$\begin{aligned} 0 \rightarrow \text{Hom}_{\mathcal{O}_X}(F, F(-D)) &\rightarrow \text{Hom}_{\mathcal{O}_X}(F, F) \rightarrow \text{Hom}_{\mathcal{O}_X}(F, F|_{wP}) \\ &\xrightarrow{\delta} \text{Ext}_{\mathcal{O}_X}^1(F, F(-D)) \rightarrow \text{Ext}_{\mathcal{O}_X}^1(F, F). \end{aligned}$$

The first term is 0, the second term is \mathbb{C} , and the third is of dimension w . Thus the rank of δ is $w - 1$, and we have only to show that $\text{Ext}_{\mathcal{O}_X}^1(F, F(-D))$ has dimension $\beta^2 + 1$.

Riemann-Roch theorem tells us that $\sum_{i=0}^2 (-1)^i \dim \text{Ext}_{\mathcal{O}_X}^i(F, F(-D)) = -\beta^2$, and we already have $\text{Hom}_{\mathcal{O}_X}(F, F(-D)) = 0$. Since $-D \sim K_X$, we have $\text{Ext}_{\mathcal{O}_X}^2(F, F(-D)) \cong \text{Ext}_{\mathcal{O}_X}^2(F, F \otimes \mathcal{O}_X(K_X))$. This is dual to $\text{Hom}_{\mathcal{O}_X}(F, F)$, which is \mathbb{C} , thus yielding the assertion.

The second assertion follows from Lemma 2.3, 2.7 and 2.8. \square

Remark 2.13 This theorem can be seen as a partial logarithmic analogue of the unobstructedness of sheaves on K3 surfaces [35].

It might also be possible to think of the discreteness of $D(\beta)$ as analogous to the obstructedness of divisor classes in the moduli space of Kähler K3 surfaces.

3 Basic stable log maps

In this section, we recall some definitions and facts about basic stable log maps from [24, 13, 1] and [16, Section 2]. We restrict to maps of genus 0 with 1 distinguished marked point, target varieties endowed with log structure coming from a divisor. The general practice in log geometry is to underline log schemes to denote the underlying scheme. We only follow this convention when we wish to emphasize the distinction. See [20] for an introduction to log geometry.

Let X be a smooth variety and D a divisor on X . We will consider a stable log map f maximally tangent to D at a smooth point of D , and study the local structure of the moduli space at $[f]$. This depends only on a neighborhood of $\text{Im } f$, so we may assume that D is smooth and connected. We view X as the log scheme (X, \mathcal{M}_X) endowed with the divisorial log structure associated to D . Let $\beta \in H_2(X, \mathbb{Z})$ be a curve class.

3.1 Basic 1-marked genus 0 stable log maps to a smooth pair

For a log scheme (X, \mathcal{M}_X) , the monoid homomorphism $\mathcal{M}_X \rightarrow \mathcal{O}_X$ will be denoted by α .

Definition 3.1 Let $(\mathcal{C}/W, \{x_1\})$ be a 1-marked pre-stable log curve (in the sense of [24, Def. 1.3]) over a log scheme W and $(\mathcal{C}/W, \{x_1\}, f)$ a stable log map to X over W (i.e., $f : \mathcal{C} \rightarrow X$ is a log morphism

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{f} & X \\ \downarrow & & \downarrow \\ W & \longrightarrow & \text{Spec } \mathbb{C} \end{array}$$

and f is a stable map over W , see [24, Def. 1.6]).

It is called a stable log map of *maximal tangency* of genus 0 and class β if the following hold for any geometric point \bar{w} of \underline{W} :

(i) $\mathcal{C}_{\bar{w}}$ is of arithmetic genus 0 and $(f_{\bar{w}})_*[C_{\bar{w}}] = \beta$.

(ii) $f(x_1(\bar{w})) \in D$, and the natural map

$$\mathbb{N} \cong \Gamma(X, \overline{\mathcal{M}}_X) \longrightarrow \overline{\mathcal{M}}_{C_{\bar{w}}, x_1(\bar{w})} \rightarrow \overline{\mathcal{M}}_{C_{\bar{w}}, x_1(\bar{w})}/Q \cong \mathbb{N}$$

is given by $1 \mapsto D.\beta$.

In many cases, the condition (ii) follows from the rest of the conditions ([16, Proposition 2.9], Proposition 4.1).

Remark 3.2 (1) In the language of [24, Def. 3.1], this is the case $g = 0$, $k = 1$, the data “A” provided by β , $Z_1 = D$ and $s_1 \in \Gamma(D, (\overline{\mathcal{M}}_D^{\text{gp}})^*)$ given by $\overline{\mathcal{M}}_D^{\text{gp}} \simeq \mathbb{Z}_D \rightarrow \mathbb{Z}_D$, $1 \mapsto D.\beta$.

(2) The log structure on X is defined in the Zariski topology. Also, we work with genus 0 domain curves, and the base scheme \underline{W} will mainly be the spectrum of a finite dimensional local algebra over an algebraically closed field containing \mathbb{C} , and in that case it suffices to consider log structures in the Zariski topology.

Now we recall the definition of basicness (minimality) for stable log maps in the case of our concern, i.e. genus 0 stable log maps of maximal tangency to a log scheme associated to a smooth pair. See [24, §§ 1.4] for the general case.

Let $\underline{W} = \text{Spec } \kappa$ for an algebraically closed field $\kappa \supseteq \mathbb{C}$ and let $(\underline{C}/\underline{W}, \{x_1\}, f)$ be the 1-marked genus 0 stable map of class β that underlies a maximally tangent stable log map. Let $\pi : \underline{C} \rightarrow \underline{W}$ denote the structure morphism. Denote the nodes of \underline{C} by R_1, \dots, R_k , the irreducible components by C_0, \dots, C_k , and their generic points by η_0, \dots, η_k . We choose maps $g_1, g_2 : \{1, \dots, k\} \rightarrow \{0, \dots, k\}$ such that $R_i = C_{g_1(i)} \cap C_{g_2(i)}$.

We consider data $(Q, \overline{\mathcal{M}}_C, \psi, \varphi)$, where

- Q is a fine saturated (fs) monoid without invertible elements, regarded as a sheaf on \underline{W} ,
- $\overline{\mathcal{M}}_C$ is a fs sheaf of monoids on \underline{C} , and
- $\psi : \pi^{-1}Q \rightarrow \overline{\mathcal{M}}_C$, $\varphi : f^{-1}\overline{\mathcal{M}}_X \rightarrow \overline{\mathcal{M}}_C$ are local homomorphisms (i.e. non-invertible elements are mapped to non-invertible elements).

Here $\overline{\mathcal{M}}_X$ is the ghost sheaf of the log structure on X , but $\overline{\mathcal{M}}_C$ is a priori just a general sheaf of monoids satisfying the conditions above.

We require that $(Q, \overline{\mathcal{M}}_C, \psi)$ endows \underline{C} with the structure of a pre-stable log curve on the level of ghost sheaves, i.e.,

- ψ is an isomorphism on $\underline{C} \setminus \{x_1, R_1, \dots, R_k\}$,
- there is an isomorphism $\overline{\mathcal{M}}_{C, x_1} \rightarrow Q \oplus \mathbb{N}$ compatible with ψ (which is unique), and
- for each i , there is an element $\rho_{R_i} \in Q \setminus \{0\}$ such that, if we consider the homomorphism $\mathbb{N} \rightarrow Q$; $1 \mapsto \rho_{R_i}$ and the diagonal map $\mathbb{N} \rightarrow \mathbb{N}^2$ and take the pushout, there is an isomorphism $\overline{\mathcal{M}}_{C, R_i} \cong Q \oplus_{\mathbb{N}} \mathbb{N}^2$ characterized by the condition that the generization map $\overline{\mathcal{M}}_{C, R_i} \rightarrow \overline{\mathcal{M}}_{C, \eta_{g_1(i)}} \cong Q$ is given by $(q, (a_1, a_2)) \mapsto q + a_j \cdot \rho_{R_i}$.

In fact, these conditions are satisfied for the ghost sheaves and their homomorphisms obtained from a stable log map.

Assume that $f(x_1) \in D$. We may also assume that $\{i \mid f(R_i) \in D\} = \{1, \dots, k'\}$ and $\{i \mid f(\eta_i) \in D\} = \{0, \dots, k''\}$ after renumbering.

Definition 3.3 The type of $(Q, \overline{\mathcal{M}}_C, \psi, \varphi)$ as above is the element

$$\mathbf{u} = (u_{x_1}, u_{R_1}, \dots, u_{R_{k'}}) \in \mathbb{N} \times \mathbb{Z}^{k'}$$

defined as follows.

- $\mathbb{N} \ni u_{x_1}$ is the image of 1 by the map

$$\mathbb{N} \cong (f^{-1}\overline{\mathcal{M}}_X)_{x_1} \xrightarrow{\varphi_{x_1}} \overline{\mathcal{M}}_{C,x_1} \longrightarrow \overline{\mathcal{M}}_{C,x_1}/Q \cong \mathbb{N}.$$

- $\mathbb{Z} \ni u_{R_i}$ is characterized by the equality

$$u_{R_i} \cdot \rho_{R_i} = \varphi_{\eta_{g_2(i)}}(\chi_{i,2}) - \varphi_{\eta_{g_1(i)}}(\chi_{i,1}) \in Q,$$

where $\chi_{i,j} = 1 \in \mathbb{N} \cong (f^{-1}\overline{\mathcal{M}}_X)_{\eta_{g_j(i)}}$ if $g_j(i) \leq k''$ (i.e. $f(\eta_{g_j(i)}) \in D$), and $\chi_{i,j} = 0$ otherwise.

Given a stable map f as above and a type \mathbf{u} , we define a monoid Q^{basic} . We regard $\mathbb{Z}^{k'} \times \mathbb{Z}^{k''+1}$ as the direct product of \mathbb{Z} , one copy for each of $R_1, \dots, R_{k'}$ and $\eta_0, \dots, \eta_{k''}$. In this additive group, let a_{R_i} be the vector whose components are u_{R_i} at R_i , 1 at $\eta_{g_i(1)}$ if $g_i(1) \leq k''$, -1 at $\eta_{g_i(2)}$ if $g_i(2) \leq k''$, and 0 elsewhere. Then let \mathfrak{R} be the saturation of the subgroup generated by $a_{R_1}, \dots, a_{R_{k'}}$. We define Q^{basic} to be the saturation of the image of the natural map

$$\mathbb{N}^{k'} \times \mathbb{N}^{k''+1} \rightarrow (\mathbb{Z}^{k'} \times \mathbb{Z}^{k''+1})/\mathfrak{R}.$$

If we change the numbering of R_i , η_i and the maps g_i in a compatible way, there is a canonical homomorphism between the corresponding monoids. For example, exchanging $g_1(i)$ and $g_2(i)$ reverses the sign of u_{R_i} , hence that of a_{R_i} and thus \mathfrak{R} does not change.

Definition 3.4 ([24, Definition 1.20, Proposition 1.19]) Let $(C/W, \{x_1\}, f)$ be a stable log map over an fs log point $W = (\text{Spec } \kappa, \mathcal{M}_W)$ and let \mathbf{u} denote its type. Then it is called basic if the induced homomorphism $f^{-1}\overline{\mathcal{M}}_X \rightarrow \overline{\mathcal{M}}_C$ is universal in the category of maps of the ghost sheaves of type \mathbf{u} from $f^{-1}\overline{\mathcal{M}}_X$. This is equivalent to the condition that the natural homomorphism $Q^{\text{basic}} \rightarrow \overline{\mathcal{M}}_W$, induced by $(\rho_{R_1}, \dots, \rho_{R_{k'}}, \varphi_{\eta_0}(1), \dots, \varphi_{\eta_{k''}}(1))$, is an isomorphism.

A stable log map $(C/W, \{x_1\}, f)$ over a general fs log scheme W is called basic if, for any geometric point $\bar{w} \rightarrow W$, the induced log map over \bar{w} is basic.

3.2 Moduli


By [24, Theorem 0.1] the stack of basic stable log maps to X is an algebraic log stack, which is locally of finite type. By [24, Theorem 0.2], imposing genus 0, class β and maximal tangency with D cuts out a proper Deligne-Mumford stack $\overline{\mathcal{M}}_\beta(X, D)$, which admits a virtual fundamental class, thus yielding the log Gromov-Witten invariant $\mathcal{N}_\beta(X, D)$ of maximal tangency of (X, D) . In fact, by [47, Corollary 1.2], the forgetful morphism from $\overline{\mathcal{M}}_\beta(X, D)$ to the moduli space of genus 0 stable maps of X of class β is finite. In what follows, we analyze special components of $\overline{\mathcal{M}}_\beta(X, D)$ corresponding to log maps whose images have two irreducible components.

The log structure at the nodes gives rise to tropical curves. We illustrate it by the example that will be relevant in the next section, cf. Corollary 4.2. Consider a chain of smooth rational curves C_1, C_0, C_2 meeting successively in nodes. We map it to (X, D) by collapsing C_0 to $P \in D$ and mapping C_1 and C_2 each to a rational curve in X meeting D only at P in maximal tangency. This gives a “tropical curve in \mathbb{R} ” as in Figure 1. All edges are parallel, weighted and satisfy the balancing condition, see Corollary 4.2. For more details, see also [16, Section 2].

4 Proof of Theorem 1.14

In this section, we prove Theorem 1.14.

Figure 1: Tropical curve



4.1 Maximally tangent genus 0 log maps with 2 non-collapsed components

The following gives a rough idea of what a genus 0, 1-marked stable log map looks like.

Proposition 4.1 ([16, Corollary 2.10]) *Let X be a divisorial log scheme given by a smooth variety X and a smooth divisor D .*

For a genus 0 stable log map $f : (C/W, x_1) \rightarrow X$ with $\underline{W} = \text{Spec } \mathbb{C}$, assume the following:

- $w := D.f_*[C] > 0$ and $d_i := D.f_*[C_i] \geq 0$ for any irreducible component C_i of C .
- If C_i is an irreducible component of C that is not collapsed by f , then $f(C_i) \not\subseteq D$.

Then it is of maximal tangency, and the following holds.

- (1) $f(C) \cap D$ consists of one point P .
- (2) If there is only 1 non-collapsed component, then $C \cong \mathbb{P}^1$ and $f^*(D) = w x_1$.
- (3) If there are at least 2 non-collapsed components, and $D.f_*[C_i] > 0$ holds for non-collapsed components, then C is given by adding $C_i = \mathbb{P}^1$ as leaves to a tree C' of \mathbb{P}^1 collapsed to P , with maps $f_i : C_i \rightarrow X$ satisfying $f_i^*(D) = d_i(C_i \cap C')$.

Now we consider the case where the image of f has 2 components. We employ the notation of §§ 3.1.

Corollary 4.2 *Let X be a divisorial log scheme given by a smooth variety X and a smooth divisor D .*

For a genus 0 stable log map $f : (C/W, x_1) \rightarrow X$ with $\underline{W} = \text{Spec } \mathbb{C}$, assume that C has 2 non-collapsed components C_1 and C_2 with $f(C_i) \not\subseteq D$, that $f|_{C_i}$ are birational, and that $f(C_1) \neq f(C_2)$.

Assume $d_i := D.f(C_i) > 0$ and let $d = \gcd(d_1, d_2)$ and $e_i = d_i/d$. Then the following holds.

- (1) C is a chain of smooth rational curves C_1, C_0 and C_2 in this order, C_0 mapping to a point $P \in D$, and we have $(f|_{C_i})^*D = d_i R_i$ for $i = 1, 2$, where $R_i = C_i \cap C_0$.

We think of C_0 as the “first” component at R_1, R_2 , i.e. we set $g_1(1) = 0$, $g_2(1) = 1$ and $g_1(2) = 0$, $g_2(2) = 2$.

- (2) With the notation in §§ 3.1, the homomorphism on the log structure at the ghost level is given by $\varphi_{R_1}(1) = \overline{(0, (d_1, 0))}$, $\varphi_{R_2}(1) = \overline{(0, (d_2, 0))}$ and $\varphi_{\eta_0}(1) = d_1 \cdot \rho_{R_1} = d_2 \cdot \rho_{R_2}$. The type of f is given by $\mathbf{u} = (u_{x_1}, u_{R_1}, u_{R_2}) = (d_1 + d_2, -d_1, -d_2)$.

- (3) The stable log map f is basic if and only if $\overline{\mathcal{M}}_W$ is isomorphic to \mathbb{N} and, identifying $\overline{\mathcal{M}}_W$ with \mathbb{N} , $\rho_{R_1} = e_2$, $\rho_{R_2} = e_1$ and $\varphi_{\eta_0}(1) = de_1e_2$. The last condition can be replaced by “either $\rho_{R_1} = e_2$, $\rho_{R_2} = e_1$ or $\varphi_{\eta_0}(1) = de_1e_2$.”

Proof (1) By the previous proposition, C is obtained by adding C_1 and C_2 as leaves to a tree of \mathbb{P}^1 's. This tree is collapsed and has only 3 special points apart from nodes, i.e. x_1 and the intersection points with C_1 and C_2 . From stability it must be \mathbb{P}^1 .

The latter half also follows from the previous proposition.

(2) Note that $k = k' = 2$ and $k'' = 0$. Let $\varphi_{R_1}(1)$ be $\overline{(q, (a, b))}$, and let w be a lift to \mathcal{M}_{C, R_1} . If z_1 and z_2 denote local coordinates near R_1 with $C_0 = (z_1 = 0)$ and $C_1 = (z_2 = 0)$, then w maps to a function that is $0^q z_1^a z_2^b$ up to unit, where 0^q is 1 if $q = 0$ and 0 otherwise. Comparing with the pullback of a local defining equation of D , we have $q = 0$, $a = d_1$ and $b = 0$. By generalization we have

$$\varphi_{\eta_{g_1(1)}}(\chi_{1,1}) = \varphi_{\eta_0}(1) = d_1 \cdot \rho_{R_1}.$$

We also have $\chi_{1,2} = 0$, so $u_{R_1} = -d_1$. We obtain $u_{R_2} = -d_2$ in the same way.

The equality $u_{x_1} = d_1 + d_2$ follows from the definition of a maximally tangent log map, or from the balancing condition([24, Proposition 1.15]).

(3) We have $a_{R_1} = (-d_1, 0, 1)$ and $a_{R_2} = (0, -d_2, 1)$. Thus \mathfrak{R} is generated by a_{R_1} and $(a_{R_1} - a_{R_2})/d = (-e_1, e_2, 0)$, and we have an isomorphism $\mathbb{Z}^3/R \rightarrow \mathbb{Z}; \overline{(a, b, c)} \mapsto ae_2 + be_1 + cde_1e_2$. This maps \mathbb{N}^3 to $\mathbb{N}e_1 + \mathbb{N}e_2$, and its saturation Q^{basic} is \mathbb{N} .

The natural homomorphism $Q^{\text{basic}} \rightarrow \overline{\mathcal{M}}_W$ is the one induced from

$$\mathbb{N}^3 \rightarrow \overline{\mathcal{M}}_W; (a, b, c) \mapsto a \cdot \rho_{R_1} + b \cdot \rho_{R_2} + c \cdot \varphi_{\eta_0}(1),$$

and therefore e_2 , e_1 and de_1e_2 are mapped to ρ_{R_1} , ρ_{R_2} and $\varphi_{\eta_0}(1)$, respectively. Thus we have the “only if” part. Conversely, if $\overline{\mathcal{M}}_W \cong \mathbb{N}$ and $\rho_{R_1} = e_2$, say, then $\mathbb{N} \cong Q^{\text{basic}} \rightarrow \overline{\mathcal{M}}_W \cong \mathbb{N}$ is a monoid homomorphism which maps e_2 to e_2 , hence the identity. \square

4.2 Coordinates and log structures

In the rest of this section, we make the following assumptions, which are part of those in Theorem 1.14 after taking a neighborhood of $Z_1 \cup Z_2$.

Assumptions 4.3 Let (X, D) be a pair of a smooth variety and a smooth divisor. Let Z_1 and Z_2 be proper integral curves in X satisfying the following:

- (1) Z_i is a rational curve of class β_i maximally tangent to D ,
- (3) $Z_1 \cap D$ and $Z_2 \cap D$ consist of the same point P , and
- (4') $Z_1 \neq Z_2$.

Write $d_i = de_i = D.Z_i$ with $\gcd(e_1, e_2) = 1$.

We will study the deformation space of stable log maps by an explicit calculation. As a preparation, let us fix coordinate systems.

In order to deal with the deformations of C , we make it stable by adding marked points. The moduli scheme $\mathbf{M}_{0,5}$ of 5-marked genus 0 stable curves has a point corresponding to our curve $C = C_1 \cup C_0 \cup C_2$, with markings $x_2, x_3 \in C_1$, $x_1 \in C_0$ and $x_4, x_5 \in C_2$.

Notation 4.4 As a formal scheme supported by 1 point, we have the following description for the formal neighborhood \mathbf{M} of $[(C, x_1, \dots, x_5)] \in \mathbf{M}_{0,5}$, the universal curve \mathcal{C} (which has the same underlying space as C) over \mathbf{M} and the marked sections $\tilde{x}_1, \dots, \tilde{x}_5$ extending x_1, \dots, x_5 .

(1) \mathbf{M} is a formal 2-disk with coordinates μ_1 and μ_2 .

(2) $\mathcal{C} = \mathcal{U}'_1 \cup \mathcal{U}_1 \cup \mathcal{U}_2 \cup \mathcal{U}'_2$, where

- $\mathcal{U}'_i = \mathbb{A}_{\mathbf{M}}^1 = \{(z_{i0}, \mu_1, \mu_2)\}$,
- $\mathcal{U}_i = \{(z_{i1}, z_{i2}, \mu_1, \mu_2) | z_{i1}z_{i2} = \mu_i\} \subset \mathbb{A}_{\mathbf{M}}^2$,
- $\{z_{i0} \neq 0\} \subset \mathcal{U}'_i$ and $\{z_{i1} \neq 0\} \subset \mathcal{U}_i$ are patched by $z_{i0} = 1/z_{i1}$, and
- $\{z_{i2} \neq 0\} \subset \mathcal{U}_1$ and $\{z_{22} \neq 0\} \subset \mathcal{U}_2$ are patched by $z_{12} = 1/z_{22}$ (and $z_{11} = \mu_1 z_{22}$, $z_{21} = \mu_2 z_{12}$).

We denote the point $(z_{i1} = z_{i2} = 0)$ by R_i .

(3) The marked sections are $\tilde{x}_1 : (z_{12} = 1)$ on \mathcal{U}_1 , $\tilde{x}_2 : (z_{10} = 0)$ on \mathcal{U}'_1 , $\tilde{x}_3 : (z_{11} = 1)$ on \mathcal{U}_1 , $\tilde{x}_4 : (z_{21} = 1)$ on \mathcal{U}_2 and $\tilde{x}_5 : (z_{20} = 0)$ on \mathcal{U}'_2 .

We denote the central fiber of \mathcal{U}_i by U_i , that of \tilde{x}_i by x_i , etc. For $i = 1$ or 2 , write C_i° for $C_i \setminus R_i$ and \mathcal{C}_i° for the corresponding open subspace of \mathcal{C} .

Lemma 4.5 Let $S = \text{Spec } R$ where R is a local finite-dimensional \mathbb{C} -algebra with a basis $\{r_1, \dots, r_n\}$, and \mathcal{C}_S the curve induced from \mathcal{C} by a morphism $S \rightarrow \mathbf{M}$.

Identifying the underlying topological spaces of \mathcal{C} and \mathcal{C}_S with that of C , let U be an open neighborhood of R_i in U_i .

Then any element F of $\mathcal{O}_{\mathcal{C}_S}(U)$ can be uniquely written as

$$F = \sum_{k=1}^n r_k \left\{ F_0^{(k)} + z_{i1} F_1^{(k)}(z_{i1}) + z_{i2} F_2^{(k)}(z_{i2}) \right\},$$

where $F_0^{(k)} \in \mathbb{C}$, $F_1^{(k)}(z_{i1}) \in \mathcal{O}_{C_i}(U \cap C_i)$ and $F_2^{(k)}(z_{i2}) \in \mathcal{O}_{C_0}(U \cap C_0)$. Here, if i is e.g. 1, we regard $\mathcal{O}_{C_1}(U \cap C_1)$ as a subring of $\mathcal{O}_{\mathcal{C}_S}(U)$ using the projection map $\mathcal{U}_1 \rightarrow \text{Spec } \mathbb{C}[z_{11}]$, etc.

Proof This basically follows from $z_{i1}z_{i2} = \mu_i$ and flatness (or an explicit calculation). \square

Notation 4.6 The standard log structures on \mathbf{M} and \mathcal{C} are the divisorial log structures defined by the degeneracy locus and its inverse image and the sections. We replace the log structure at $\tilde{x}_2, \dots, \tilde{x}_5$ with the one induced from \mathbf{M} . They are explicitly given as follows.

- $\mathcal{M}_{\mathbf{M},0} = \coprod_{a,b=0}^{\infty} \mu_1^a \mu_2^b \mathcal{O}_{\mathbf{M},0}^\times$, with $\alpha(\mu_1) = \mu_1$ and $\alpha(\mu_2) = \mu_2$.
- $\mathcal{M}_{\mathcal{C},R_1} = \coprod_{a,b,c=0}^{\infty} z_{11}^a z_{12}^b \mu_2^c \mathcal{O}_{\mathcal{C},R_1}^\times$ and $\mathcal{M}_{\mathcal{C},R_2} = \coprod_{a,b,c=0}^{\infty} z_{21}^a z_{22}^b \mu_1^c \mathcal{O}_{\mathcal{C},R_2}^\times$, with $\alpha(z_{ij}) = z_{ij}$.
- $\mathcal{M}_{\mathcal{C},x_1} = \coprod_{a,b,c=0}^{\infty} z'^a \mu_1^b \mu_2^c \mathcal{O}_{\mathcal{C},x_1}^\times$ with $\alpha(z') = z_{12} - 1$.
- At other points, $\mathcal{M}_{\mathcal{C}} = \coprod_{a,b=0}^{\infty} \mu_1^a \mu_2^b \mathcal{O}_{\mathcal{C}}^\times$.

The structure homomorphism and the generization maps are given by the following rule:

- μ_i at various points are identified.
- μ_i maps to $z_{i1}z_{i2} \in \mathcal{M}_{\mathcal{C},R_i}$.
- If $\{j, j'\} = \{1, 2\}$, z_{ij} and $z_{ij'}$ generizes to z_{ij} and $\mu_i z_{ij}^{-1}$ on $\{z_{ij} \neq 0\}$. z' generizes to $z_{12} - 1$.

The parameter spaces we will be concerned with are of the following form.

Notation 4.7 For a nonnegative integer n and $m(s) \in s \cdot \mathbb{C}[s]/(s^{n+1})$, let $S_{n,m(s)}$ be the log scheme $(\underline{S}, \mathcal{M})$ defined as follows.

- $\underline{S} = S_n = \text{Spec } \mathbb{C}[s]/(s^{n+1})$.

- $\mathcal{M} = \mathbb{N} \times \mathcal{O}_{\underline{S}}^{\times}$, with $\alpha(k, u(s)) = m(s)^k u(s)$.

For coprime positive integers r_1, r_2 and $u_1(s), u_2(s) \in \mathcal{O}_{\underline{S}}^{\times}$, let $\mathcal{C}_{n,m(s),(r_1,u_1(s)),(r_2,u_2(s))}$ denote the log scheme over $S_{n,m(s)}$ obtained from \mathcal{C} by taking the fiber product with the log morphism $S_{n,m(s)} \rightarrow \mathbf{M}$ defined by $\mu_i \mapsto (r_i, u_i(s))$ (and $\mu_i \mapsto m(s)^{r_i} u_i(s)$) in the category of log schemes.

Lemma 4.8 Write S for $S_{n,m(s)}$ and \mathcal{C}_S for $\mathcal{C}_{n,m(s),(r_1,u_1(s)),(r_2,u_2(s))}$. Then S and \mathcal{C}_S are fs log schemes and \mathcal{C}_S/S with the induced section is a 1-marked pre-stable log curve.

As a scheme, it is represented as in Notation 4.4 with μ_i replaced by $m(s)^{r_i} u_i(s)$.

The induced log structure can be described as follows. (We use the same symbols for points and functions on \mathcal{C} and \mathcal{C}_S etc.)

- $\mathcal{M}_{S,0} = \coprod_{a=0}^{\infty} \mathbf{m}^a \mathcal{O}_{S,0}^{\times}$, with $\alpha(\mathbf{m}) = m(s)$.
- $\mathcal{M}_{\mathcal{C}_S,R_i} = \bigcup_{a,b,c=0}^{\infty} \mathbf{z}_{i1}^a \mathbf{z}_{i2}^b \mathbf{m}^c \mathcal{O}_{\mathcal{C}_S,R_i}^{\times} = \coprod_{a,b=0}^{\infty} \coprod_{c=0}^{r_i-1} \mathbf{z}_{i1}^a \mathbf{z}_{i2}^b \mathbf{m}^c \mathcal{O}_{\mathcal{C}_S,R_i}^{\times}$ subject to the relation $\mathbf{z}_{i1} \mathbf{z}_{i2} = \mathbf{m}^{r_i} u_i(s)$, with $\alpha(\mathbf{z}_{ij}) = z_{ij}$.
- $\mathcal{M}_{\mathcal{C}_S,x_1} = \coprod_{a,b=0}^{\infty} \mathbf{z}'^a \mathbf{m}^b \mathcal{O}_{\mathcal{C}_S,x_1}^{\times}$ with $\alpha(\mathbf{z}') = z_{12} - 1$.
- At other points, $\mathcal{M}_{\mathcal{C}_S} = \coprod_{a=0}^{\infty} \mathbf{m}^a \mathcal{O}_{\mathcal{C}_S}^{\times}$.

The structure homomorphism and the generization maps are given by the following relations:

- \mathbf{m} at various places are to be identified.
- If $\{j, j'\} = \{1, 2\}$, \mathbf{z}_{ij} and $\mathbf{z}_{ij'}$ generizes to z_{ij} and $\mathbf{m}^{r_i} u_i(s) z_{ij}^{-1}$ on $\{z_{ij} \neq 0\}$. \mathbf{z}' generizes to $z_{12} - 1$.

Proof One thing that is not so obvious is that the fiber product in the category of log schemes is a fs log scheme in this case, but this is a known fact in the study of moduli of log curves. In fact, the ghost sheaf at R_1 is the pushout

$$\overline{\mathcal{M}}_S \oplus_{\overline{\mathcal{M}}_M} \overline{\mathcal{M}}_{\mathcal{C}, R_1} \cong \mathbb{N} \oplus_{\mathbb{N}^2} \mathbb{N}^3$$

defined from $\mathbb{N}^2 \rightarrow \mathbb{N}; (a, b) \mapsto r_1 a + r_2 b$ and $\mathbb{N}^2 \rightarrow \mathbb{N}^3; (a, b) \mapsto (a, a, b)$. This is isomorphic to the submonoid of \mathbb{N}^2 generated by $(r_1, 0)$, $(0, r_1)$ and $(1, 1)$ (corresponding to $\bar{\mathbf{z}}_{11}, \bar{\mathbf{z}}_{12}$ and $\bar{\mathbf{m}}$, respectively). \square

Notation 4.9 Let (w_1, w_2, \dots, w_N) be étale coordinates on an affine open neighborhood W of P in X such that $w_1 = 0$ defines D in W .

Let $f^{(i)} : \mathbb{P}^1 \rightarrow Z_i \subset X$ be the normalization map. We may assume that $\infty \notin (f^{(i)})^{-1}(W)$.

For $i = 1$ or 2 , we identify the domain of $f^{(i)}$ with C_i , and by extending to C_0 by a constant map, we obtain a stable map $f_0 : (C, x_1) \rightarrow X$. Denote $f_0^{-1}(W)$ by V , and then we have $C_0 \subset V$ (and in particular $R_1, R_2 \in V$) and $x_2, x_5 \notin V$. We define $A_i(z_{i1})$ and $B_{ji}(z_{i1})$, $i = 1, 2$, by

$$\begin{aligned} w_1 &\mapsto z_{i1}^{d_i} A_i(z_{i1}), & A_i(z_{i1}) &\in \mathcal{O}_{C_i}(V \cap C_i)^{\times}, \\ w_j &\mapsto B_{ji}(z_{i1}), & B_{ji}(z_{i1}) &\in \mathcal{O}_{C_i}(V \cap C_i), & 2 \leq j \leq N. \end{aligned}$$

By the assumptions, we have $B_{j1}(0) = B_{j2}(0)$. We write

$$\begin{aligned} V_0 &= U_1 \cap U_2, \\ V_1 &= V \cap (U_1 \setminus x_1), \\ V_2 &= V \cap (U_2 \setminus x_1), \end{aligned}$$

and denote the corresponding open subspaces of \mathcal{C} by \mathcal{V} and \mathcal{V}_i .

4.3 Calculation

Recall that $\overline{M}_\beta = \overline{M}_\beta(X, D)$ is the moduli stack of maximally tangent genus 0 basic stable log maps of class β to the log scheme associated to (X, D) .

We consider $S = S_n := \text{Spec } \mathbb{C}[s]/(s^{n+1})$. This will be sufficient since the tangent space to \overline{M}_β is at most 1-dimensional as we will see later. (In fact, most of what we state below generalize to the spectrum of any local finite-dimensional \mathbb{C} -algebra.)

When referring to a base change of something to S , we sometimes drop S , e.g. we write just \mathcal{V}_i for $(\mathcal{V}_i)_S$.

Lemma 4.10 *Let $(\mathcal{C}_S/S, \{\tilde{x}_1\}, f)$ be a basic stable log map over S such that the cycle theoretic image of the central fiber is $Z_1 + Z_2$.*

(1) *One can identify the restriction $\underline{f}|_0$ to the central fiber with f_0 constructed in Notation 4.9. There exist $m(s) = \sum_{k=1}^n m^{(k)} s^k$ and $u_i(s) = \sum_{k=0}^n u_i^{(k)} s^k$ such that \mathcal{C}_S/S is isomorphic to $\mathcal{C}_{n,m(s),(e_2,u_1(s)),(e_1,u_2(s))}/S_{n,m(s)}$. (Note that e_2 comes first.)*

(2) *Via an isomorphism as in (1), the log map f gives rise to*

$$\begin{aligned}\tilde{A}_i &\in \mathcal{O}_{\mathcal{C}_S}(\mathcal{V}_i)^\times, & i &= 1, 2 \\ \tilde{B}_{2i}, \dots, \tilde{B}_{Ni} &\in \mathcal{O}_{\mathcal{C}_S}(\mathcal{V}_i), & i &= 1, 2,\end{aligned}$$

with $z_{i1}^{d_i} \tilde{A}_i \equiv z_{i1}^{d_i} A_i \pmod{(s)}$ and $\tilde{B}_{ji} \equiv B_{ji} \pmod{(s)}$, characterized by the condition that

$$\begin{aligned}(f|_{\mathcal{V}_i})^\flat w_1 &= z_{i1}^{d_i} \tilde{A}_i, \\ (\underline{f}|_{\mathcal{V}_i})^* w_j &= \tilde{B}_{ji} \quad j = 2, \dots, N.\end{aligned}$$

Here, w_1 is w_1 considered as a local section of \mathcal{M}_X .

Conversely, such data uniquely determines f (under the conditions described in the next lemma).

Proof (1) By Corollary 4.2 (1), we can identify $\underline{f}|_0$ with f_0 . By Corollary 4.2 (3), $\mathcal{M}_{S,0} \cong \mathbb{N}$ and hence S is isomorphic to $S_{n,m(s)}$ for some $m(s)$.

Since the central fiber has 3 components, by [26], (modulo adding 4 sections and then subtracting), \mathcal{C}_S is induced from some log morphism $S \rightarrow \mathbf{M}$. By Corollary 4.2 (3), μ_1 maps to e_2 and μ_2 maps to e_1 at the level of ghost sheaves. Thus $S \rightarrow \mathbf{M}$ is of the asserted form.

(2) By Corollary 4.2 (2), $(f|_{\mathcal{V}_i})^\flat w_1$ is of the form as above.

For the converse: By the compatibility of f^* and f^\flat , we have $(\underline{f}|_{\mathcal{V}_i})^* w_1 = z_{i1}^{d_i} \tilde{A}_i$. Since w_1, \dots, w_N are étale coordinates on W , this determines $f|_{\mathcal{V}_S}$. If $j : V \rightarrow C$ denotes the inclusion map, then $\mathcal{O}_{\mathcal{C}_S} \rightarrow j_* \mathcal{O}_{\mathcal{V}_S}$ is injective, and therefore $\tilde{A}_i, \tilde{B}_{2i}, \dots, \tilde{B}_{Ni}$ determine f . \square

By Lemma 4.5, we can expand \tilde{A}_i and \tilde{B}_i as

$$\begin{aligned}\tilde{A}_i &= \sum_{k=0}^n s^k \left\{ A_{i0}^{(k)} + z_{i1} A_{i1}^{(k)}(z_{i1}) + z_{i2} A_{i2}^{(k)}(z_{i2}) \right\} \\ \tilde{B}_{ji} &= \sum_{k=0}^n s^k \left\{ B_{ji0}^{(k)} + z_{j1} B_{j1}^{(k)}(z_{j1}) + z_{j2} B_{j2}^{(k)}(z_{j2}) \right\}\end{aligned}$$

with

$$\begin{aligned} A_{i0}^{(k)}, B_{j0}^{(k)} &\in \mathbb{C}, \\ A_{i1}^{(k)}(z_{i1}), B_{j1}^{(k)}(z_{i1}) &\in \mathcal{O}_{C_i}(V_i \cap C_i) = \mathcal{O}_{C_i}(V \cap C_i), \\ A_{i2}^{(k)}(z_{i2}), B_{j2}^{(k)}(z_{i2}) &\in \mathcal{O}_{C_0}(V_i \cap C_0) = \mathcal{O}_{C_0}((U_i \cap C_0) \setminus x_1). \end{aligned}$$

Let us write down the conditions for these data to actually give a log map.

Lemma 4.11 *The above data give a family in $\overline{\mathbf{M}}_\beta$ if and only if the following hold.*

- (a) (w_1 on \mathcal{V}_0) $z_{12}^{-d_1} u_1(s)^{d_1} \tilde{A}_1 = z_{22}^{-d_2} u_2(s)^{d_2} \tilde{A}_2$ on $\mathcal{V}_0 \setminus x_1$, and it extends to a function on \mathcal{V}_0 with vanishing order $d_1 + d_2$ along \tilde{x}_1 .
- (b) (w_2, \dots, w_N on \mathcal{V}_0) For $j = 2, \dots, N$, $\tilde{B}_{j1}|_{\mathcal{V}_0 \setminus x_1} = \tilde{B}_{j2}|_{\mathcal{V}_0 \setminus x_1}$, and it extends to a regular function at x_1 .
- (c) (w_1, \dots, w_N on C_1°, C_2°) The morphism $\mathcal{V} \cap C_i^\circ \rightarrow S$ given by $z_{i1}^{d_1} \tilde{A}_i$ and $\tilde{B}_{2i}, \dots, \tilde{B}_{Ni}$ extends to a morphism $C_i^\circ \rightarrow X$.

Proof We first show that the conditions are necessary:

For (a), on \mathcal{V}_0 we have $z_{11} = \mathbf{m}^{e_2} z_{12}^{-1} u_1(s)$ and $z_{21} = \mathbf{m}^{e_1} z_{22}^{-1} u_2(s)$, and so

$$\begin{aligned} z_{11}^{d_1} \tilde{A}_1 &= \mathbf{m}^{de_1 e_2} z_{12}^{-d_1} u_1(s)^{d_1} \tilde{A}_1, \\ z_{21}^{d_2} \tilde{A}_2 &= \mathbf{m}^{de_1 e_2} z_{22}^{-d_2} u_2(s)^{d_2} \tilde{A}_2. \end{aligned}$$

These must be equal, so we have the first assertion. At x_1 it is equal to $\mathbf{m}^{de_1 e_2} (\mathbf{z}')^{d_1 + d_2} \cdot (\text{unit})$, hence the second assertion. It is obvious that (b) and (c) must hold in order to define \underline{f} .

The conditions are also sufficient: From (a), by the equalities above we have $z_{11}^{d_1} \tilde{A}_1 = z_{21}^{d_2} \tilde{A}_2$ on $\mathcal{V}_0 \setminus x_1$, and it extends to x_1 . Together with (b) and (c), these data give a stable map \underline{f} . By (a), we can lift it to a log map by sending w_1 to $z_{11}^{d_1} \tilde{A}_1$, $z_{21}^{d_2} \tilde{A}_2$ and $\mathbf{m}^{de_1 e_2} (\mathbf{z}')^{d_1 + d_2} \cdot (\text{unit})$ at R_1, R_2 and x_1 , respectively. \square

Let c_0 be a d_2 -th roots of $(-1)^{d_1+d_2} A_1(0)/A_2(0)$ and set $c_p := e^{2\pi p\sqrt{-1}/d_2} c_0$. The following theorem proves Theorem 1.14.

Theorem 4.12 *Let (X, D) and Z_i be as in Assumption 4.3 and $f^{(i)}, A_i, B_{ji}$ be as in Notation 4.9. For $n < \min\{e_1, e_2\}$, let $S_{n,s}$ and $\mathcal{C}_{n,p} = \mathcal{C}_{n,s,(e_2,1),(e_1,c_p)}$ be as defined in Notation 4.7.*

- (1) *The underlying pre-stable curve of $\mathcal{C}_{n,p}/S_{n,s}$ is the trivial family.*
- (2) *The functions*

$$\begin{aligned} \tilde{A}_i &= A_i(z_{i1}) + A_i(0) \{(1 - z_{i2})^{d_1 + d_2} - 1\}, \\ \tilde{B}_{j,i} &= B_{j,i}(z_{i1}) \end{aligned}$$

defines a stable log map $f_{n,p} : \mathcal{C}_{n,p} \rightarrow X$, whose underlying stable map is the trivial family.

- (3) *The stable log maps $f_{0,p}$ and $f_{0,p'}$ are isomorphic if and only if $p \equiv p' \pmod{d}$. Each $f_{0,p}$ has no nontrivial automorphisms.*
- (4) *The stable log map $f_{n,p}$ gives a closed immersion of $S_{n,s}$ into $\overline{\mathbf{M}}_\beta$.*
- (5) *The stable log maps $f_{0,p}$ exhaust the maps whose image cycles are $Z_1 + Z_2$.*
- (6) *Under the assumptions of Theorem 1.14, the stable log map $f_{\min\{e_1, e_2\}-1, p}$ gives an isomorphism of $S_{\min\{e_1, e_2\}-1}$ with a connected component.*

Proof (5) and (6) will be proven in the subsections that follow.

(1) Since $e_1, e_2 > n$, we have $s^{e_1} = s^{e_2} = 0$ in $\mathbb{C}[s]/(s^{n+1})$. Thus the underlying pre-stable curve is trivial.

(2) We have $z_{i1}z_{i2} = 0$ from (1). Thus $z_{i1}^{d_i}\tilde{A}_i = z_{i1}^{d_i}A_i(z_{i1})$, and these data give a trivial stable map. In particular, the conditions (b) and (c) of Lemma 4.11 are clearly satisfied.

Let us check Lemma 4.11 (a). We have $u_1(s) = 1$ and $u_2(s) = c_p$ and therefore, on \mathcal{V}_0 ,

$$\begin{aligned} z_{12}^{-d_1}u_1(s)^{d_1}\tilde{A}_1 &= z_{12}^{-d_1}A_1(z_{11}) + z_{12}^{-d_1}A_1(0)\{(1-z_{12})^{d_1+d_2}-1\} \\ &= z_{12}^{-d_1}A_1(0) + z_{12}^{-d_1}A_1(0)\{(1-z_{12})^{d_1+d_2}-1\} \\ &= A_1(0)z_{12}^{-d_1}(1-z_{12})^{d_1+d_2}, \end{aligned}$$

and, similarly,

$$z_{22}^{-d_2}u_2(s)^{d_2}\tilde{A}_2 = c_p^{d_2}A_2(0)z_{22}^{-d_2}(1-z_{22})^{d_1+d_2}.$$

Since $c_p^{d_2} = (-1)^{d_1+d_2}A_1(0)/A_2(0)$, they are equal.

Clearly they vanish to order $d_1 + d_2$ at x_1 .

(3) An isomorphism of $f_{0,p}$ and $f_{0,p'}$ is given by a compatible pair of isomorphisms $\varphi : \mathcal{C}_{0,p} \rightarrow \mathcal{C}_{0,p'}$ and $\psi : S_{0,s} \rightarrow S_{0,s}$.

The underlying stable map of φ is the identity map, since $Z_1 \neq Z_2$ by assumption. Therefore there exist functions $f_1(z_{12})$ with $f_1(0) = 1$ such that $\varphi^b z_{11} = z_{11}f_1(z_{12})$, and $f_2(z_{11})$ with $f_2(0) = 1$ such that $\varphi^b z_{12} = z_{12}f_2(z_{11})$.

On the other hand, φ is determined by a nonzero complex number γ such that $\psi^b \mathbf{m} = \mathbf{m}\gamma$.

Then we have

$$\begin{aligned} z_{11}z_{12}f_1(z_{12})f_2(z_{11}) &= \varphi^b(z_{11}z_{12}) \\ &= \psi^b(\mathbf{m}^{e_2}) \\ &= \mathbf{m}^{e_2}\gamma^{e_2}. \end{aligned}$$

Thus $f_1 \equiv f_2 \equiv 1$ and $\gamma^{e_2} = 1$.

Similarly, there exist functions $g_1(z_{22})$ with $g_1(0) = 1$ such that $\varphi^b z_{21} = z_{21}g_1(z_{22})$, and $g_2(z_{21})$ with $g_2(0) = 1$ such that $\varphi^b z_{22} = z_{22}g_2(z_{21})$.

$$\begin{aligned} z_{21}z_{22}g_1(z_{22})g_2(z_{21}) &= \varphi^b(z_{21}z_{22}) \\ &= \psi^b(\mathbf{m}^{e_1}c_{p'}) \\ &= \mathbf{m}^{e_1}\gamma^{e_1}c_{p'}. \end{aligned}$$

Thus $g_1 \equiv g_2 \equiv 1$ and $c_p = \gamma^{e_1}c_{p'}$. Thus $c_p/c_{p'}$ is an e_2 -th root of 1, and $p \equiv p' \pmod{d}$ follows. We can easily see that the converse is also true.

In the case $p = p'$, the calculation above shows $\varphi^b z_{ij} = z_{ij}$ and $\psi^b \mathbf{m} = \mathbf{m}$, and therefore that $f_{0,p}$ has no nontrivial automorphisms.

(4) By (3), $\overline{\mathbf{M}}_\beta$ is an algebraic space near $[f_{0,p}]$. Therefore, the assertion is clear if $n = 0$. If $n \geq 1$, the truncation $f_{1,p}$ is nontrivial, since $\alpha(\mathbf{m}) \neq 0$. Thus $S_n \rightarrow \overline{\mathbf{M}}_\beta$ has nonzero tangent map, and it is a closed immersion by Nakayama's Lemma. \square

4.4 Central fiber

Now we prove Theorem 4.12 (5).

We know that the curve is as in Lemma 4.10 (1) with $n = 0$ and the maps are given as in Lemma 4.10 (2) satisfying the conditions of Lemma 4.11. Here $m(s) = 0$ since it is in the maximal ideal. Note that by replacing \mathbf{m} (which does not affect $m(s)$) we may suppose that $u_1 = 1$.

Reducing modulo (s) , $\tilde{A}_i = A_i^{(0)} + z_{i1}A_{i1}^{(0)}(z_{i1}) + z_{i2}A_{i2}^{(0)}(z_{i2})$ and the condition of Lemma 4.10 (2) states

$$\begin{aligned} z_{i1}^{d_i}(A_i^{(0)} + z_{i1}A_{i1}^{(0)}(z_{i1}) + z_{i2}A_{i2}^{(0)}(z_{i2})) &= z_{i1}^{d_i}A_i(z_{i1}), \\ B_{ji0}^{(0)} + z_{i1}B_{ji1}^{(0)}(z_{i1}) + z_{i2}B_{ji2}^{(0)}(z_{i2}) &= B_{ji}(z_{i1}). \end{aligned}$$

Reducing modulo (s) , we are looking at $f|_0$, which is constant on C_0 and thus the terms in z_{i2} vanish. Hence $A_i^{(0)} + z_{i1}A_{i1}^{(0)}(z_{i1}) = A_i(z_{i1})$ and in particular $A_i^{(0)} = A_i(0)$. Thus it remains to show that $A_i(0) + z_{i2}A_{i2}^{(0)}(z_{i2}) = A_i(0)(1 - z_{i2})^{d_1+d_2}$ and that u_2 is one of c_p , the latter being equivalent to $u_2^{d_2} = (-1)^{d_1+d_2}A_1(0)/A_2(0)$.

In fact, for the function $A_{12}^{(0)}(z_{12}) \in \mathbb{C}[z_{12}, 1/(z_{12} - 1)]$, Lemma 4.11 (a) says that $A_1(0) + z_{12}A_{12}^{(0)}(z_{12})$ is regular also at $z_{12} = 1$. Therefore $A_{12}^{(0)}(z_{12})$ is in fact a polynomial. Similarly for $A_{22}^{(0)}(z_{22})$. Restricting to \mathcal{V}_0 so that $z_{11} = 0$ and $z_{21} = 0$, by Lemma 4.11 (a) we have

$$\begin{aligned} (4-1) \quad A_1(0) + z_{12}A_{12}^{(0)}(z_{12}) &= z_{12}^{d_1}u_1^{-d_1} \cdot z_{22}^{-d_2}u_2^{d_2}(A_2(0) + z_{22}A_{22}^{(0)}(z_{22})) \\ &= u_2^{d_2}z_{12}^{d_1+d_2}(A_2(0) + z_{12}^{-1}A_{22}^{(0)}(z_{12}^{-1})), \end{aligned}$$

and it follows that this is a polynomial of degree $d_1 + d_2$. It also must have vanishing order $d_1 + d_2$ at $z_{12} = 1$, and therefore $A_1(0) + z_{12}A_{12}^{(0)}(z_{12}) = A_1(0)(1 - z_{12})^{d_1+d_2}$. By symmetry we also have $A_2(0) + z_{22}A_{22}^{(0)}(z_{22}) = A_2(0)(1 - z_{22})^{d_1+d_2}$, and (4-1) implies that $(-1)^{d_1+d_2}A_1(0) = u_2^{d_2}A_2(0)$. (This is basically the same calculation as we did in the proof of (2).)

4.5 Extending deformations

Now we make the same assumption as in Theorem 1.14. In particular $N = 2$, and we denote B_{2i} by B_i etc. We may assume that

$$(4-2) \quad B'_i(0) \neq 0.$$

In fact, since Z_i is smooth at the intersection point P , this always holds if $d_i > 1$. If $d_i = 1$, this holds after changing coordinates if necessary.

By symmetry, assume that $d_1 \leq d_2$. Recall that $d_i = de_i$ with $\gcd(e_1, e_2) = 1$.

Lemma 4.13 Theorem 4.12 (6) (and hence Theorem 1.14) is reduced to the following claims.

- (A) If $e_1 = 1$, any extension of $f_{0,p}$ to S_1 is trivial.
- (B) If $e_1 \geq 2$, the set of extensions of $f_{0,p}$ to S_1 forms an (at most) 1-dimensional vector space.
- (C) For $e_1 \geq 2$, the family $f_{e_1-1,p}$ cannot be extended to a family over S_{e_1} .

Proof Since $f_{0,p}$ has no nontrivial automorphisms, $\overline{\mathcal{M}}_\beta$ is an algebraic space near $f_{0,p}$. If $e_1 = 1$, it is a reduced point by (A).

If $e_1 > 1$ the tangent space is 1-dimensional by (B). Thus the moduli space is étale locally a closed subspace of $\text{Spec } \mathbb{C}[s]$. If it contains S_{e_1} , our family $f_{e_1-1,p}$ induces a map $\iota : S_{e_1-1} \rightarrow S_{e_1}$ such that $f_{e_1-1,p}$ is isomorphic to the pullback by ι of a basic stable log map over S_{e_1} , and its tangent map is nonzero since $m(s) = s \not\equiv 0 \pmod{s^2}$. By composing with an automorphism of S_{e_1} , we may assume that ι is the standard closed immersion, and we would have an extension of $f_{e_1-1,p}$ to S_{e_1} . \square

So, for $1 \leq n \leq e_1$, we consider the family $f_{n-1,p}$ and study its extensions to S_n . We write

$$v := m^{(1)}$$

and we have

$$m(s) = \begin{cases} vs & \text{if } n = 1, \\ vs + m^{(n)}s^n & \text{and } v = 1 \text{ if } n \geq 2, \end{cases}$$

$$\begin{aligned} u_1(s) &= 1 + u_1^{(n)}s^n, \\ u_2(s) &= c_p + u_2^{(n)}s^n, \\ \tilde{A}_i &= A_i(z_{i1}) + A_i(0)\{(1 - z_{i2})^{d_1+d_2} - 1\} \\ &\quad + s^n(A_{i0}^{(n)} + z_{i1}A_{i1}^{(n)}(z_{i1}) + z_{i2}A_{i2}^{(n)}(z_{i2})), \\ \tilde{B}_i &= B_i(z_{i1}) + s^n(B_{i0}^{(n)} + z_{i1}B_{i1}^{(n)}(z_{i1}) + z_{i2}B_{i2}^{(n)}(z_{i2})) \end{aligned}$$

with $s^{n+1} = 0$, and see when the conditions of Lemma 4.11 are satisfied.

We have

$$\begin{aligned} z_{11}z_{12} &= m(s)^{e_2}u_1(s) \\ &= v^{e_2}s^{e_2}, \\ z_{21}z_{22} &= m(s)^{e_1}u_2(s) \\ &= c_p v^{e_1}s^{e_1}. \end{aligned}$$

Lemma 4.14 Condition (b) of Lemma 4.11 holds if and only if the following hold:

$$B_{10}^{(n)} = B_{20}^{(n)}$$

and

$$B_{12}^{(n)}(z_{12}) \equiv B_{12}^{(n)}(0) \text{ and } B_{22}^{(n)}(z_{22}) \equiv B_{22}^{(n)}(0)$$

with

$$\begin{aligned} B_{12}^{(n)}(0) &= \begin{cases} B'_2(0)c_p v^n & \text{if } n = e_1, \\ 0 & \text{otherwise,} \end{cases} \\ B_{22}^{(n)}(0) &= \begin{cases} B'_1(0)v^n & \text{if } n = e_2, \text{ i.e., } n = e_1 = e_2 = 1, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Proof Lemma 4.11 (b) says that

$$\begin{aligned} &B_1(0) + s^n(B_{10}^{(n)} + z_{12}B_{12}^{(n)}(z_{12})) + s^{e_2}B'_1(0)v^{e_2}z_{12}^{-1} \\ &= B_2(0) + s^n(B_{20}^{(n)} + z_{12}^{-1}B_{22}^{(n)}(z_{12}^{-1})) + s^{e_1}B'_2(0)c_p v^{e_1}z_{12}, \end{aligned}$$

and that both sides are extendable to $z_{12} = 1$.

From the regularity at $z_{12} = 1$, we see that $B_{i2}^{(n)}(z_{i2})$ are polynomials, and comparing the coefficients, we obtain the assertion.

For the other implication, note that $B_1(0) = B_2(0)$ is assumed. \square

We have a natural trivialization $\mathcal{C}_i^\circ \cong C_i^\circ \times S \cong (\mathbb{P}^1 \setminus \{0\}) \times S$ given by the functions z_{i0} and z_{i1} . Then the morphism $\mathcal{C}_i^\circ \cap \mathcal{V} \rightarrow X$ by data \tilde{A}_i and \tilde{B}_{ji} can be considered as a deformation of $f^{(i)}|_{\mathcal{C}_i^\circ \cap V}$. Since it is trivial modulo s^n and $(s^n)^2 = 0$, we obtain a vector field $\mathbf{v}_i \in \Gamma(C_i^\circ \cap V, (f^{(i)})^*T_X)$ to X along $f^{(i)}|_{\mathcal{C}_i^\circ \cap V}$. Specifically, if φ is a local regular

function on X we can write $(f|_{C_i^\circ})^* \varphi = (f^{(i)})^* \varphi + s^n D(\varphi)$. Then it is easy to check that D defines a derivation $(f^{(i)}|_{C_i^\circ \cap V})^{-1} \mathcal{O}_X \rightarrow \mathcal{O}_{C_i^\circ \cap V}$, hence a section $\mathbf{v}_i \in \Gamma(C_i^\circ \cap V, (f^{(i)})^* T_X)$.

By the same reasoning, if the condition (c) of Lemma 4.11 holds, then \mathbf{v}_i extends to a section in $\Gamma(C_i^\circ, (f^{(i)})^* T_X)$.

Let us explicitly write down \mathbf{v}_i . On $C_1^\circ \cap V$, from $s^{n+1} = 0$, $z_{12} = z_{11}^{-1} v^{e_2} s^{e_2}$ and $e_2 \geq n$ we have

$$z_{11}^{d_1} \tilde{A}_1 = z_{11}^{d_1} A_1(z_{11}) + s^n z_{11}^{d_1} (A_{10}^{(n)} + z_{11} A_{11}^{(n)}(z_{11})) + s^{e_2} z_{11}^{d_1-1} (-(d_1 + d_2) A_1(0)) v^{e_2}$$

and

$$\tilde{B}_1 = B_1(z_{11}) + s^n (B_{10}^{(n)} + z_{11} B_{11}^{(n)}(z_{11})).$$

Thus we have

$$\mathbf{v}_1 = z_{11}^{d_1} (A_{10}^{(n)} + z_{11} A_{11}^{(n)}(z_{11})) \partial_{w_1} + (B_{10}^{(n)} + z_{11} B_{11}^{(n)}(z_{11})) \partial_{w_2}$$

if $n < e_2$ and

$$\mathbf{v}_1 = z_{11}^{d_1} (z_{11}^{-1} (-(d_1 + d_2) A_1(0)) v^n + A_{10}^{(n)} + z_{11} A_{11}^{(n)}(z_{11})) \partial_{w_1} + (B_{10}^{(n)} + z_{11} B_{11}^{(n)}(z_{11})) \partial_{w_2}$$

if $n = e_2$ (i.e. $n = e_1 = e_2 = 1$).

Similarly we have

$$\mathbf{v}_2 = z_{21}^{d_2} (A_{20}^{(n)} + z_{21} A_{21}^{(n)}(z_{21})) \partial_{w_1} + (B_{20}^{(n)} + z_{21} B_{21}^{(n)}(z_{21})) \partial_{w_2}$$

if $n < e_1$ and

$$\mathbf{v}_2 = z_{21}^{d_2} (z_{21}^{-1} (-(d_1 + d_2) A_2(0)) c_p v^n + A_{20}^{(n)} + z_{21} A_{21}^{(n)}(z_{21})) \partial_{w_1} + (B_{20}^{(n)} + z_{21} B_{21}^{(n)}(z_{21})) \partial_{w_2}$$

if $n = e_1$.

Lemma 4.15 Let \mathcal{E}_i be the \mathcal{O}_{C_i} -submodule of $(f^{(i)})^* T_X$ generated by $(f^{(i)})^* T_X(-\log D)$ and T_{C_i} .

- (1) The sheaf \mathcal{E}_i can also be described as the \mathcal{O}_{C_i} -submodule of $(f^{(i)})^* T_X$ generated by $(f^{(i)})^* T_X(-\log D)$ and $z_{i1}^{d_1-1} \partial_{w_1}$ at R_i .
- (2) If the condition (c) of Lemma 4.11 holds, then \mathbf{v}_i extends to a global section of \mathcal{E}_i .
- (3) There is a commutative diagram with exact rows and columns as follows.

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \downarrow & & \downarrow & & \\
0 & \longrightarrow & T_{C_i}(-\log R_i) & \longrightarrow & (f^{(i)})^* T_X(-\log D) & \longrightarrow & \mathcal{O}_{C_i}(-1) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \parallel \\
0 & \longrightarrow & T_{C_i} & \longrightarrow & \mathcal{E}_i & \longrightarrow & \mathcal{O}_{C_i}(-1) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \\
& & \mathbb{C} & \xlongequal{\quad} & \mathbb{C} & & \\
& & \downarrow & & \downarrow & & \\
& & 0 & & 0 & &
\end{array}$$

- (4) The natural map $\Gamma(T_{C_i}) \rightarrow \Gamma(\mathcal{E}_i)$ is an isomorphism.

Proof (1) Direct calculations.

(2) As explained right after the definition of \mathbf{v}_i , if we assume Lemma 4.11 (c), then \mathbf{v}_i extends to a section of $(f^{(i)})^*T_X$ over C_i° . Note that $(f^{(i)})^*T_X|_{C_i^\circ} = \mathcal{E}_i|_{C_i^\circ}$.

We see that \mathbf{v}_i also belongs to $(\mathcal{E}_i)_{R_i}$ from our explicit description of \mathbf{v}_i and (1), and we have the assertion.

(3) The first homomorphism in the first row is injective with an invertible cokernel since $f^{(i)}$ is a log map which is immersive (although the immersivity at R_i is not needed *here*). From $(K_X + D).Z_i = 0$ we see that the cokernel is $\mathcal{O}_{C_i}(-1)$. Then the assertion is easy to prove.

(4) follows from (3). \square

Thus we have global vector fields $a_1(z_{11})\partial_{z_{11}}$ and $a_2(z_{21})\partial_{z_{21}}$ on C_1 and C_2 respectively such that

$$\mathbf{v}_1 = a_1(z_{11})(f^{(1)})_*\partial_{z_{11}}, \mathbf{v}_2 = a_2(z_{21})(f^{(2)})_*\partial_{z_{21}}.$$

Comparing at R_i , we have

$$(4-3) \quad a_1(0) = \begin{cases} 0 & \text{if } n < e_2, \\ -\frac{d_1+d_2}{d_1}v^n & \text{if } n = e_2 \text{ (i.e. } n = e_1 = e_2 = 1\text{),} \end{cases}$$

$$(4-4) \quad a_2(0) = \begin{cases} 0 & \text{if } n < e_1, \\ -\frac{d_1+d_2}{d_2}c_p v^n & \text{if } n = e_1, \end{cases}$$

$$\begin{aligned} a_1(0)B'_1(0) &= B_{10}^{(n)}, \\ a_2(0)B'_2(0) &= B_{20}^{(n)}. \end{aligned}$$

We also have $B_{10}^{(n)} = B_{20}^{(n)}$ by Lemma 4.14.

Lemma 4.16 If $n = e_1$, then $v = 0$.

Proof If $n = e_1 = e_2$, then they are all 1. By $(Z_1.Z_2)_P = d$ we have $A_1(0)(B'_2(0))^d \neq A_2(0)(B'_1(0))^d$. Also, $c_p^d = A_1(0)/A_2(0)$ by definition. From these we obtain $v = 0$.

If $n = e_1 < e_2$, then we have $a_1(0) = 0$, and then $v = 0$ as we have $B'_2(0) \neq 0$ (since $d_2 \geq 2$ and C_2 is smooth at the point of intersection P). \square

Proof of Lemma 4.13 (C) We are considering the case $n = e_1 \geq 2$, so $v = m^{(1)} = 1$ by our setup. But we also have $v = 0$ by Lemma 4.16, which yields a contradiction. \square

Lemma 4.17 The deformation of the underlying prestable curve is trivial, $B_{12}^{(n)}(z_{12}) = B_{22}^{(n)}(z_{22}) = 0$, and $a_1(0) = a_2(0) = 0$.

Proof If $n = e_1$, then $v = 0$ by Lemma 4.16, and from equations (4-3) and (4-4) we have $a_1(0) = a_2(0) = 0$ in any case. Similarly, from Lemma 4.14 it follows that $B_{12}^{(n)}(z_{12}) = B_{22}^{(n)}(z_{22}) = 0$.

The deformation parameters of the underlying curve are

$$m(s)^{e_2}u_1(s) = v^{e_2}s^{e_2} = 0 \text{ and } m(s)^{e_1}u_2(s) = c_p v^{e_1}s^{e_1} = 0,$$

for $v = 0$ if $n = e_1$ and $s^{e_1} = s^{e_2} = 0$ if $n < e_1$. So the deformation of the curve is trivial. \square

Since $a_i(0) = 0$, $a_1(z_{11})\partial_{z_{11}}$ and $a_2(z_{21})\partial_{z_{21}}$ define an automorphism of \mathcal{C}_{S_n} over S_n , and by untwisting we may assume that the underlying stable map is a trivial deformation, i.e. $\mathbf{v}_1 = \mathbf{v}_2 = 0$, or more explicitly,

$$A_{10}^{(n)} = A_{11}^{(n)}(z_{11}) = B_{10}^{(n)} = B_{11}^{(n)}(z_{11}) = A_{20}^{(n)} = A_{21}^{(n)}(z_{21}) = B_{20}^{(n)} = B_{21}^{(n)}(z_{21}) = 0.$$

Lemma 4.18 Assuming $A_{10}^{(n)} = A_{20}^{(n)} = 0$, condition (a) of Lemma 4.11 implies $A_{12}^{(n)}(z_{12}) = A_{22}^{(n)}(z_{22}) = 0$ and $d_1 u_1^{(n)} = d_2 c_p^{-1} u_2^{(n)}$.

Proof Again noting that either $v = 0$ or $s^{e_1} = s^{e_2} = 0$, and also that $z_{11} = z_{21} = 0$ on \mathcal{V}_0 , we have

$$\begin{aligned} z_{12}^{-d_1} u_1(s)^{d_1} \tilde{A}_1 &= A_1(0) z_{12}^{-d_1} (1 - z_{12})^{d_1+d_2} \\ &\quad + s^n z_{12}^{-d_1} (A_{10}^{(n)} + z_{12} A_{12}^{(n)}(z_{12}) + d_1 A_1(0) u_1^{(n)} (1 - z_{12})^{d_1+d_2}) \end{aligned}$$

and

$$\begin{aligned} z_{22}^{-d_2} u_2(s)^{d_2} \tilde{A}_2 &= A_2(0) c_p^{d_2} z_{22}^{-d_2} (1 - z_{22})^{d_1+d_2} \\ &\quad + s^n c_p^{d_2} z_{22}^{-d_2} (A_{20}^{(n)} + z_{22} A_{22}^{(n)}(z_{22}) + d_2 A_2(0) c_p^{-1} u_2^{(n)} (1 - z_{22})^{d_1+d_2}) \end{aligned}$$

on $\mathcal{V}_0 \setminus \{x_1\}$.

By the regularity condition at $z_{12} = 1$ from Lemma 4.11 (a), $A_{i2}^{(n)}(z_{i2})$ are polynomials. The equality in Lemma 4.11 (a) is equivalent to

$$\begin{aligned} &A_{10}^{(n)} + z_{12} A_{12}^{(n)}(z_{12}) + d_1 A_1(0) u_1^{(n)} (1 - z_{12})^{d_1+d_2} \\ &= c_p^{d_2} z_{12}^{d_1+d_2} (A_{20}^{(n)} + z_{12}^{-1} A_{22}^{(n)}(z_{12}^{-1})) + d_2 A_2(0) c_p^{d_2-1} u_2^{(n)} (z_{12} - 1)^{d_1+d_2}, \end{aligned}$$

and from this we see that $\deg(A_{10}^{(n)} + z_{12} A_{12}^{(n)}(z_{12})) \leq d_1 + d_2$. Similarly, $\deg(A_{20}^{(n)} + z_{22} A_{22}^{(n)}(z_{22})) \leq d_1 + d_2$. From the condition on the vanishing order at \tilde{x}_1 , we see that $A_{10}^{(n)} + z_{12} A_{12}^{(n)}(z_{12}) = A_{10}^{(n)} (1 - z_{12})^{d_1+d_2}$, $A_{20}^{(n)} + z_{22} A_{22}^{(n)}(z_{22}) = A_{20}^{(n)} (1 - z_{22})^{d_1+d_2}$, with

$$A_{10}^{(n)} + d_1 A_1(0) u_1^{(n)} = (-1)^{d_1+d_2} c_p^{d_2} (A_{20}^{(n)} + d_2 c_p^{-1} A_2(0) u_2^{(n)}).$$

But $A_{10}^{(n)} = A_{20}^{(n)} = 0$, so $A_{12}^{(n)}(z_{12}) = A_{22}^{(n)}(z_{22}) = 0$ and $d_1 A_1(0) u_1^{(n)} = (-1)^{d_1+d_2} c_p^{d_2} d_2 c_p^{-1} A_2(0) u_2^{(n)}$, i.e. $d_1 u_1^{(n)} = d_2 c_p^{-1} u_2^{(n)}$. \square

Proof of Lemma 4.13 (A) and (B) By Lemma 4.17, the curve \mathcal{C}_{S_n}/S_n is trivial and we may assume that

$$\begin{aligned} A_{10}^{(n)} &= A_{11}^{(n)}(z_{11}) = A_{12}^{(n)}(z_{12}) = 0, \\ B_{10}^{(n)} &= B_{11}^{(n)}(z_{11}) = B_{12}^{(n)}(z_{12}) = 0, \\ A_{20}^{(n)} &= A_{21}^{(n)}(z_{21}) = A_{22}^{(n)}(z_{22}) = 0, \\ B_{20}^{(n)} &= B_{21}^{(n)}(z_{21}) = B_{22}^{(n)}(z_{22}) = 0, \end{aligned}$$

i.e. \tilde{A}_i and \tilde{B}_i are constant with respect to s .

Let $\mathbf{m}' := \mathbf{m}(1 + (u_1^{(n)}/e_2)s^n)$, then we have

$$\alpha(\mathbf{m}') = \alpha(\mathbf{m}) \left(1 + \frac{u_1^{(n)}}{e_2} s^n \right) = \alpha(\mathbf{m})$$

and

$$\mathbf{m}^{e_2} u_1(s) = (\mathbf{m}')^{e_2}.$$

So, if we replace \mathbf{m} by \mathbf{m}' , we see that $m(s)$ does not change and that $u_1^{(n)} = 0$. By Lemma 4.18, $u_2^{(n)} = 0$. Since \tilde{A}_i and \tilde{B}_i do not depend on \mathbf{m} , they do not change either.

If $n = e_1 = 1$, we also have $v = 0$, and the deformation is trivial.

If $n = 1 < e_1$, this says that the only deformation parameter is v . \square

Example 4.19 We illustrate by an example the difference in our setting between the moduli space of stable log maps and the relative maps of [30, 31]. For this example the divisor is normal crossing, so the more appropriate moduli space is the one of [38], but since the calculations are local, it is not relevant.

Start with \mathbb{P}^2 with its toric boundary $D = L_1 \cup L_2 \cup L_3$. Choose $P \in L_1$ a smooth point of D . Choose some local coordinates w_1, w_2 around P such that L_1 is given by $w_1 = 0$ and $P = (0, 0)$. We consider Z'_1 and Z'_2 two general smooth conics tangent to L_1 at P . For example, consider the conics given near P by

$$\begin{aligned} Z'_1 : \quad & (w_1 - 1)^2 + w_2^2 - 1 = 0, \\ Z'_2 : \quad & 2(w_1 - 2)^2 + w_2^2 - 8 = 0. \end{aligned}$$

In particular, $(Z'_1 \cdot Z'_2)_P = 2$ and Z'_1 and Z'_2 meet in 2 other points. Also, we may assume that $Z'_1 \cup Z'_2$ is disjoint from the torus-fixed points. Write P_1, \dots, P_8 for the distinct points of intersection of $(Z'_1 \cup Z'_2) \setminus P$ with $L_2 \cup L_3$.

Let X be the blowup of \mathbb{P}^2 in P_1, \dots, P_8 with exceptional divisors E_1, \dots, E_8 and by abusing notation slightly we denote by D the proper transform of $L_1 \cup L_2 \cup L_3$. Let Z_1 and Z_2 be the proper transforms of Z'_1 and Z'_2 . Then Z_1 and Z_2 satisfy the hypotheses of Theorem 1.14 for (X, D) and $Z_1 + Z_2$ is in the class $\beta = 4H - \sum_{i=1}^8 E_i$, where H is the pullback of the hyperplane class.

Consider the chain of \mathbb{P}^1 's given by $C = C_1 \cup_{R_1} C_0 \cup_{R_2} C_2$ with R_i nodes with local equations $z_{11}z_{12} = 0$ and $z_{21}z_{22} = 0$, where z_{i1} are local coordinates for C_i ($i = 1, 2$) and z_{i2} are inhomogeneous coordinates on C_0 satisfying $z_{12}z_{22} = 1$ on $C_0 \setminus \{R_1, R_2\}$. The nodes R_1 and R_2 are the points at infinity of the projective completion of the affine curve $z_{12}z_{22} = 1$. Choosing parametrizations of Z_1 and Z_2 , there is only one 1-marked stable map $f : C_1 \cup C_0 \cup C_2 \rightarrow X$ with $f_* C = Z_1 + Z_2$ which underlies a stable log map of maximal tangency. Locally near P and for $f_i := f|_{C_i}$,

$$\begin{aligned} f_1(z_{11}) &= \left(\frac{2z_{11}^2}{z_{11}^2 + 1}, \frac{2z_{11}}{z_{11}^2 + 1} \right), \\ f_2(z_{21}) &= \left(\frac{8z_{21}^2}{2z_{21}^2 + 1}, \frac{8z_{21}}{2z_{21}^2 + 1} \right), \end{aligned}$$

and C_0 is mapped to P .

By Theorem 4.12, we have two nonisomorphic stable log maps over f each of which has multiplicity 1 in $\overline{\mathcal{M}}_\beta$. In what follows, we illustrate the log structure assuming $n = 0$.

Pulling back along f_i as in Notation 4.9,

$$w_1 \mapsto z_{i1}^2 A_i(z_{i1}),$$

so that we have $A_1(z_{11}) = 2/(z_{11}^2 + 1)$ and $A_2(z_{21}) = 8/(2z_{21}^2 + 1)$ with $A_1(0)/A_2(0) = 1/4$.

The base is $S = \text{Spec } \mathbb{C}$ with log structure $\mathcal{M}_S = \coprod_{a=0}^\infty \mathbf{m}^a \mathbb{C}^\times$, with $\alpha(\mathbf{m}^a) = 0^a$. At each node of C , the log structure \mathcal{M}_{C, R_i} is $\coprod_{a,b,c=0}^\infty z_{i1}^a z_{i2}^b \mathbf{m}^c \mathcal{O}_{C, R_i}^\times$ subject to the relation $z_{i1} z_{i2} = \mathbf{m} u_i$ with $\alpha(z_{ij}) = z_{ij}$. Up to isomorphism, we may set $u_1 = 1$.

Near P , let \mathbf{w}_1 be the unique lift of w_1 to \mathcal{M}_X . Recall that x_1 is the marked point. By Theorem 4.12, near R_i away from x_1 , we have

$$\begin{aligned} f_1^\flat(\mathbf{w}) &= z_{11}^2 \left(\frac{2}{z_{11}^2 + 1} + 2((1 - z_{12})^4 - 1) \right), \\ f_2^\flat(\mathbf{w}) &= z_{21}^2 \left(\frac{8}{2z_{21}^2 + 1} + 8((1 - z_{22})^4 - 1) \right). \end{aligned}$$

By the generization maps to C_0 , sending z_{i1} to $\mathbf{m} u_i z_{i2}^{-1}$ respectively, they have to agree on $C_0 \setminus \{R_1, R_2, x_1\}$. Thus,

$$\mathbf{m}^2 z_{12}^{-2} (2((1 - z_{12})^4)) = \mathbf{m}^2 u_2^2 z_{22}^{-1} (8((1 - z_{22})^4))$$

as $z_{i1} = 0$ on $C_0 \setminus \{R_1, R_2, x_1\}$. Since $z_{12}z_{22} = 1$, we conclude $u_2 = \pm 1/2$. Therefore we have two stable log maps.

Example 4.20 Let us look at the relative side. Since we work locally near $Z_1 \cup Z_2$, we may think of D as smooth. Before proceeding with the example, we quickly recall some of the notation and ideas of the theory of relative stable maps, referring the reader to [30] for more detail.

In the theory of relative stable maps to X , the target space X is allowed to deform in a specific family over \mathbb{A}^n to singular limits $X[n]$. The central fiber $X[n]_0$ is a chain of $n + 1$ smooth components obtained by gluing X to \mathbb{P}^1 -bundles $\mathbb{P}_1, \dots, \mathbb{P}_n$ over D . The gluings are along $D \subset X$ and sections of the \mathbb{P}_j . There are n singular divisors D_1, \dots, D_n in $X[n]$. The divisor D_1 is the intersection of X and \mathbb{P}_1 , while D_k is the intersection of \mathbb{P}_{k-1} and \mathbb{P}_k for $k > 1$. We will describe $X[1]_0$ in more detail below.

If $f : C \rightarrow X[n]_0$ is a relative stable map, then no component of C is mapped into any D_k . Furthermore, there is the *predeformability* condition: if $p \in f^{-1}(D_k)$, then p is a node of C . One component C_i of C containing p satisfies $f(C_i) \subset \mathbb{P}_{k-1}$, while another such component C_j satisfies $f(C_j) \subset \mathbb{P}_k$, with an obvious modification if $k = 1$. Then the predeformability requirement is that $\text{mult}_p(f|_{C_i}^*(D_k)) = \text{mult}_p(f|_{C_j}^*(D_k))$. For the predeformability of a family of maps, we require a little more ([30, Definition 2.9, Lemma 2.4, Definition 2.3]). In the case at hand, we explain the condition later.

In the current case, it turns out to be sufficient to consider $X[1]_0$, the central fiber of the blow-up of $X \times \mathbb{A}^1$ along $D \times \{0\}$. This is also obtained by gluing X and the \mathbb{P}^1 -bundle $\mathbb{P}(N_{D/X} \oplus \mathcal{O}_D)$ over D .

We can describe $X[1]_0$ explicitly as follows. Let $X'_0 = \mathbb{A}^2 \setminus (L_2 \cup L_3)$, where $L_2 \cup L_3$ is the union of two distinct lines that will corresponds to $L_2 \cup L_3 \subset \mathbb{P}^2$. Consider \mathbb{A}^3 with coordinates w_1, w_2, v_1 and projection $p : \mathbb{A}^3 \rightarrow \mathbb{A}^2$ given by w_1, w_2 , let $X' = p^{-1}(X'_0) \cap V(w_1 v_1)$ and $U = (L_1 \cap X'_0) \times \mathbb{A}^1 \subset \mathbb{A}^2$ with coordinates w_2, u_1 . Then we may think of $X[1]_0$ as covered by open sets X' , U and $X \setminus D$: The open sets $(v_1 \neq 0) \subset X'$ and $(u_1 \neq 0) \subset U$ are identified by $u_1 v_1 = 1$, and $(w_1 \neq 0) \subset X'$ is identified with an open subset of $X \setminus D$ in a natural way.

We have a projection map $\pi : X[1]_0 \rightarrow X$ with $\pi|_{X'} = p|_{X'}$, which identifies $(v_1 = 0) \subset X'$ with X'_0 and contracts U to D . The divisor $(u_1 = 0) \subset U$ will be considered as the boundary divisor.

Let $S = \text{Spec } \mathbb{C}[s]/(s^2)$. We will define a nontrivial family of relative stable maps $\mathcal{C}_S \rightarrow X[1]_0$ whose central fiber, composed with π , gives \underline{f} in the previous example. To do this, we take \mathcal{C}_S to be the family of curves defined by $\mu_1 = s$ and $\mu_2 = s/4$ (hence $z_{11} z_{12} = s$ and $z_{21} z_{22} = s/4$) and give a stable map $\mathcal{C}_S \rightarrow X[1]_0$ over S such that the following hold:

- (pullback of the boundary) The pullback of $(u_1 = 0)$ is $4 \cdot \tilde{x}_1$.
- (predeformability) Near R_i , one can write $w_1 = z_{i1}^2 \tilde{A}_i$, $v_1 = z_{i2}^2 \tilde{B}_i$ with \tilde{A}_i, \tilde{B}_i invertible and $\tilde{A}_i \tilde{B}_i \in \mathbb{C}[s]/(s^2)$.

Once such a family of relative stable maps is obtained, it is nontrivial since \mathcal{C}_S is a nontrivial deformation.

We set

$$\begin{aligned} \tilde{A}_1 &= \frac{2(1 - z_{12})^4}{z_{11}^2 + 1}, & \tilde{B}_1 &= \frac{z_{11}^2 + 1}{(1 - z_{12})^4}, \\ \tilde{A}_2 &= \frac{8(1 - z_{22})^4}{2z_{21}^2 + 1}, & \tilde{B}_2 &= \frac{2z_{21}^2 + 1}{(1 - z_{22})^4}, \\ w_2 &= \frac{2z_{11} + s(4z_{11}^2 - 4 + 2z_{12})}{z_{11}^2 + 1} \text{ near } R_1, \\ w_2 &= \frac{8z_{21} + s(8z_{21}^2 - 4 + 2z_{22})}{2z_{21}^2 + 1} \text{ near } R_2. \end{aligned}$$

These data give a stable map $\mathcal{C}_S \rightarrow X[1]_0$. In fact, let $\mathcal{V} \subset \mathcal{C}_S$ be the open subscheme supported on $\underline{f}^{-1}(X'_0) \setminus \{x_1\}$. We first see that the two sets of w_1, v_1 (defined from \tilde{A}_i and \tilde{B}_i as above) and w_2 coincide on $\underline{C}_0 \setminus \{R_1, R_2, x_1\}$

and give a morphism $\mathcal{V} \rightarrow \mathbb{A}^3$: On $C_0 \setminus \{R_1, R_2, x_1\}$, $z_{11}^2 \tilde{A}_1 = z_{21}^2 \tilde{A}_2 = 0$, $z_{12}^2 \tilde{B}_1 = z_{12}^2 / (1 - z_{12})^4$ is equal to $z_{22}^2 \tilde{B}_2 = z_{22}^2 / (1 - z_{22})^4$, and the two expressions for w_2 are

$$\frac{2z_{11} + s(4z_{11}^2 - 4 + 2z_{12})}{z_{11}^2 + 1} = \frac{2sz_{12}^{-1} + s(4(sz_{12}^{-1})^2 - 4 + 2z_{12})}{(sz_{12}^{-1})^2 + 1} = s(2z_{12}^{-1} - 4 + 2z_{12})$$

and

$$\frac{8z_{21} + s(8z_{21}^2 - 4 + 2z_{22})}{2z_{21}^2 + 1} = \frac{8(s/4)z_{22}^{-1} + s(8((s/4)z_{22}^{-1})^2 - 4 + 2z_{22})}{2((s/4)z_{22}^{-1})^2 + 1} = s(2z_{22}^{-1} - 4 + 2z_{22}),$$

which are equal.

Since $w_1 v_1 = z_{i1}^2 z_{i2}^2 \tilde{A}_i \tilde{B}_i = 0$, we have a morphism $\tilde{f}' : \mathcal{V} \rightarrow X'$.

On $\mathcal{V} \cap (C_1 \setminus R_1)$,

$$(w_1, w_2) = \left(\frac{2z_{11}^2 - 8sz_{11}}{z_{11}^2 + 1}, \frac{2z_{11} + 4s(z_{11}^2 - 1)}{z_{11}^2 + 1} \right),$$

and this is the reparametrization of f_1 by $z_{11} \mapsto z_{11} - 2s(z_{11}^2 + 1)$. Similarly, the restriction of \tilde{f}' to $\mathcal{V} \cap (C_2 \setminus R_2)$ is given by the reparametrization of f_2 by $z_{21} \mapsto z_{21} - (1/2)s(2z_{21}^2 + 1)$. This implies that \tilde{f}' extends to a morphism $\mathcal{C}_S \setminus \{x_1\} \rightarrow X[1]_0$. (We may also verify that w_1 and w_2 satisfy the equation for Z'_i .)

On $C_0 \setminus \{R_1, R_2\}$, we have $u_1 = (v_1)^{-1} = (1 - z_{12})^4 / z_{12}^2$, so \tilde{f}' extends to a morphism $\mathcal{C}_S \rightarrow X[1]_0$, and the condition on the pullback of the boundary is satisfied. Predeformability also holds since $\tilde{A}_1 \tilde{B}_1 = 2$ and $\tilde{A}_2 \tilde{B}_2 = 8$.

It follows from the main result of [41] that the corresponding point of the moduli space of relative stable map is actually isomorphic to $S = \text{Spec } \mathbb{C}[s]/(s^2)$, in contrast to the 2 reduced points in the moduli space of basic stable log maps.

References

- [1] D Abramovich, Q Chen, *Stable logarithmic maps to Deligne-Faltings pairs II*, Asian J. Math. 18 (2014) 465–488
- [2] D Abramovich, S Marcus, J Wise, *Comparison theorems for Gromov-Witten invariants of smooth pairs and of degenerations*, Ann. Inst. Fourier (Grenoble) 64 (2014) 1611–1667
- [3] A B Altman, A Iarrobino, S L Kleiman, *Irreducibility of the compactified Jacobian*, from “Real and complex singularities (Proc. Ninth Nordic Summer School/NAVF Sympos. Math., Oslo, 1976)”, Sijthoff and Noordhoff, Alphen aan den Rijn (1977) 1–12
- [4] P S Aspinwall, D R Morrison, *Topological field theory and rational curves*, Comm. Math. Phys. 151 (1993) 245–262
- [5] L Barrott, N Nabijou, *Tangent curves to degenerating hypersurfaces* In preparation, 2020
- [6] A Beauville, *Counting rational curves on K3 surfaces*, Duke Math. J. 97 (1999) 99–108
- [7] P Bousseau, *A proof of N. Takahashi’s conjecture on genus zero Gromov-Witten theory of (\mathbb{P}^2, E)* arXiv:1909.02992, 2019
- [8] P Bousseau, *Scattering diagrams, stability conditions, and coherent sheaves on \mathbb{P}^2* arXiv:1909.02985, 2019
- [9] P Bousseau, A Brini, M van Garrel, *On the log-local principle for the toric boundary* arXiv:1908.04371, 2019
- [10] P Bousseau, A Brini, M van Garrel, *Stable maps to Looijenga pairs*, in preparation (2020)
- [11] T Bridgeland, *Stability conditions on a non-compact Calabi-Yau threefold*, Comm. Math. Phys. 266 (2006) 715–733
- [12] L Caporaso, J Coelho, E Esteves, *Abel maps of Gorenstein curves*, Rend. Circ. Mat. Palermo (2) 57 (2008) 33–59
- [13] Q Chen, *Stable logarithmic maps to Deligne-Faltings pairs I*, Ann. of Math. (2) 180 (2014) 455–521
- [14] X Chen, *Rational curves on K3 surfaces*, J. Algebraic Geom. 8 (1999) 245–278

- [15] **T-M Chiang, A Klemm, S-T Yau, E Zaslow**, *Local mirror symmetry: calculations and interpretations*, Adv. Theor. Math. Phys. 3 (1999) 495–565
- [16] **J Choi, M van Garrel, S Katz, N Takahashi**, *Log BPS numbers of log Calabi-Yau surfaces*, to appear in Trans. Amer. Math. Soc. [arXiv:1810.02377](https://arxiv.org/abs/1810.02377), 2018
- [17] **J Choi, M van Garrel, S Katz, N Takahashi**, *Local BPS Invariants: Enumerative Aspects and Wall-Crossing*, Int. Math. Res. Not. (2018)Rny171
- [18] **B Fantechi, L Göttsche, D van Straten**, *Euler number of the compactified Jacobian and multiplicity of rational curves*, J. Algebraic Geom. 8 (1999) 115–133
- [19] **M v Garrel, T Graber, H Ruddat**, *Local Gromov-Witten invariants are log invariants*, Adv. Math. 350 (2019) 860–876
- [20] **M v Garrel, D P Overholser, H Ruddat**, *Enumerative aspects of the Gross-Siebert program*, from “Calabi-Yau varieties: arithmetic, geometry and physics”, Fields Inst. Monogr. 34, Fields Inst. Res. Math. Sci., Toronto, ON (2015) 337–420
- [21] **M van Garrel, T W H Wong, G Zaimi**, *Integrality of relative BPS state counts of toric del Pezzo surfaces*, Commun. Number Theory Phys. 7 (2013) 671–687
- [22] **T Gräfnitz**, *Tropical correspondence for smooth del Pezzo log Calabi-Yau pairs* [arXiv:2005.14018](https://arxiv.org/abs/2005.14018), 2020
- [23] **M Gross, R Pandharipande, B Siebert**, *The tropical vertex*, Duke Math. J. 153 (2010) 297–362
- [24] **M Gross, B Siebert**, *Logarithmic Gromov-Witten invariants*, J. Amer. Math. Soc. 26 (2013) 451–510
- [25] **E-N Ionel, T H Parker**, *The Gopakumar-Vafa formula for symplectic manifolds*, Ann. of Math. (2) 187 (2018) 1–64
- [26] **F Kato**, *Log smooth deformation and moduli of log smooth curves*, Internat. J. Math. 11 (2000) 215–232
- [27] **S Katz**, *Genus zero Gopakumar-Vafa invariants of contractible curves*, J. Differential Geom. 79 (2008) 185–195
- [28] **A Klemm, D Maulik, R Pandharipande, E Scheidegger**, *Noether-Lefschetz theory and the Yau-Zaslow conjecture*, J. Amer. Math. Soc. 23 (2010) 1013–1040
- [29] **M Kontsevich**, *Enumeration of rational curves via torus actions*, from “The moduli space of curves (Texel Island, 1994)”, Progr. Math. 129, Birkhäuser Boston, Boston, MA (1995) 335–368
- [30] **J Li**, *Stable morphisms to singular schemes and relative stable morphisms*, J. Differential Geom. 57 (2001) 509–578
- [31] **J Li**, *A degeneration formula of GW-invariants*, J. Differential Geom. 60 (2002) 199–293
- [32] **Y-S Lin**, *Correspondence Theorem between Holomorphic Discs and Tropical Discs on K3 Surfaces* To appear in J. Diff. Geom.
- [33] **Y-S Lin**, *Open Gromov-Witten invariants on elliptic K3 surfaces and wall-crossing*, Comm. Math. Phys. 349 (2017) 109–164
- [34] **Y I Manin**, *Generating functions in algebraic geometry and sums over trees*, from “The moduli space of curves (Texel Island, 1994)”, Progr. Math. 129, Birkhäuser Boston, Boston, MA (1995) 401–417
- [35] **S Mukai**, *Symplectic structure of the moduli space of sheaves on an abelian or K3 surface*, Invent. Math. 77 (1984) 101–116
- [36] **N Nabijou, D Ranganathan**, *Gromov-Witten theory with maximal contacts* [arXiv:1908.04706](https://arxiv.org/abs/1908.04706), 2019
- [37] **R Pandharipande**, *Maps, sheaves and K3 surfaces*, from “Lectures on geometry”, Clay Lect. Notes, Oxford Univ. Press, Oxford (2017) 159–185
- [38] **D Ranganathan**, *Logarithmic Gromov-Witten theory with expansions* [arXiv:1903.09006](https://arxiv.org/abs/1903.09006), 2019
- [39] **V Shende**, *Hilbert schemes of points on a locally planar curve and the Severi strata of its versal deformation*, Compos. Math. 148 (2012) 531–547
- [40] **N Takahashi**, *Curves in the complement of a smooth plane cubic whose normalizations are \mathbb{A}^1* [arXiv:alg-geom/9605007](https://arxiv.org/abs/alg-geom/9605007), 1996
- [41] **N Takahashi**, *On the multiplicity of reducible relative stable morphisms* [arXiv:1711.08173](https://arxiv.org/abs/1711.08173), 2017
- [42] **N Takahashi**, *Log mirror symmetry and local mirror symmetry*, Comm. Math. Phys. 220 (2001) 293–299
- [43] **Y Toda**, *Limit stable objects on Calabi-Yau 3-folds*, Duke Math. J. 149 (2009) 157–208

- [44] **Y Toda**, *Curve counting theories via stable objects I. DT/PT correspondence*, J. Amer. Math. Soc. 23 (2010) 1119–1157
- [45] **Y Toda**, *Stability conditions and curve counting invariants on Calabi-Yau 3-folds*, Kyoto J. Math. 52 (2012) 1–50
- [46] **C Voisin**, *A mathematical proof of a formula of Aspinwall and Morrison*, Compositio Math. 104 (1996) 135–151
- [47] **J Wise**, *Uniqueness of minimal morphisms of logarithmic schemes*, Algebr. Geom. 6 (2019) 50–63
- [48] **S-T Yau, E Zaslow**, *BPS states, string duality, and nodal curves on K3*, Nuclear Phys. B 471 (1996) 503–512

Department of Mathematics & Research Institute of Natural Sciences, Sookmyung Women’s University, Cheongpa-ro 47-gil 100, Youngsan-gu, Seoul 04310, Republic of Korea

Mathematics Institute, Zeeman Building, University of Warwick, Coventry CV4 7AL, UK

Department of Mathematics, MC-382, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

Department of Mathematics, Graduate School of Science, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima, 739-8526 JAPAN

jwchoi@sookmyung.ac.kr, michel.van-garrel@warwick.ac.uk, katz@math.uiuc.edu,
tkhsnbys@hiroshima-u.ac.jp