

*A constructive approach to a conjecture by
Voskresenskii*

Mathieu Florence & Michel van Garrel

Selecta Mathematica

New Series


ISSN 1022-1824

Volume 23

Number 4

Sel. Math. New Ser. (2017) 23:2899-2906

DOI 10.1007/s00029-017-0330-x



Your article is protected by copyright and all rights are held exclusively by Springer International Publishing. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".



A constructive approach to a conjecture by Voskresenskii

Mathieu Florence¹ · Michel van Garrel²

Published online: 12 May 2017
© Springer International Publishing 2017

Abstract Voskresenskii conjectured that stably rational tori are rational. Klyachko proved this assertion for a wide class of tori by general principles. We re-prove Klyachko's result by providing simple explicit birational isomorphisms, and elaborate on some links to torus-based cryptography.

Keywords Linear algebraic groups · Stable rationality · Rationality · Algebraic tori · Voskresenskii conjecture · Torus-based cryptography

Mathematics Subject Classification 14E08 · 14M20 · 14L15 · 14L30 · 14G50

Contents

1 Introduction	2900
2 Setup and statement of result	2900
3 Proof of the Theorem	2902
4 An application to Cryptography	2905
References	2906

The first author is partially supported by the French National Agency (Project GeoLie ANR-15-CE40-0012).

✉ Michel van Garrel
michel.van.garrel@uni-hamburg.de

Mathieu Florence
mathieu.florence@imj-prg.fr

¹ Institut de Mathématiques de Jussieu, Université Paris 6, Place Jussieu 4, 75005 Paris, France

² Fachbereich Mathematik, Universität Hamburg, Bundesstrasse 55, 20146 Hamburg, Germany

1 Introduction

Let k be an infinite field of any characteristic. We denote by \bar{k} an algebraic closure of k . A variety X over k is said to be *rational* if it is birational to a projective space \mathbb{P}_k^n . A strictly weaker notion is that of *stable rationality*.

Definition 1.1 Let X be a variety over k . X is said to be *stably rational* if $X \times_k \mathbb{P}_k^m$ is rational for some $m \geq 0$.

Let T be a linear (=affine) algebraic group over k . Then T is said to be an *algebraic torus* if, over an algebraic closure of k , it becomes isomorphic to a product of \mathbb{G}_m 's. A conjecture of Voskresenskii (see [5, p. 68]) states that a stably rational torus over k ought to be rational. This conjecture is widely open. A result of Klyachko ([2], see also [5, sec. 6.3]) gives a positive answer for a special type of stably rational tori, which we describe now (see Sect. 2 for a more detailed description).

Let A and B be étale k -algebras of coprime dimension over k . Denote by $\mathrm{GL}_1(A)$ the algebraic group of invertible elements in A . Let T be the quotient of $\mathrm{GL}_1(A \otimes_k B)$ by the subgroup generated by $\mathrm{GL}_1(A)$ and $\mathrm{GL}_1(B)$. Then T is a stably rational k -torus and Klyachko shows that it is in fact rational. However, his proof by general principles does not provide a simple explicit birational isomorphism from T to a projective space.

We remedy to this by re-proving Klyachko's result, constructing a simple birational map from T to a projective space. We expect our construction to generalize to the situation where A and B are *any* not necessarily commutative finite-dimensional k -algebras, of coprime dimension over k (in that case T is not necessarily a torus, or even an algebraic group).

In Sect. 4, we explore applications of our explicit birational maps to torus-based cryptography. Following the methods developed by Rubin-Silverberg in [4], we propose more general compression algorithms.

2 Setup and statement of result

Let k be an infinite field of any characteristic and let V be a finite-dimensional k -vector space. We start by recalling some k -schemes that are associated to V . The *affine space* of V , denoted by $\mathbb{A}(V)$, is defined as the functor

$$X \mapsto \mathbb{A}(V)(X) := V \otimes_k \Gamma(X, \mathcal{O}_X),$$

from k -schemes to sets. It is represented by the affine scheme $\mathrm{Spec}(\mathrm{Sym} V^*)$.

The *projective space* of V , denoted $\mathbb{P}(V)$, represents the (functor of) locally free submodules of rank one $N \subset V$, such that the quotient V/N is locally free. It is defined to be

$$\mathbb{P}(V) := (\mathbb{A}(V) - \{0\})/\mathbb{G}_m = \mathrm{Proj}(\mathrm{Sym} V^*).$$

Let A be a not necessarily commutative (unital) k -algebra of finite dimension. The linear algebraic group $\mathrm{GL}_1(A)$ is defined as the functor

$$X \mapsto \mathrm{GL}_1(A)(X) := (A \otimes_k \Gamma(X, \mathcal{O}_X))^\times,$$

from k -schemes to groups. It is represented by the closed subscheme of $\mathbb{A}(A \oplus A)$ given by the equation $xy = 1$. One has a canonical injective homomorphism of algebraic groups

$$1 \longrightarrow \mathbb{G}_m \longrightarrow \mathrm{GL}_1(A),$$

and can form the quotient

$$\mathrm{PGL}_1(A) := \mathrm{GL}_1(A)/\mathbb{G}_m,$$

which is a linear algebraic group.

For the remainder, assume that A is commutative.

Then, $\mathrm{GL}_1(A)$ is canonically isomorphic to the Weil restriction of scalars $\mathrm{Res}_{A/k}(\mathbb{G}_m)$. Let M be an A -module, which is locally free of finite rank. The projective space $\mathbb{P}(M)$ is defined over $\mathrm{Spec}(A)$. In this work, it shall be viewed as a k -variety, by Weil scalar restriction. More explicitly, we set

$$\mathbb{P}_A(M) := \mathrm{Res}_{A/k}(\mathbb{P}(M)).$$

Consider two finite-dimensional commutative k -algebras A and B . We have an exact sequence

$$1 \longrightarrow \mathbb{G}_m \xrightarrow{i} \mathrm{GL}_1(A) \times \mathrm{GL}_1(B) \xrightarrow{j} \mathrm{GL}_1(A \otimes_k B),$$

$$i(x) = (x, x^{-1}), \quad j(a, b) = a \otimes b.$$

Put $H(A, B) = \mathrm{im}(j)$. We will consider the quotient

$$Q(A, B) := \mathrm{GL}_1(A \otimes_k B)/H(A, B). \tag{2.1}$$

It follows from [5, section 6.1, Theorem 1] that $Q(A, B)$ is stably rational.

Recall that a k -algebra A is said to be étale if one of the two following equivalent conditions holds:

- $A \cong \prod_{i=1}^n k_i$, where the k_i are finite separable field extensions of k .
- $A \otimes_k \bar{k}$, as a \bar{k} -algebra, is isomorphic to a finite product of copies of \bar{k} .

The main result of this paper is to re-prove, in a constructive fashion, the following result.

Theorem 2.1 (Klyachko in [2], see also [5], section 6.3) *Let k be an infinite field of any characteristic and let A and B be two étale k -algebras of finite dimension. Assume that $\dim(A)$ and $\dim(B)$ are coprime. Then $Q(A, B)$ is k -rational.*

Note that the proof of Theorem 2.1 that we provide in Sect. 3 is via explicit birational isomorphisms, whereas Klyachko's original proof is by general principles. Recall that an algebraic k -torus of dimension d is a k -group scheme T such that

$$T \times_k \bar{k} \cong \mathbb{G}_{m,\bar{k}}^d.$$

The following conjecture states that for algebraic tori, stable rationality is equivalent to rationality.

Conjecture 2.2 (Voskresenskii, see [5], section 6.2) *Stably rational k -tori are k -rational.*

Theorem 2.1 thus provides a positive proof of Conjecture 2.2, in a particular case.

3 Proof of the Theorem

Let A and B be étale k -algebras of coprime dimensions (over k) a and b , respectively. Being invertible is an open condition, so that $\mathrm{GL}_1(A \otimes B)$ is a nonempty open subvariety of $\mathbb{A}(A \otimes B)$. Choose integers $0 < u \leq b$ and $0 < v \leq a$ such that

$$ua + vb = ab + 1.$$

This is possible since a and b are chosen to be coprime to each other. For a k -vector subspace $W \subset A \otimes_k B$, containing 1, denote by

$$\mathbb{P}_1(W) \subset \mathbb{P}(W)$$

the non-empty open subvariety consisting of lines directed by an invertible element of W .

Proposition 3.1 *There exist k -vector subspaces $U \in \mathrm{Gr}(u, B)(k)$ and $V \in \mathrm{Gr}(v, A)(k)$, both containing 1, such that the morphism below is a birational isomorphism:*

$$\begin{aligned} \phi_1 : \mathbb{P}_1(V \otimes_k B) \times \mathbb{P}_1(A \otimes_k U) &\dashrightarrow \mathbb{P}_1(A \otimes_k B) = \mathrm{PGL}_1(A \otimes_k B), \\ (x, y) &\longmapsto xy^{-1}. \end{aligned} \tag{3.1}$$

Proof in the case of fields. We first prove the assertion in the case that A and B are fields. Then $A \otimes_k B$ is a field as well, because a and b are coprime. Take arbitrary U and V as in the statement. We claim that ϕ_1 then is a birational isomorphism. Consider the fibers of ϕ_1 . An invertible k -rational point of $\mathrm{PGL}_1(A \otimes_k B)$ is given by the class of $t \in (A \otimes_k B)^\times$. The fiber over that class consists of (the projectivization of)

$$\{(x, y) \in (V \otimes_k B) \oplus (A \otimes_k U) \mid x = yt\},$$

where $(V \otimes_k B) \oplus (A \otimes_k U)$ is a vector k -space of dimension $vb + au = ab + 1$. Hence the equation $x = yt$ in $A \otimes_k B$ breaks down into a homogeneous linear system

of ab equations in $ab + 1$ variables. It follows that it has a non-trivial solution (x, y) over k . Since $A \otimes_k B$ is a field, both x and y are invertible. This shows that the fiber of ϕ_1 at t is non-empty, even isomorphic to a non-empty open of a projective space. But one may base-change from k to the function field K of $\text{PGL}_1(A \otimes_k B)$, and reproduce the previous arguments with K instead of k (note that K/k is purely transcendental, hence $A \otimes_k K$ and $B \otimes_k K$ are still fields). We thus get that the generic fiber of ϕ_1 is K -rational. But the source and target of ϕ_1 have the same dimension $ab - 1$. Hence, as asserted, ϕ_1 is a birational isomorphism. \square

Proof in the general case. It is a specialization argument as follows. We start by introducing the polynomial algebra (in $a + b$ variables)

$$\mathcal{K} := k[x_0, \dots, x_{a-1}, y_0, \dots, y_{b-1}],$$

and denote by $\tilde{\mathcal{K}}$ its field of fractions. Set

$$\mathcal{A} := K[T]/\langle T^a + x_{a-1}T^{a-1} + \dots + x_1T + x_0 \rangle$$

and

$$\mathcal{B} := K[T]/\langle T^b + y_{b-1}T^{b-1} + \dots + y_1T + y_0 \rangle,$$

and put

$$\tilde{\mathcal{A}} := \mathcal{A} \otimes_{\mathcal{K}} \tilde{\mathcal{K}}$$

as well as

$$\tilde{\mathcal{B}} := \mathcal{B} \otimes_{\mathcal{K}} \tilde{\mathcal{K}}.$$

Then $\tilde{\mathcal{A}}$ (resp. $\tilde{\mathcal{B}}$) is an étale $\tilde{\mathcal{K}}$ -algebra of degree a (resp. b). It is clearly a field. Pick $\tilde{\mathcal{K}}$ -subspaces $\tilde{\mathcal{U}} \in \text{Gr}(u, \tilde{\mathcal{B}})(\tilde{\mathcal{K}})$ and $\tilde{\mathcal{V}} \in \text{Gr}(v, \tilde{\mathcal{A}})(\tilde{\mathcal{K}})$, both containing 1. By what precedes, the $\tilde{\mathcal{K}}$ -morphism

$$\begin{aligned} \tilde{\Phi}_1 : \mathbb{P}_1(\tilde{\mathcal{V}} \otimes_{\tilde{\mathcal{K}}} \tilde{\mathcal{B}}) \times \mathbb{P}_1(\tilde{\mathcal{A}} \otimes_{\tilde{\mathcal{K}}} \tilde{\mathcal{U}}) &\dashrightarrow \mathbb{P}_1(\tilde{\mathcal{A}} \otimes_{\tilde{\mathcal{K}}} \tilde{\mathcal{B}}), \\ (x, y) &\longmapsto xy^{-1} \end{aligned}$$

is a birational isomorphism. Since all above schemes are of finite presentation over $\tilde{\mathcal{K}}$, they, as well as Φ_1 , are actually defined over a nonempty open subscheme of $\text{Spec}(\mathcal{K})$. More precisely, there exists a nonzero element $F \in \mathcal{K}$, such that, denoting by $\mathcal{K}_{(F)}$ the k -algebra obtained by inverting F in \mathcal{K} , the following holds:

- (a) The $\mathcal{K}_{(F)}$ -algebras $\mathcal{A}_{(F)}$ and $\mathcal{B}_{(F)}$ are étale.
- (b) The subspaces $\tilde{\mathcal{U}}$ and $\tilde{\mathcal{V}}$ are defined over $\mathcal{K}_{(F)}$, i.e., are given by elements $\mathcal{U} \in \text{Gr}(u, \mathcal{B})(\mathcal{K}_{(F)})$ and $\mathcal{V} \in \text{Gr}(v, \mathcal{A})(\mathcal{K}_{(F)})$, respectively.

(c) The $\mathcal{K}_{(F)}$ -morphism

$$\begin{aligned} \Phi_1 : \mathbb{P}_1(\mathcal{V} \otimes_{\mathcal{K}_{(F)}} \mathcal{B}_{(F)}) \times \mathbb{P}_1(\mathcal{A}_{(F)} \otimes_{\mathcal{K}_{(F)}} \mathcal{U}) &\dashrightarrow \mathbb{P}_1(\mathcal{A}_{(F)} \otimes_{\mathcal{K}_{(F)}} \mathcal{B}_{(F)}), \\ (x, y) &\longmapsto xy^{-1} \end{aligned}$$

is a birational isomorphism.

But the étale $\tilde{\mathcal{K}}$ -algebras $\tilde{\mathcal{A}}$ and $\tilde{\mathcal{B}}$ are *versal*, in the sense of [1, Definition 5.1, see also section 24.6]. Hence, there exists a k -morphism

$$\theta : \mathcal{K}_{(F)} \longrightarrow k$$

such that $\mathcal{A}_{(F)} \otimes_{\theta} k$ is isomorphic to A (resp. such that $\mathcal{B}_{(F)} \otimes_{\theta} k$ is isomorphic to B). Put $V := \mathcal{V} \otimes_{\theta} k$ and $U := \mathcal{U} \otimes_{\theta} k$. Then U (resp. V) belongs to $\text{Gr}(u, B)(k)$ (resp. to $\text{Gr}(v, A)(k)$), and the specialization of Φ_1 via θ yields the birational isomorphism ϕ_1 . This finishes the proof of Proposition 3.1. \square

Note that $\mathbb{P}_1(V \otimes_k B) \times \mathbb{P}_1(A \otimes_k U)$ is open in $\mathbb{P}(V \otimes_k B) \times \mathbb{P}(A \otimes_k U)$, and that $\mathbb{P}_1(A \otimes_k B)$ is open in $\mathbb{P}(A \otimes_k B)$. Hence the map ϕ_1 of (3.1) extends to a birational isomorphism

$$\phi : \mathbb{P}(V \otimes_k B) \times \mathbb{P}(A \otimes_k U) \dashrightarrow \mathbb{P}(A \otimes_k B). \tag{3.2}$$

Generically, $G := \text{GL}_1(A)/\mathbb{G}_m \times \text{GL}_1(B)/\mathbb{G}_m$ acts freely on both sides of (3.2). We have the identifications as birational quotients:

$$\begin{aligned} \mathbb{P}(V \otimes_k B) / (\text{GL}_1(B)/\mathbb{G}_m) &\equiv (V \otimes_k B / \mathbb{G}_m) / (\text{GL}_1(B)/\mathbb{G}_m) \\ &\equiv (V \otimes_k B) / \text{GL}_1(B) \equiv \mathbb{P}_B(V \otimes_k B). \end{aligned}$$

Since the action of $\text{GL}_1(B)/\mathbb{G}_m$ on $V \otimes_k B / \mathbb{G}_m$ is generically free, we conclude that $\dim \mathbb{P}_B(V \otimes_k B) = vb - b$. Similarly,

$$\mathbb{P}(A \otimes_k U) / \text{GL}_1(A) \equiv \mathbb{P}_A(A \otimes_k U)$$

is of dimension $au - a$.

On the right hand side of the map of (3.2), we take the following birational quotient:

$$\mathbb{P}(A \otimes_k B) / G \equiv (A \otimes_k B / \mathbb{G}_m) / G.$$

As G acts generically freely, the dimension of this quotient is $ab - a - b + 1$. For an A -module M , recall from Sect. 2 that we defined $\mathbb{P}_A(M)$ to be the Weil scalar restriction $\text{Res}_{A/k}(\mathbb{P}(M))$.

Lemma 3.2 *The map ϕ of (3.2) induces a birational isomorphism*

$$\bar{\phi} : \mathbb{P}_B(V \otimes_k B) \times \mathbb{P}_A(A \otimes_k U) \dashrightarrow \mathbb{P}(A \otimes_k B) / \text{GL}_1(A) \times \text{GL}_1(B).$$

Proof The dimensions of both quotients agree. Since the map is a birational isomorphism before taking the quotient, we only need to show that it descends to the quotient. But that is clear since the map is given by taking the inverse and multiplication. \square

Finally, note that $\mathbb{P}(A \otimes_k B)/GL_1(A) \times GL_1(B)$ is birational to $Q(A, B)$. This then completes the proof of Theorem 2.1, as both $\mathbb{P}_B(V \otimes_k B)$ and $\mathbb{P}_A(A \otimes_k U)$ are rational.

4 An application to Cryptography

Our explicit birational maps open up some new venues for torus-based cryptography. Using finite cyclic groups for public key encryption is an old idea, cf. [3, chapter 8]. Rubin-Silverberg in [4] suggested using rational algebraic tori defined over finite fields. The advantage is in term of computational gain. Representing most elements of the torus as elements of an affine space over a finite field yields efficiency gains in the transmitted information. Let q be a prime power and choose $n \geq 1$ to be a square-free integer. If $\mathbb{F}_q \subseteq L \subsetneq \mathbb{F}_{q^n}$ is an intermediate field, recall that there is a norm map

$$N_{\mathbb{F}_{q^n}/L} : \text{Res}_{\mathbb{F}_{q^n}/\mathbb{F}_q} \mathbb{G}_m \rightarrow \text{Res}_{L/\mathbb{F}_q} \mathbb{G}_m.$$

Following [4], consider

$$T_n := \bigcap_{\mathbb{F}_q \subseteq L \subsetneq \mathbb{F}_{q^n}} \ker \left(N_{\mathbb{F}_{q^n}/L} \right) \text{ and } G_{q,n} := T_n(\mathbb{F}_q).$$

For encryption purposes, $G_{q,n}$ is the cryptographically most significant part of $\mathbb{F}_{q^n}^\times$ and $G_{q,n}$, albeit smaller, inherits the security of $\mathbb{F}_{q^n}^\times$. See [4] for more details. $G_{q,n}$ is a torus over \mathbb{F}_q of dimension $\phi(n)$, where ϕ denotes Euler's phi function. Assuming that it is rational, one then would like to (computationally) compress elements of $G_{q,n}$ via a *compression map* (birational map)

$$f : G_{q,n} \dashrightarrow \mathbb{F}_q^{\phi(n)}$$

that has an efficiently computable inverse j . Since $G_{q,n} < \mathbb{F}_{q^n}^\times$, the latter being of dimension n over \mathbb{F}_q , sending $f(x)$ instead of $x \in G_{q,n}$ yields an efficiency gain (in bits) of $n/\phi(n)$. Based on this idea, Rubin-Silverberg introduce two compression algorithms inducing efficient public key cryptosystems that they name \mathbb{T}_2 and CEILIDH. They also explain how to extend their algorithms to all $G_{q,n}$, provided that a compression map is known. Note that the encryption is restricted to the open part of $G_{q,n}$ where f and j are mutually inverse. This part is large if q is large, see the discussion in [4]. Moreover, they limit their discussion to when n is the product of up to two distinct primes. In particular, they consider $n = 2$ for \mathbb{T}_2 and $n = 6$ for CEILIDH (to yield secure encryption, q should be large). If n is the product of at least three primes, it is not known, though conjectured by Voskresenskii, that $G_{q,n}$ is rational.

\mathbb{T}_2 and CEILIDH are based on explicit birational compression maps that Rubin-Silverberg construct from Galois extensions. They rely on choosing generators for

these extensions. Our setting extends the groups beyond $G_{q,n}$ and does not rely on the extension being Galois, nor on choosing generators.

For the remainder, let $A = \mathbb{F}_{q^a}$ and $B = \mathbb{F}_{q^b}$, where a and b are coprime. Our birational decomposition map

$$\phi(A, B) : \mathbb{P}_B(V \otimes_{\mathbb{F}_q} B) \times \mathbb{P}_A(A \otimes_{\mathbb{F}_q} U) \dashrightarrow Q(A, B)$$

solely depends on the choice of the \mathbb{F}_q -vector subspaces $U \in \text{Gr}(u, B)(\mathbb{F}_q)$ and $V \in \text{Gr}(v, A)(\mathbb{F}_q)$ of Proposition 3.1. Note that, though Theorem 2.1 is a priori stated for infinite fields, it is easy to see that it actually holds for k finite, when A and B are fields. Furthermore, since a and b are coprime to each other, $A \otimes_{\mathbb{F}_q} B = \mathbb{F}_{q^{ab}}$ and

$$Q(A, B)(\mathbb{F}_q) = \mathbb{F}_{q^{ab}}^\times / \langle \mathbb{F}_{q^a}^\times, \mathbb{F}_{q^b}^\times \rangle,$$

where, cf. (2.1), $\langle \mathbb{F}_{q^a}^\times, \mathbb{F}_{q^b}^\times \rangle = H(\mathbb{F}_{q^a}, \mathbb{F}_{q^b})(\mathbb{F}_q)$ is the subgroup generated by $\mathbb{F}_{q^a}^\times$ and $\mathbb{F}_{q^b}^\times$. If in addition a and b are distinct primes (or $b = 1$ and a is prime), then

$$Q(A, B)(\mathbb{F}_q) \cong Q_{q,ab},$$

which is the case developed in [4]. Note that our compression maps differ, and work for all choices of primes a and b .

In order to have a computationally efficient extension of Rubin-Silverberg's algorithms to $Q(A, B)$, two conditions must be satisfied. Firstly, the ratio $ab/\phi(ab)$ should be large. Second and most crucially, the \mathbb{F}_q -vector subspaces U and V should be chosen such that the birational inverse of $\phi(A, B)$ is computed fast. As explained in the proof of Proposition 3.1, calculating this inverse is obtained through solving linear equations. Suitable choices of U and V will lead to computationally efficient algorithms. We leave the specifics of implementation to future considerations.

Acknowledgements A large part of this work was accomplished while the first author was visiting the Korea Institute for Advanced Study (KIAS), where the second author at the time was a research fellow. The authors thank KIAS for the hospitality and excellent research environment. The comments of the referee greatly enhanced the quality of the paper. The authors are particularly thankful for the connection made with cryptography.

References

1. Garibaldi, S., Merkurjev, A., Serre, J.-P.: Cohomological Invariants in Galois Cohomology. University Lecture Series, vol. 28. AMS (2003)
2. Klyachko, A.A.: On Rationality of Tori with a Cyclic Splitting Field. Arithmetic and Geometry of Varieties. Kuibyshev University Press, Kuibyshev (1988). **(Russian)**
3. Menezes, A.J., van Oorschot, P.C., Vanstone, S.A.: Handbook of Applied Cryptography. CRC Press, Boca Raton (1997)
4. Rubin, K., Silverberg, A.: Compression in finite fields and torus-based cryptography. SIAM J. Comput. **37**(5), 1401–1428 (2008)
5. Voskresenskii, V.E.: Algebraic Groups and Their Birational Invariants. Translations of Mathematical Monographs. AMS, Providence (1998)