

EXAMPLE SHEET 3

All rings are commutative with a 1 unless stated otherwise.

Matsushima  
14 H pg 92

1. A chain of prime ideals is maximal if it is not a proper subset of another chain of primes. Prove that all maximal chains of prime ideals in a finitely generated  $k$ -algebra  $T$  which is an integral domain are of the same length, and that  $\text{ht}P + \dim T/P = \dim T$  for any prime ideal  $P$  of  $T$ .

2. Give an example of a finitely generated algebra  $T$  with a prime ideal  $P$  for which  $\text{ht}P + \dim T/P < \dim T$ .

3. Let  $R$  be a Noetherian regular local ring. Show that  $R[[X]]$  is a regular local ring of dimension  $\dim R + 1$ . Deduce that if  $k$  is a field then  $k[[X_1, \dots, X_n]]$  of formal power series in  $n$  indeterminates is a regular local ring of dimension  $n$ .

Nakayama

4. For a not necessarily commutative ring  $R$  show that the following are equivalent for a right ideal  $I$ : (i)  $I \leq \text{Jac}R$ ; (ii) if  $M$  is a finitely generated  $R$ -module and a submodule  $N$  satisfies  $N + MI = M$  then  $N = M$ ; (iii) the set of elements  $1 + x$  for  $x \in I$  form a subgroup of the unit group of  $R$ .

5. For a not necessarily commutative ring  $R$  show that if an  $R$ -module  $M$  is a sum of simple submodules then  $M$  may be expressed as the direct sum of some simple submodules.

6. Let  $R$  be a semisimple right Artinian ring and  $M$  be an Artinian right  $R$ -module. Show that  $M$  is a direct sum of finitely many simple  $R$ -modules.

Use AW.

7. Let  $R$  be a  $k$ -algebra where  $k$  is an algebraically closed field, and suppose that  $R$  is finite dimensional as a  $k$ -vector space and semisimple. Define a Lie bracket on  $R$  by  $[x, y] = xy - yx$ . Show that the  $k$  vector space dimension of  $R/[R, R]$  is equal to the number of isomorphism classes of simple right  $R$ -modules.

8. Let  $k$  be a field of characteristic  $p > 0$  and let  $G$  be a finite group of order a power of  $p$ . Show that the augmentation ideal of  $kG$  (the kernel of the ring homomorphism from  $kG$  to  $k$  sends each  $g$  to 1) is nilpotent and that up to isomorphism the only simple module of  $kG$  is the trivial module, one dimensional as a  $k$  vector space.

9. Let  $G = S_3$  and let  $k$  be a field of characteristic 2. Describe the simple modules, the socle series and the Jacobson radical of  $kG$ .

10. Show that a ring  $R$  with an exhaustive and separated filtration is an integral domain if the associated graded ring  $\text{gr}R$  is an integral domain. Assume that the filtration of  $R$  is

positive and show that  $R$  is Noetherian if  $\text{gr}R$  is Noetherian. Is the same true for negative filtrations, for example the  $P$ -adic filtration of  $R$  where  $P$  is a prime ideal?

11. Let  $R$  be a Noetherian ring with ideal  $I$ . Show that the Rees ring of  $R$  with respect to the  $I$ -adic filtration is Noetherian. Let  $M$  be a finitely generated  $R$ -module. A filtration of  $M$  with respect to the  $I$ -adic filtration of  $R$  is said to be *good* if its Rees module  $\text{Rees}(M)$  is a Noetherian  $\text{Rees}(R)$ -module. Show that this is equivalent to it being *stable* (i.e there is some  $J$  such that  $M_{-j} = I^j M_{-j}$  for all  $j > 0$ ).

12. (Artin, Rees) Let  $R$  be a Noetherian ring, and let  $I$  be an ideal. Let  $M$  be a finitely generated  $R$ -module with submodule  $N$ . Show that there exists  $r \geq 0$  such that  $N \cap I^a M = I^{a-r}(N \cap I^r M)$  for  $a \geq r$ .

13. (Krull) Let  $R$  be a Noetherian local ring, and let  $I$  be a proper ideal. Let  $M$  be a finitely generated  $R$ -module. Show that the intersection of all the submodules  $I^n M$  is zero. In particular the intersection of all ideals  $I^n$  is zero.

14. Let  $R$  be a Noetherian ring and let  $I$  be an ideal. Let  $S = 1 + I$ . Show that the kernel of the canonical map from  $R$  to  $S^{-1}R$  is the intersection of the positive powers of  $I$ .

brookes@dpnms.cam.ac.uk