

NONLINEAR SCHRÖDINGER EVOLUTIONS FROM LOW REGULARITY INITIAL DATA

J. Colliander

University of Toronto

Warwick, June 2009

- 1 CUBIC NLS ON \mathbb{R}^2
- 2 HIGH-LOW FOURIER TRUNCATION
- 3 BILINEAR STRICHARTZ ESTIMATE
- 4 THE I -METHOD OF ALMOST CONSERVATION

1. CUBIC NLS INITIAL VALUE PROBLEM ON \mathbb{R}^2

1. CUBIC NLS INITIAL VALUE PROBLEM ON \mathbb{R}^2

We consider the initial value problems:

$$\begin{cases} (i\partial_t + \Delta)u = \pm|u|^2u \\ u(0, x) = u_0(x). \end{cases} \quad (NLS_3^\pm(\mathbb{R}^2))$$

The + case is called **defocusing**; - is **focusing**. NLS_3^\pm is ubiquitous in physics. The solution has a dilation symmetry

$$u^\lambda(\tau, y) = \lambda^{-1}u(\lambda^{-2}\tau, \lambda^{-1}y).$$


which is invariant in $L^2(\mathbb{R}^2)$. This problem is L^2 -critical.

(This talk mostly addresses the defocusing case.)

TIME INVARIANT QUANTITIES

$$\text{Mass} = \int_{\mathbb{R}^d} |u(t, x)|^2 dx.$$

$$\text{Momentum} = 2\Im \int_{\mathbb{R}^2} \bar{u}(t) \nabla u(t) dx.$$

$$\text{Energy} = H[u(t)] = \frac{1}{2} \int_{\mathbb{R}^2} |\nabla u(t)|^2 dx \pm \frac{1}{2} |u(t)|^4 dx.$$


Hamiltonian

kinetic

potential

- Mass is L^2 ; Momentum is close to $H^{1/2}$; Energy involves H^1 .
- Dynamics on a sphere in L^2 ; **focusing/defocusing** energy.
- Local conservation laws express **how** quantity is conserved: e.g., $\partial_t |u|^2 = \nabla \cdot 2\Im(\bar{u} \nabla u)$. Frequency Localizations?

LINEAR SCHRÖDINGER PROPAGATOR AND ESTIMATES

The solution of the linear Schrödinger initial value problem

$$\begin{cases} (i\partial_t + \Delta)u = 0 \\ u(0, x) = u_0(x). \end{cases} \quad (LS(\mathbb{R}^d))$$

is denoted $u(t, x) = e^{it\Delta}u_0$. The solution can be given explicitly

- Fourier Multiplier Representation:

$$e^{it\Delta}u_0(x) = c_\pi \int_{\mathbb{R}^d} e^{ix \cdot \xi} e^{-it|\xi|^2} \widehat{u}_0(\xi) d\xi.$$

- Convolution Representation:

$$e^{it\Delta}u_0(x) = c_\pi \frac{1}{(it)^{d/2}} \int_{\mathbb{R}^d} e^{i\frac{|x-y|^2}{4t}} u_0(y) dy.$$

LINEAR SCHRÖDINGER PROPAGATOR AND ESTIMATES

The solution of the linear Schrödinger initial value problem

$$\begin{cases} (i\partial_t + \Delta)u = 0 \\ u(0, x) = u_0(x). \end{cases} \quad (LS(\mathbb{R}^d))$$

is denoted $u(t, x) = e^{it\Delta}u_0$. The solution can be given explicitly

- **Fourier Multiplier Representation:**

$$e^{it\Delta}u_0(x) = c_\pi \int_{\mathbb{R}^d} e^{ix \cdot \xi} \underbrace{e^{-it|\xi|^2}}_{\text{Modulus 1 Multiplier}} \widehat{u}_0(\xi) d\xi.$$

- Convolution Representation:

$$e^{it\Delta}u_0(x) = c_\pi \frac{1}{(it)^{d/2}} \int_{\mathbb{R}^d} e^{i\frac{|x-y|^2}{4t}} u_0(y) dy.$$

LINEAR SCHRÖDINGER PROPAGATOR

The solution of the linear Schrödinger initial value problem

$$\begin{cases} (i\partial_t + \Delta)u = 0 \\ u(0, x) = u_0(x). \end{cases} \quad (LS(\mathbb{R}^d))$$

is denoted $u(t, x) = e^{it\Delta}u_0$. The solution can be given explicitly

- Fourier Multiplier Representation:

$$e^{it\Delta}u_0(x) = c_\pi \int_{\mathbb{R}^d} e^{ix \cdot \xi} e^{-it|\xi|^2} \widehat{u}_0(\xi) d\xi.$$

- Convolution Representation:

$$e^{it\Delta}u_0(x) = c_\pi \frac{1}{(it)^{d/2}} \int_{\mathbb{R}^d} e^{i\frac{|x-y|^2}{4t}} u_0(y) dy.$$

 Time Decay

ESTIMATES FOR LINEAR SCHRÖDINGER PROPAGATOR

- Fourier Multiplier Representation \implies Unitary in H^s :

$$\|D_x^s e^{it\Delta} u_0\|_{L_x^2} = \|D_x^s u_0\|_{L_x^2}.$$

- Convolution Representation \implies Dispersive estimate:

$$\|e^{it\Delta} u_0\|_{L_x^\infty} \leq \frac{C}{t^{d/2}} \|u_0\|_{L_x^1}.$$

- Spacetime estimates? **Strichartz estimates** hold, for example,

$$\|e^{it\Delta} u_0\|_{L^4(\mathbb{R}_t \times \mathbb{R}_x^2)} \leq C \|u_0\|_{L^2(\mathbb{R}_x^2)}.$$

(Strichartz estimates linked to Fourier restriction phenomena.)

LOCAL-IN-TIME THEORY FOR $NLS_3^\pm(\mathbb{R}^2)$

- $\forall u_0 \in L^2(\mathbb{R}^2) \exists T_{lwp}(u_0)$ determined by

$$\|e^{it\Delta} u_0\|_{L_{tx}^4([0, T_{lwp}] \times \mathbb{R}^2)} < \frac{1}{100} \text{ such that}$$

\exists unique $u \in C([0, T_{lwp}]; L^2) \cap L_{tx}^4([0, T_{lwp}] \times \mathbb{R}^2)$ solving $NLS_3^+(\mathbb{R}^2)$.

- $\forall u_0 \in H^s(\mathbb{R}^2), s > 0, T_{lwp} \sim \|u_0\|_{H^s}^{-\frac{2}{s}}$ and regularity persists: $u \in C([0, T_{lwp}]; H^s(\mathbb{R}^2))$.
- Define the **maximal forward existence time** $T^*(u_0)$ by

$$\|u\|_{L_{tx}^4([0, T^* - \delta] \times \mathbb{R}^2)} < \infty$$

for all $\delta > 0$ but diverges to ∞ as $\delta \searrow 0$.

- \exists **small data scattering threshold** $\mu_0 > 0$

$$\|u_0\|_{L^2} < \mu_0 \implies \|u\|_{L_{tx}^4(\mathbb{R} \times \mathbb{R}^2)} < 2\mu_0.$$

GLOBAL-IN-TIME THEORY?

What is the ultimate fate of the local-in-time solutions?

L^2 -critical Scattering Conjecture:

$L^2 \ni u_0 \mapsto u$ solving $NLS_3^+(\mathbb{R}^2)$ is global-in-time and

$$\|u\|_{L_{t,x}^4} < A(u_0) < \infty.$$

Moreover, $\exists u_{\pm} \in L^2(\mathbb{R}^2)$ such that

$$\lim_{t \rightarrow \pm\infty} \|e^{\pm it\Delta} u_{\pm} - u(t)\|_{L^2(\mathbb{R}^2)} = 0.$$

Same statement for focusing $NLS_3^-(\mathbb{R}^2)$ if $\|u_0\|_{L^2} < \|Q\|_{L^2}$.

Remarks:

- Known for small data $\|u_0\|_{L^2(\mathbb{R}^2)} < \mu_0$.
- Known for **large radial data** [Killip-Tao-Visan 07].

$NLS_3^\pm(\mathbb{R}^2)$: PRESENT STATUS FOR GENERAL DATA

regularity	idea	reference
$s > \frac{2}{3}$ $s > \frac{4}{7}$	high/low frequency decomposition $H(lu)$	[Bourgain98] [CKSTT02]
$s > \frac{1}{2}$ $s \geq \frac{1}{2}$ $s > \frac{2}{5}$	resonant cut of 2nd energy $H(lu)$ & Interaction Morawetz $H(lu)$ & Interaction l -Morawetz	[CKSTT07] [Fang-Grillakis05] [CGTz07]
$s > \frac{1}{3}$	resonant cut & l -Morawetz	[C-Roy08]
$s > 0?$		

- Morawetz-based arguments are only for defocusing case.
- Focusing results assume $\|u_0\|_{L^2} < \|Q\|_{L^2}$.
- Unify theory of focusing-under-ground-state and defocusing?

2. BOURGAIN'S HIGH-LOW FOURIER TRUNCATION

2. BOURGAIN'S HIGH-LOW FOURIER TRUNCATION

IMRN International Mathematics Research Notices
1998, No. 5

Refinements of Strichartz' Inequality and Applications to 2D-NLS with Critical Nonlinearity

J. Bourgain

Summary

Consider the 2D IVP

$$\begin{cases} iu_t + \Delta u + \lambda|u|^2u = 0 \\ u(0) = \varphi \in L^2(\mathbb{R}^2). \end{cases} \quad (t)$$

The theory on the Cauchy problem asserts a unique maximal solution

$$u \in C([0, T], T^*[L^2(\mathbb{R}^2)]) \cap L^4([0, T], T^*[L^4(\mathbb{R}^2)])$$

2. BOURGAIN'S HIGH-LOW FOURIER TRUNCATION

Consider the Cauchy problem for defocusing cubic NLS on \mathbb{R}^2 :

$$\begin{cases} (i\partial_t + \Delta)u = +|u|^2u \\ u(0, x) = \phi_0(x). \end{cases} \quad (NLS_3^+(\mathbb{R}^2))$$

We describe the first result to give global well-posedness below H^1 .

- $NLS_3^+(\mathbb{R}^2)$ is GWP in H^s for $s > \frac{2}{3}$ [Bourgain 98].
- First use of Bilinear Strichartz estimate was in this proof.
- Proof cuts solution into low and high frequency parts.
- For $u_0 \in H^s$, $s > \frac{2}{3}$, Proof gives (and crucially exploits),

$$u(t) - e^{it\Delta}\phi_0 \in H^1(\mathbb{R}_x^2).$$

SETTING UP; DECOMPOSING DATA

- Fix a large target time T .
- Let $N = N(T)$ be large to be determined.
- Decompose the initial data:

$$\phi_0 = \phi_{low} + \phi_{high}$$

where

$$\phi_{low}(x) = \int_{|\xi| < N} e^{ix \cdot \xi} \widehat{\phi_0}(\xi) d\xi.$$

- Our plan is to evolve:

$$\phi_0 = \phi_{low} + \phi_{high}$$

$$u(t) = u_{low}(t) + u_{high}(t).$$

SETTING UP; DECOMPOSING DATA

Low Frequency Data Size:

- Kinetic Energy:

$$\begin{aligned}\|\nabla \phi_{low}\|_{L^2}^2 &= \int_{|\xi| < N} |\xi|^2 |\widehat{\phi_0}(\xi)|^2 dx \\ &= \int_{|\xi| < N} |\xi|^{2(1-s)} |\xi|^{2s} |\widehat{\phi_0}(\xi)|^2 dx \\ &\leq N^{2(1-s)} \|\phi_0\|_{H^s}^2 \leq C_0 N^{2(1-s)}.\end{aligned}$$

- Potential Energy: $\|\phi_{low}\|_{L_x^4} \leq \|\phi_{low}\|_{L^2}^{1/2} \|\nabla \phi_{low}\|_{L^2}^{1/2}$
 $\implies H[\phi_{low}] \leq CN^{2(1-s)}.$

High Frequency Data Size:

$$\|\phi_{high}\|_{L^2} \leq C_0 N^{-s}, \quad \|\phi_{high}\|_{H^s} \leq C_0.$$

LWP OF LOW FREQUENCY EVOLUTION ALONG NLS

The NLS Cauchy Problem for the low frequency data

$$\begin{cases} (i\partial_t + \Delta)u_{low} = +|u_{low}|^2 u_{low} \\ u_{low}(0, x) = \phi_{low}(x) \end{cases}$$

is well-posed on $[0, T_{lwp}]$ with $T_{lwp} \sim \|\phi_{low}\|_{H^1}^{-2} \sim N^{-2(1-s)}$.

We obtain, as a consequence of the local theory, that

$$\|u_{low}\|_{L^4_{[0, T_{lwp}], x}} \leq \frac{1}{100}.$$

LWP OF HIGH FREQUENCY EVOLUTION ALONG DE

The NLS Cauchy Problem for the low frequency data

$$\begin{cases} (i\partial_t + \Delta)u_{high} = +2|u_{low}|^2u_{high} + \text{similar} + |u_{high}|^2u_{high} \\ u_{high}(0, x) = \phi_{high}(x) \end{cases}$$

is also well-posed on $[0, T_{lwp}]$.

Remark: The LWP lifetime of *NLS* evolution of u_{low} AND the LWP lifetime of the *DE* evolution of u_{high} are controlled by $\|u_{low}(0)\|_{H^1}$.

EXTRA SMOOTHING OF NONLINEAR DUHAMEL TERM

The high frequency evolution may be written

$$u_{high}(t) = e^{it\Delta} u_{high} + w.$$

The local theory gives $\|w(t)\|_{L^2} \lesssim N^{-s}$. Moreover, due to smoothing (obtained via bilinear Strichartz), we have that

$$w \in H^1, \quad \|w(t)\|_{H^1} \lesssim N^{1-2s+}. \quad (\text{SMOOTH!})$$

Let's postpone the proof of (SMOOTH!).

NONLINEAR HIGH FREQUENCY TERM HIDING STEP!

- $\forall t \in [0, T_{lwp}]$, we have

$$u(t) = u_{low}(t) + e^{it\Delta} \phi_{high} + w(t).$$

- At time T_{lwp} , we define data for the progressive scheme:

$$u(T_{lwp}) = u_{low}(T_{lwp}) + w(T_{lwp}) + e^{iT_{lwp}\Delta} \phi_{high}.$$

$$u(t) = u_{low}^{(2)}(t) + u_{high}^{(2)}(t)$$

for $t > T_{lwp}$.

HAMILTONIAN INCREMENT: $\phi_{low}(0) \mapsto u_{low}^{(2)}(T_{lwp})$

The Hamiltonian increment due to $w(T_{lwp})$ being added to low frequency evolution can be calculated. Indeed, by Taylor expansion, using the bound (SMOOTH!) and energy conservation of u_{low} evolution, we have using

$$\begin{aligned} H[u_{low}^{(2)}(T_{lwp})] &= H[u_{low}(0)] + (H[u_{low}(T_{lwp}) + w(T_{lwp})] - H[u_{low}(T_{lwp})]) \\ &\sim N^{2(1-s)} + N^{2-3s} \sim N^{2(1-s)}. \end{aligned}$$

Moreover, we can accumulate N^s increments of size N^{2-3s} before we double the size $N^{2(1-s)}$ of the Hamiltonian. During the iteration, Hamiltonian of “low frequency” pieces remains of size $\lesssim N^{2(1-s)}$ so the LWP steps are of uniform size $N^{-2(1-s)}$. We advance the solution on a time interval of size:

$$N^s N^{-2(1-s)} = N^{-2+3s}.$$

For $s > \frac{2}{3}$, we can choose N to go past target time T .

HOW DO WE PROVE (SMOOTH!)?

Bourgain's Bilinear Strichartz Estimate: For (dyadic) $N \leq L$

$$\|e^{it\Delta} f_L e^{it\Delta} g_N\|_{L_{t,x}^2} \leq \frac{N^{\frac{2-1}{2}}}{L^{\frac{1}{2}}} \|f_L\|_{L_x^2} \|g_N\|_{L_x^2}.$$

COROLLARY

For $s \geq \frac{1}{2}$

$$\begin{aligned} \|D_x^s(u_1 u_2)\|_{L_{[0,\delta],x}^2} &\leq C(\|u_1\|_{X_{[0,\delta]}^{s,1/2+}} \|u_2\|_{X_{[0,\delta]}^{0,1/2+}} \\ &\quad + \|u_1\|_{X_{[0,\delta]}^{1/2,1/2+}} \|u_2\|_{X_{[0,\delta]}^{s-1/2,1/2+}}). \end{aligned}$$

Thus, the Bilinear Estimate allows us move half a derivative off the high frequency part and instead onto of the low frequency part.

TREATMENT OF A TYPICAL TERM IN w

- Using the controls we have on u_{low} , u_{high} from the local theory on $[0, T_{low}]$, we want to prove for

$$w = \int_0^t e^{i(t-t')\Delta} |u_{low}|^2 u_{high}(t') dt'$$

that $\sup_{t \in [0, T_{low}]} \|\nabla w\|_{L^2} < N^{1-2S+}$.

- By Sobolev embedding, we have

$$\|w\|_{L_{[0, T_{low}]}^\infty H^1} \leq \|w\|_{X_{[0, T_{low}]}^{1, 1/2+}}$$

- The mapping $f \mapsto \int_0^t e^{i(t-t')\Delta} f$ is formally $f \mapsto (i\partial_t + \Delta)^{-1} f$ which, due to time localization, is essentially $\widehat{f} \mapsto \langle \tau + |\xi|^2 \rangle \widehat{f}$. It suffices to control $\|D_x |u_{low}|^2 u_{high}\|_{X^{0, -1/2+}}$. Proceed by duality....

TREATMENT OF A TYPICAL TERM IN w

$$\begin{aligned} \|w\|_{L^\infty_{[0, T_{low}]} H^1} &\leq \sup_{\|g\|_{X^{0,1/2-}} \leq 1} \langle g, D_x(|u_{low}|^2 u_{high}) \rangle. \\ &\lesssim \sup_g \langle g D_x u_{low}, u_{low} u_{high} \rangle + \sup_g \langle g u_{low}, D_x(u_{low} u_{high}) \rangle \\ &= \text{easier} + \sup_g \langle D_x^{1/2}(g u_{low}), D_x^{1/2}(u_{low} u_{high}) \rangle. \end{aligned}$$

The corollary and the available bounds then give (SMOOTH!).

3. BOURGAIN'S BILINEAR STRICHARTZ ESTIMATE

3. BOURGAIN'S BILINEAR STRICHARTZ ESTIMATE

- Recall the Strichartz estimate for $(i\partial_t + \Delta)$ on \mathbb{R}^2 :

$$\|e^{it\Delta} u_0\|_{L^4(\mathbb{R}_t \times \mathbb{R}_x^2)} \leq C \|u_0\|_{L^2(\mathbb{R}_x^2)}.$$

- We can view this trivially as a bilinear estimate by writing

$$\|e^{it\Delta} u_0 e^{it\Delta} v_0\|_{L^2(\mathbb{R}_t \times \mathbb{R}_x^2)} \leq C \|u_0\|_{L^2(\mathbb{R}_x^2)} \|v_0\|_{L^2(\mathbb{R}_x^2)}.$$

- Bourgain refined this trivial bilinear estimate for functions having certain Fourier support properties.

BOURGAIN'S BILINEAR STRICHARTZ ESTIMATE

Shrinks Constant

THEOREM

For (dyadic) $N \leq L$ and for $x \in \mathbb{R}^2$,

$$\|e^{it\Delta} f_L e^{it\Delta} g_N\|_{L_{t,x}^2} \leq \frac{N^{\frac{1}{2}}}{L^{\frac{1}{2}}} \|f_L\|_{L_x^2} \|g_N\|_{L_x^2}.$$

- Here $\text{spt}(\widehat{f}_L) \subset \{|\xi| \sim L\}$, g_N similar.
- Observe that $\sqrt{\frac{N}{L}} \ll 1$ when $N \ll L$.

3. BOURGAIN'S PROOF

B98:IMRN

Proof. Since the standard Strichartz inequality yields (112) without the

$$\left(\frac{M_1}{M_2}\right)^{\frac{1}{2}} \text{-factor,}$$

we may assume $M_2 \gg M_1$.

Writing

$$(e^{it\Delta}\psi_1)(e^{it\Delta}\psi_2) = \int \widehat{\psi}_1(\xi_1)\widehat{\psi}_2(\xi_2)e^{i[(\xi_1+\xi_2)\cdot x + (|\xi_1|^2+|\xi_2|^2)t]} d\xi_1 d\xi_2,$$

it follows from Parseval's identity and Cauchy-Schwarz that

$$\begin{aligned} \|(e^{it\Delta}\psi_1)(e^{it\Delta}\psi_2)\|_2^2 &= \int d\xi d\lambda \left| \int \widehat{\psi}_1(\xi_1)\widehat{\psi}_2(\xi - \xi_1)\delta_0(|\xi_1|^2 + |\xi - \xi_1|^2 - \lambda) d\xi_1 \right|^2 \\ &\leq \|\psi_1\|_2^2 \|\psi_2\|_2^2 \left[\sup_{\lambda, |\xi| \sim M_2} \text{mes}_{(1)}[|\xi_1|, |\xi_1| \sim M_1 \right. \\ &\quad \left. \text{and } |\xi_1|^2 + |\xi - \xi_1|^2 = \lambda] \right] \\ &< C \frac{M_1}{M_2}. \end{aligned}$$

PROOF BASED ON CHANGE OF VARIABLES

Ideas from (Kenig-Ponce-Vega); see [C-Delort-Kenig-Staffilani].

Recall the Fourier multiplier representation of the propagator:

$$\begin{aligned} e^{it\Delta} f(x) &= c_\pi \int_{\mathbb{R}^2} e^{ix \cdot \xi} e^{-it|\xi|^2} \widehat{f}(\xi) d\xi \\ &= c_\pi \int_{\mathbb{R}^{1+2}} e^{i(x \cdot \xi + t\tau)} \delta_0(\tau + |\xi|^2) \widehat{f}(\xi) d\tau d\xi. \end{aligned}$$

spacetime inverse
Fourier transform

With $f = f_L$ and $g = g_N$, we wish to estimate

$$\|e^{it\Delta} f e^{it\Delta} g\|_{L_{t,x}^2} = \|\mathcal{F}[e^{it\Delta} f e^{it\Delta} g]\|_{L_{\tau,\xi}^2}.$$

Using Fourier transform property, $\mathcal{F}(ab) = \widehat{a} * \widehat{b}$, we find....

FOURIER MANIPULATIONS; DIRAC EVALUATIONS

We wish to estimate (in $L^2_{\tau, \xi}$) the expression

$$\int \delta_0(\tau_1 + |\xi_1|^2) \widehat{f}(\xi_1) \delta_0(\tau_2 + |\xi_2|^2) \widehat{g}(\xi_2).$$
$$\tau = \tau_1 + \tau_2$$
$$\xi = \xi_1 + \xi_2$$

Evaluating the δ functions, we find $\tau_j = -|\xi_j|^2$, so

$$\int \widehat{f}(\xi_1) \widehat{g}(\xi_2)$$
$$\tau = -|\xi_1|^2 - |\xi_2|^2$$
$$\xi = \xi_1 + \xi_2$$

We proceed by **duality**. Let's test this against $d(\tau, \xi)$

DUALITY REDUCES MATTERS TO CERTAIN INTEGRAL

$$\begin{aligned} \|e^{it\Delta} f e^{it\Delta} g\|_{L^2_{t,x}} &= \sup_{\|d\|_{L^2_{\tau,\xi}} \leq 1} \left\langle d(\tau, \xi), \int_{\substack{\tau = -|\xi_1|^2 - |\xi_2|^2 \\ \xi = \xi_1 + \xi_2}} \widehat{f}(\xi_1) \widehat{g}(\xi_2) \right\rangle. \\ &= \sup_d \int d(-|\xi_1|^2 - |\xi_2|^2, \xi_1 + \xi_2) \widehat{f}(\xi_1) \widehat{g}(\xi_2) d\xi_1 d\xi_2. \end{aligned}$$

The preceding Fourier manipulations have reduced matters to **bounding a certain integral**. Our task is to show the integral above is bounded by

$$\lesssim \sqrt{\frac{N}{L}} \|f\|_{L^2} \|g\|_{L^2} \|d\|_{L^2}.$$

SETTING UP THE CHANGE OF VARIABLES

Let's define a change of variables motivated by the arguments of d :

$$u = -|\xi_1|^2 - |\xi_2|^2, \quad v = \xi_1 + \xi_2.$$

- Note that $u \in \mathbb{R}$ and $v \in \mathbb{R}^2$. Thus, $dudv$ is a measure in 3d while $d\xi_1 d\xi_2$ is a measure in 4d.
- Note also that ξ_2 is the argument of $g = g_N$ so it is localized to the smaller dyadic shell $|\xi_2| \sim N \ll L$.
- Let's denote the components of $\xi_j \in \mathbb{R}^2$ with superscripts:

$$\xi_j = (\xi_j^1, \xi_j^2).$$

- The full change of variables is the defined via

$$dudv \, d\xi_2^1 = |J| \, d\xi_1^1 d\xi_1^2 d\xi_2^2 \, d\xi_2^1.$$

We have an **extra** variable **outside** the changed integral.

THE JACOBIAN

The Jacobian matrix J is calculated as

$$J = \begin{bmatrix} \frac{\partial u}{\partial \xi_1^1} & \frac{\partial v^1}{\partial \xi_1^1} & \frac{\partial v^2}{\partial \xi_1^1} \\ \frac{\partial u}{\partial \xi_2^1} & \frac{\partial v^1}{\partial \xi_2^1} & \frac{\partial v^2}{\partial \xi_2^1} \\ \frac{\partial u}{\partial \xi_2^2} & \frac{\partial v^1}{\partial \xi_2^2} & \frac{\partial v^2}{\partial \xi_2^2} \end{bmatrix}.$$

The explicit forms for u, v permit calculating

$$|J| = 2|\xi_1^1 - \xi_1^2|.$$

Since $|\xi_1| \sim L$, we may assume by rotation that $|J| \sim L$.

CHANGING VARIABLES

Our task: Estimate, for $|\xi_1| \sim L$, $|\xi_2| \sim N$, the integral

$$\int_{|\xi_2^2| \lesssim N} \int_{\xi_1, \xi_2^1} d(-|\xi_1|^2 - |\xi_2|^2, \xi_1 + \xi_2) \widehat{f}(\xi_1) \widehat{g}(\xi_2) d\xi_1^1 d\xi_1^2 d\xi_2^1 d\xi_2^2.$$


We insert the Jacobian and reexpress inner integration as

$$\int_{\xi_1, \xi_2^1} d(-|\xi_1|^2 - |\xi_2|^2, \xi_1 + \xi_2) \frac{\widehat{f}(\xi_1) \widehat{g}(\xi_2)}{|J|} |J| d\xi_1^1 d\xi_1^2 d\xi_2^1.$$

Changing variables, we observe this equals

$$\int_{u, v} d(u, v) H(u, v; \xi_2^2) |J| du dv$$

where

$$H(u, v; \xi_2^2) = \frac{\widehat{f}(\xi_1) \widehat{g}(\xi_2)}{|J|}.$$


CAUCHY-SCHWARZ; JACOBIAN REMNANT

We apply Cauchy-Schwarz in u, v to bound by

$$\|d\|_{L^2} \left(\int_{u,v} |H(u, v; \xi_2^2)|^2 dudv \right)^{1/2}.$$

We drop $\|d\|_{L^2} \leq 1$ by duality and change variables back. We get

$$\left(\int_{\xi_1, \xi_2^2} \left| \frac{\widehat{f}(\xi_1) \widehat{g}(\xi_2)}{|J|} \right|^2 |J| d\xi_1^1 d\xi_1^2 d\xi_2^1 \right)^{1/2}.$$

One factor of the Jacobian denominator remains! We gain $L^{-1/2}$.

We still have the extra outside integration....

TRIVIAL CAUCHY-SCHWARZ ON EXTRA INTEGRAL

Recalling what we must control, using what we have obtained....

$$\int_{|\xi_2^2| \lesssim N} \left(\int_{\xi_1, \xi_2^2} \left| \frac{\widehat{f}(\xi_1) \widehat{g}(\xi_2)}{|J|} \right|^2 |J| d\xi_1^1 d\xi_1^2 d\xi_2^1 \right)^{1/2} d\xi_2^2.$$

Apply Cauchy-Schwarz in ξ_2^2 and pay the penalty in the numerator of $N^{1/2}$.

We gain over the trivial bilinear estimate by the factor

$$\sqrt{\frac{(\text{measure of extra support})}{|J|}} = \sqrt{\frac{N}{L}}.$$

4. THE I -METHOD OF ALMOST CONSERVATION

4. THE I -METHOD OF ALMOST CONSERVATION

Let $H^s \ni u_0 \mapsto u$ solve *NLS* for $t \in [0, T_{IWP}]$, $T_{IWP} \sim \|u_0\|_{H^s}^{-2/s}$.

Consider two ingredients (to be defined):

- A **smoothing operator** $I = I_N : H^s \mapsto H^1$. The *NLS* evolution $u_0 \mapsto u$ induces a **smooth reference evolution** $H^1 \ni Iu_0 \mapsto Iu$ solving $I(\text{NLS})$ equation on $[0, T_{IWP}]$.
- A **modified energy** $\tilde{E}[Iu]$ built using the reference evolution.

We postpone how we actually choose these objects.

FIRST VERSION OF THE I -METHOD: $\tilde{E} = H[lu]$

For $s < 1$, $N \gg 1$ define smooth monotone $m : \mathbb{R}_\xi^2 \rightarrow \mathbb{R}^+$ s.t.

$$m(\xi) = \begin{cases} 1 & \text{for } |\xi| < N \\ \left(\frac{|\xi|}{N}\right)^{s-1} & \text{for } |\xi| > 2N. \end{cases}$$

The associated Fourier multiplier operator, $(\widehat{lu})(\xi) = m(\xi)\widehat{u}(\xi)$, satisfies $I : H^s \rightarrow H^1$. Note that, pointwise in time, we have

$$\|u\|_{H^s} \lesssim \|lu\|_{H^1} \lesssim N^{1-s}\|u\|_{H^s}.$$

Set $\tilde{E}[lu(t)] = H[lu(t)]$. Other choices of \tilde{E} are mentioned later.

AC LAW DECAY AND SOBOLEV GWP INDEX

- 1 Modified LWP.** Initial v_0 s.t. $\|\nabla I v_0\|_{L^2} \sim 1$ has $T_{IWP} \sim 1$.
- 2 Goal.** $\forall u_0 \in H^s, \forall T > 0$, construct $u : [0, T] \times \mathbb{R}^2 \rightarrow \mathbb{C}$.
- 3 \iff Dilated Goal.** Construct $u^\lambda : [0, \lambda^2 T] \times \mathbb{R}^2 \rightarrow \mathbb{C}$.
- 4 Rescale Data.** $\|I \nabla u_0^\lambda\|_{L^2} \lesssim N^{1-s} \lambda^{-s} \|u_0\|_{H^s} \sim 1$ provided we choose $\lambda = \lambda(N) \sim N^{\frac{1-s}{s}} \iff N^{1-s} \lambda^{-s} \sim 1$.
- 5 Almost Conservation Law.** $\|I \nabla u(t)\|_{L^2} \lesssim H[Iu(t)]$ and

$$\sup_{t \in [0, T_{IWP}]} H[Iu(t)] \leq H[Iu(0)] + N^{-\alpha}.$$

- 6 Delay of Data Doubling.** Iterate modified LWP N^α steps with $T_{IWP} \sim 1$. We obtain rescaled solution for $t \in [0, N^\alpha]$.

$$\lambda^2(N) T < N^\alpha \iff T < N^{\alpha + \frac{2(s-1)}{s}} \text{ so } s > \frac{2}{2 + \alpha} \text{ suffices.}$$

FIRST VERSION OF THE I -METHOD: $\tilde{E} = H[lu]$

A Fourier analysis established the almost conservation property of $\tilde{E} = H[lu]$ with $\alpha = \frac{3}{2}$ which led to...

THEOREM (CKSTT 02)

$NLS_3^+(\mathbb{R}^2)$ is globally well-posed for data in $H^s(\mathbb{R}^2)$ for $\frac{4}{7} < s < 1$.

Moreover, $\|u(t)\|_{H^s} \lesssim \langle t \rangle^{\beta(s)}$ for appropriate $\beta(s)$.

- The smoothing property $u(t) - e^{it\Delta}u_0 \in H^1$ is **not** obtained.
- Same result for $NLS_3^-(\mathbb{R}^2)$ if $\|u_0\|_{L^2} < \|Q\|_{L^2}$. Here Q is the **ground state** (unique positive solution of $-Q + \Delta Q = -Q^3$).
- Fourier analysis leading to $\alpha = \frac{3}{2}$ in fact gives $\alpha = 2$ for most frequency interactions.

ALMOST CONSERVATION LAW FOR $H[lu]$

PROPOSITION

Given $s > \frac{4}{7}$, $N \gg 1$, and initial data $\phi_0 \in C_0^\infty(\mathbb{R}^2)$ with $E(I_N u_0) \leq 1$, then there exists a $T_{lwp} \sim 1$ so that the solution

$$u(t, x) \in C([0, T_{lwp}], H^s(\mathbb{R}^2))$$

of $NLS_3^+(\mathbb{R}^2)$ satisfies

$$E(I_N u)(t) = E(I_N u)(0) + O(N^{-\frac{3}{2}+}),$$

for all $t \in [0, T_{lwp}]$.

IDEAS IN THE PROOF OF ALMOST CONSERVATION

- Standard Energy Conservation Calculation:

$$\begin{aligned}\partial_t H(u) &= \Re \int_{\mathbb{R}^2} \overline{u}_t (|u|^2 u - \Delta u) dx \\ &= \Re \int_{\mathbb{R}^2} \overline{u}_t (|u|^2 u - \Delta u - iu_t) dx = 0.\end{aligned}$$

cancellation

- For the smoothed reference evolution, we imitate....

$$\begin{aligned}\partial_t H(lu) &= \Re \int_{\mathbb{R}^2} \overline{lu}_t (|lu|^2 lu - \Delta lu - i|lu_t) dx \\ &= \Re \int_{\mathbb{R}^2} \overline{lu}_t (|lu|^2 lu - I(|u|^2 u)) dx \neq 0.\end{aligned}$$

commutator!

- The increment in modified energy involves a commutator,

$$H(lu)(t) - H(lu)(0) = \Re \int_0^t \int_{\mathbb{R}^2} \overline{lu}_t (|lu|^2 lu - I(|u|^2 u)) dx dt.$$

- Littlewood-Paley, Case-by-Case, (Bi)linear Strichartz, $X_{s,b}$

REMARKS

- The almost conservation property

$$\sup_{t \in [0, T_{IWP}]} \tilde{E}[Iu(t)] \leq \tilde{E}[Iu_0] + N^{-\alpha}$$

leads to GWP for

$$s > s_\alpha = \frac{2}{2 + \alpha}.$$

- The I -method is a *subcritical method*. To prove the Scattering Conjecture at $s = 0$ via the I -method would require $\alpha = +\infty$.
- The I -method *localizes the conserved density in frequency*. Similar ideas appear in recent critical scattering results.
- There is a *multilinear corrections algorithm* for defining other choices of \tilde{E} which yield a better AC property.