



High Salt Intake Linked to Social Inequalities

Published: January 8, 2013. by [University of Warwick](#)

People from low socio-economic positions in Britain eat more salt than the well off, irrespective of where they live, states a paper led by Warwick Medical School published on Tuesday in the *BMJ Open* journal .

The research was carried out by the World Health Organization Collaborating Centre for Nutrition , based in the Division of Mental Health & Wellbeing of Warwick Medical School at the University of Warwick.

The study looked at the geographical distribution of habitual dietary salt intake in Britain and its association with manual occupations and educational attainments, both indicators of socio-economic position and key determinants of health.

The researchers used the British National Diet and Nutrition Survey (2000-1), a national representative sample of 2,105 men and women aged 19-64 years living in Britain. Salt intake was assessed with two independent methods: a 7-day dietary record and the 'gold standard' 24h urine collections for sodium determination (direct marker of salt intake).

The study provides evidence for the first time that salt intake is significantly higher in those with low educational attainment and in manual occupations, when the effects of geographical variations are stripped out (people living in Scotland had higher salt intake than those in England & Wales).

Professor Francesco Cappuccio, senior author and Director of the WHO Collaborating Centre, said: "These results are important as they explain in part why people of low socio-economic background are more likely to develop high blood pressure (hypertension) and to suffer disproportionately from strokes, heart attacks and renal failure."

Ms Teresa Morris of The Bupa Foundation, which funded the study, said: "Habitual salt intake in most adult populations around the world exceeds 10 g per day and the World Health Organization recommends that daily intake should not exceed 5 g .

"Population salt reduction programmes are a cost-effective way of reducing the burden of cardiovascular disease nationally and globally."

Professor Cappuccio continued: "We have seen a reduction in salt intake in Britain from 9.5 to 8.1 g per day in the period 2004-2011, thanks to an effective policy which included awareness campaigns, food reformulation and monitoring.

"Whilst this is an achievement to celebrate, our results suggest the presence of social inequalities in levels of salt intake that would underestimate the health risks in people who are worse off - and these are the people who need prevention most.

Related Content

- [External link to University of Warwick](#)
- [More news from University of Warwick](#)

Related Articles »

Salt 7/6/11 **Cutting Down on Salt Doesn't Reduce Your Chance of Dying**

Wiley-Blackwell
Moderate reductions in the amount of salt people eat doesn't reduce their likelihood of dying or experiencing cardiovascular disease. This is the main conclusion from a systematic review published in the latest edition of The Cochrane Library. ...

Salt 11/1/10 **Mandatory Curbs on Food Salt Content 20 Times More Effective Than Voluntary Curbs**

BMJ-British Medical Journal
Imposing statutory limits on the salt content of processed foods could be 20 times more effective than voluntary curbs by industry, finds research published online in the journal Heart. The Australian researchers assessed the public health benefits and ...

Salt 12/21/11 **Salt Policy Makers Eat Too Much Salt at Work**

BMJ-British Medical Journal
Salt policy makers in the Netherlands are consuming more than the average daily recommended salt intake of 6 grams in one hot meal at their work canteens, finds a study in the Christmas issue published on bmj.com today. ...

Salt 6/16/10 **Some People Do Not Taste Salt Like Others**

Penn State
Low-salt foods may be harder for some people to like than others, according to a study by a Penn State College of Agricultural Sciences food scientist. The research indicates that genetics influence some of the difference in the levels ...

Recommendation 12/15/10 **Sticking to Dietary Recommendations Would Save 33,000 Lives a Year in the UK**

BMJ-British Medical Journal
If everyone in the UK ate their "five a day," and cut their dietary salt and unhealthy fat intake to recommended levels, 33,000 deaths could be prevented or delayed every year, reveals research published online in the Journal of ...

[More »](#)

"The diet of disadvantaged socio-economic groups tends to be made up of low-quality, salt-dense, high-fat, high-calorie unhealthy cheap foods.

"Behavioural approaches to healthy eating are unlikely to bring about the changes necessary to halt the cardiovascular epidemic and would also widen inequalities.

"Since the majority of dietary salt is added during commercial food production, widespread and continued food reformulation is necessary through both voluntary as well as regulatory means to make sure that salt reduction is achieved across all socio-economic groups", Professor Cappuccio concluded.

Show Footnotes »

Read Less (Back to summary page) »

Translate this page: [Chinese](#) [French](#) [German](#) [Italian](#) [Japanese](#) [Korean](#) [Portuguese](#) [Russian](#) [Spanish](#)

Disclaimer: The views expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the ScienceNewsline.com.

Most Popular - Medicine »

GME »

Most Physicians Do Not Meet Medicare Quality Reporting Requirements

Washington, DC – A new Harvey L. Neiman Health Policy Institute study shows that fewer than one-in-five healthcare providers meet Medicare Physician Quality Reporting System (PQRS) requirements. Those that ...

BLOOD »



Protein Essential for Healthy Eyes Described by Hebrew University, US Researchers

DEIMINATION »

Identifying the Molecular Causes of Vision Loss in Demyelinating Disease

Demyelinating diseases, such as multiple sclerosis (MS), are frequently associated with the progressive loss of vision. The retinal nerve damage is thought to be caused by immune system-mediated inflammation; ...

PARKINSON »

Study: Parkinson's Disease Itself Does Not Increase Risk of Gambling, Shopping Addiction

MINNEAPOLIS – Parkinson's disease itself does not increase the risk of impulse control problems such as compulsive gambling and shopping that have been seen in people taking certain drugs ...

NEURAL »



Transplanted Neural Stem Cells Treat ALS in Mouse Model

[ScienceNewsline.com](#) | [About](#) | [Privacy Policy](#) | [Feedback](#) | [Mobile](#)

The selection and placement of stories are determined automatically by a computer program. All contents are copyright of their owners except U.S. Government works. U.S. Government works are assumed to be in the public domain unless otherwise noted. Everything else copyright ScienceNewsline.com.