






















TESS Duotransit Candidates from the Southern Ecliptic Hemisphere

Faith Hawthorn^{*1,2} , Sam Gill^{†1,2} , Daniel Bayliss^{‡1,2} , Hugh P. Osborn³ , Ingrid Pelisoli¹ , Toby Rodel^{1,2,4} , Kaylen Smith Darnbrook¹, Peter J. Wheatley^{1,2} , David R. Anderson^{1,2} , Ioannis Apergis^{1,2} , Matthew P. Battley⁵ , Matthew R. Burleigh⁶ , Sarah L. Casewell⁶ , Philipp Eigmüller⁷ , Maximilian N. Günther⁸ , James S. Jenkins^{9,10} , Monika Lendl⁵ , Maximiliano Moyano¹¹ , Ares Osborn^{1,2} , Gavin Ramsay¹² , Solène Ulmer-Moll³ , Jose I. Vines¹¹ , Richard West^{1,2}

The authors' affiliations are shown in Appendix B.

E-mails: ^{*}faith.hawthorn@warwick.ac.uk, [†]samuel.gill@warwick.ac.uk, [‡]d.bayliss@warwick.ac.uk

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

Discovering transiting exoplanets with long orbital periods allows us to study warm and cool planetary systems with temperatures similar to the planets in our own Solar system. The *TESS* mission has photometrically surveyed the entire Southern Ecliptic Hemisphere in Cycle 1 (August 2018 - July 2019), Cycle 3 (July 2020 - June 2021) and Cycle 5 (September 2022 - September 2023). We use the observations from Cycle 1 and Cycle 3 to search for exoplanet systems that show a single transit event in each year - which we call *duotransits*. The periods of these planet candidates are typically in excess of 20 days, with the lower limit determined by the duration of individual *TESS* observations. We find 85 duotransit candidates, which span a range of host star brightnesses between $8 < T_{\text{mag}} < 14$, transit depths between 0.1 per cent and 1.8 per cent, and transit durations between 2 and 10 hours with the upper limit determined by our normalisation function. Of these candidates, 25 are already known, and 60 are new. We present these candidates along with the status of photometric and spectroscopic follow-up.

Key words: techniques: photometric — planets and satellites: detection — planets and satellites: general

1 INTRODUCTION

Longer period transiting exoplanets allow us to measure the densities of warm and cool planets, assisting with studies of atmospheric composition from transit spectroscopy (e.g. Albrecht et al. 2012; Wang et al. 2021). Planetary obliquity is an indicator of planetary migration mechanisms (e.g. Albrecht et al. 2012), and long period planets are amenable to studies of the Rossiter-McLaughlin effect (e.g. Ulmer-Moll et al. 2023) to enable determination of the spin-orbit alignment of the system. When compared to short period hot Jupiter planets, the few well-studied long period planets discovered to date appear to be more aligned (Rice et al. 2022), indicating possible migration mechanisms operating within the protoplanetary disk (e.g. Madhusadhan et al. 2017). Since the launch of *TESS* (Transiting Exoplanet Survey Satellite; Ricker et al. 2015), a total of 392¹ exoplanets have been confirmed using *TESS* data (NASA Exoplanet Archive; Akeson et al. 2013). Of these systems, only 56 have orbital periods longer than the typical *TESS* Sector time span of 27 days.

The *Kepler* mission (Borucki et al. 2010) monitored a single 105 square degree field for approximately four years, and was therefore able to discover long period transiting exoplanets. However, most of the host stars are fainter than the capabilities of modern spectrographs to measure planetary masses and the planets cannot be confirmed and fully characterised. Hsu et al. (2019) placed a limit on the occurrence rate of $\lesssim 0.247$ planets per FGK star on orbital periods from 237-500 days. The *Kepler* exoplanet sample includes

42 planets with determined periods longer than 20 days with a robust mass determination from radial velocity confirmation.

The *TESS* spacecraft has completed surveys of the Southern Ecliptic Hemisphere on two occasions – in Cycle 1 (July 2018 to July 2019) and in Cycle 3 (July 2020 to July 2021). *TESS* has also re-observed the Southern Ecliptic Hemisphere in Cycle 5 (September 2022 to September 2023), but many of these observations are unavailable at the time of writing and thus are not included in this work. During these surveys a number of stars have shown a single transit (a *monotransit*), whereby one transit like feature is found in the lightcurve but is not seen to repeat. With only one transit we are not able to determine the orbital period of the planet, although with prior knowledge of the star we can calculate the probability distribution for the orbital period, e.g. Osborn et al. (2022a). A small number of these *TESS* monotransit candidates have been confirmed, e.g. Gill et al. (2020a,c) and Lendl et al. (2020a).

In addition to the *TESS* monotransit candidates, there exist candidates that transit exactly once in Cycle 1 and once in Cycle 3. These candidates are *biennial duotransits*, as opposed to candidates that may have two transits in a single *TESS* cycle. For the remainder of this paper we use the term *duotransit* to refer only to such biennial duotransits. These duotransits confer two main advantages over monotransit candidates. Firstly, the true period of a duotransit candidate is limited to a discrete set of possible periods, P_n , given by

$$P_n \in \left(\frac{\Delta T}{n} \right), n = 1, \dots, n_{\text{max}}, \quad (1)$$

where ΔT is the time difference between the Cycle 1 and the Cycle 3

¹ <https://exoplanetarchive.ipac.caltech.edu> (2023 Oct 09)

transit event and n is a whole number ranging from $n=1$ for the longest possible period to n_{max} for the shortest possible period. Since the duration of a TESS sector is ≈ 27 days, we find $P_{n_{max}}$ is typically in the region of 20 days unless there is more than one Sector of data in the TESS Cycle 1 or Cycle 3 observations. The limited set of possible periods means it is possible to check specific orbital periods via photometric monitoring of the duotransit candidate at specific times. This is much more efficient than the continuous monitoring required to find the periods of monotransit candidates.

Secondly, duotransits are more robust against false positive signals. We can check that the two transit signals for a duotransit candidate match in depth, total duration, ingress and egress duration, limb darkening parameters, and impact parameter. By contrast, we are far less certain that a single monotransit event is not caused by some systematic or non-planetary astrophysical event.

Cooke et al. (2018) predicted that hundreds of long-period planets would be detectable as monotransit events in the TESS sample, with similar results found by Villanueva et al. (2019). Furthermore, Cooke et al. (2021) found that many of these will have a second detectable transit in Cycle 3 of TESS data, making the true orbital period one of an average of 38 period aliases.

Both mono- and duotransits require extensive follow-up with both photometric and radial velocity observations in order to confirm the transit signal, orbital period, radius and mass. Constraining the period of a candidate can greatly reduce the number of radial velocity measurements needed to confirm the exoplanet and measure its mass (e.g. Gill et al. 2020c).

In this paper we set out our search for duotransits from the Cycle 1 and Cycle 3 TESS FFI (Full-Frame Image) lightcurves. In Section 2 we describe the TESS data that we use to search for duotransit candidates. In Section 3 we outline our search algorithm and vetting procedure. In Section 4 we set out the results of our search, including the details of the 85 duotransit candidates. Finally in Section 5 we discuss the prospects for confirming these duotransit candidates via follow-up photometric and spectroscopic programmes.

2 TESS OBSERVATIONS

TESS is a space-based NASA mission focusing on the discovery of transiting exoplanets around bright stars (Ricker et al. 2015). TESS has four cameras, each equipped with a grid of four CCDs for a total combined field-of-view of TESS of $24^\circ \times 96^\circ$. In Cycle 1 (2018 Jul 25 - 2019 Jul 17), TESS observed almost the entire Southern Ecliptic Hemisphere in 13 Sectors, each observed for a period of approximately 27 days. Each TESS sector has a gap of approximately 2.4 days, during which time the satellite downlinks data. There is a similar gap in between Sectors. Other gaps in the data can be due to technical problems with the spacecraft or scattered light (mostly from the Earth) making some image frames unusable (e.g. Dalba et al. 2020). Select stars were observed with 2 minute cadence, while the full frame images (FFIs) were observed with a cadence of 30 minutes.

TESS reobserved the Southern Ecliptic Hemisphere in Cycle 3 (2020 Jul 5 - 2021 Jun 24), with the FFIs collected with a shorter cadence of 10 minutes. During Cycle 3, TESS slightly shifted its survey fields in order to cover spatial gaps in between the Cycle 1 Sectors.

For our search for duotransit candidates, we use the TESS full-frame image (FFI) light curves produced by the Science Processing Operations Center pipeline (SPOC; Jenkins et al. 2016), which are publicly available from the Mikulski Archive for Space Telescopes

(MAST)². The SPOC pipeline produces FFI light curves for approximately 160,000 stars per sector, based on selection criteria set out in Caldwell et al. (2020).

We focus solely on the Southern Ecliptic Hemisphere, as at the time of writing TESS had completed its campaign of the Southern Ecliptic in two cycles. We use Sector 1-13 data from Cycle 1, and Sector 27-38 data from Cycle 3. In total, this results in a sample of 1,422,473 stars. We search light curves from the Presearch Data Conditioning Simple Aperture Photometry (PDCSAP; Stumpe et al. 2012, 2014; Smith et al. 2012) lightcurves, which have instrumental systematic trends removed from the Simple Aperture Photometry (SAP) flux, but should retain stellar variability including any transit events.

In addition to using flux measurements from the FFI PDCSAP lightcurves, we also make use of the sky background and momentum centroiding measurements provided in the PDCSAP lightcurve files via the keywords SAPBK and MOMCENTR1/MOMCENTR2 respectively. These data allow us to rule out a large number of false candidates as described in Section 3.2.

3 DUOTRANSIT SEARCH

Our search for TESS duotransit candidates arises out of a more general search for TESS monotransits. We therefore begin by searching the entire set of 1,422,473 SPOC PDCSAP FFI light-curves for monotransit events, and then we match up events to detect duotransit candidates.

3.1 Searching for Monotransit Events

We begin by downloading the SAP and PDCSAP SPOC FFI light-curve products from MAST. To remove bad data points from the light-curves we discarded data points with QUALITY bit values above 0. We read the meta information for each light-curve, including the TIC ID, effective temperature, stellar radius, and metallicity if available. We assume a solar metallicity of 0 dex for stars with no metallicity value available. We split each lightcurve into segments defined by gaps in excess of 2.4 hours and normalise each separately. This accounts for gaps in individual orbits due to spacecraft data download times and regions of data flagged by the SPOC algorithm. For each segment, we smooth the lightcurve using an iterative Savitzky–Golay filter (5 iterations using a threshold of $3\text{-}\sigma$) with a window size 48 hours (e.g. Hattori et al. 2022). This window size strikes a balance of removing stellar activity but ultimately sets an upper limit on the width of transit events our algorithm can detect. Our tests show that transit widths below ~ 2.3 days are easily detectable using this filter width. For events wider than this, we find that the Savitzky–Golay filter begins to confuse transit signals with stellar activity and thus the event is not recovered. In addition, our tests showed that the change in cadence between Cycle 1 (30 minutes) and Cycle 3 (10 minutes) had a negligible effect on the filter. For a warm Jupiter in a circular orbit around a Solar type star, we estimate an upper limit to the possible recovered orbital periods of in excess of 1000 days.

Once the lightcurves are normalised and flattened, we pass them to `monofind`, our custom monotransit detection algorithm. `monofind` searches for monotransit events in the lightcurve where three consecutive 30-minute data points are 3 median absolute deviations below the normalised median of the lightcurve. An example of this method

² Accessible at <https://mast.stsci.edu/>




Figure 1. Upper panel: The *TESS* Sector 32 lightcurve of TOI-2447 (black) with detrending model (red solid line) and the 3-MAD detection threshold (red dashed line). A single event was detected highlighted in blue. **Lower panel:** An inspection of the transit event with best fitting transit model (red solid line).

is shown in Figure 1, as applied to TOI-2447 (Gill et al., submitted). For the Cycle 1 light-curves we use the native 30 minute data points, while for the Cycle 3 lightcurves we bin the native 10-minute cadence by three to a 30-minute cadence to achieve a consistent detection threshold for transit events.

3.2 Selecting Duotransit Candidates

After searching all the *TESS* Cycle 1 and Cycle 3 SPOC FFI lightcurves for monotransit events, we cross match detections to search for stars which show exactly one monotransit event in Cycle 1 and one monotransit event in Cycle 3. This yielded a list of 9718 duotransit candidates.

We then performed a visual inspection of these 9718 lightcurves to check which of these showed two events that could possibly be transit events. This allowed us to remove obvious false positives that had triggered the `mono.find` algorithm but did not have two transit-shaped events in the lightcurve. In most instances these were some kind of variable star. This quick visual inspection took us down to a total of 736 reasonable duotransit candidates. The majority of candidates




Figure 2. Three example duotransit candidates ruled out during the visual inspection of the transit events in the *TESS* light curves, with an arbitrary flux offset for clarity. The dotted red line indicates the 3 median absolute deviations detection threshold used in `mono.find`. The red triangles indicate the event that triggered the detection by `mono.find`. From top to bottom, these example events are caused by an asteroid passing through the target and background aperture, systematic noise caused by the spacecraft, and changes in amplitude of a variable stars.

that did not pass this vetting step were caused by asteroids passing through the target and/or background pixel apertures, systematics noise events due to the spacecraft, or changes in the amplitude of variable stars (see examples presented in Figure 2).

We examined each of these duotransit candidates in order to ascertain which appeared to be good transiting exoplanet candidates, and which were false-positives. Examples of false-positive scenarios included those causing transit-like events such as eclipsing binary stars, *TESS* spacecraft systematics (in particular momentum dumps), stellar variability, and Solar System asteroids passing through the background and target photometric aperture.

We are able to identify events caused by asteroids crossing the photometric aperture by checking for variations in the sky background (SAPBKG) that correlate with the monotransit event. We also noticed that such events were typically very asymmetrical in shape, and therefore unlike a bona-fide monotransit event. In total, we identified 170 of our initial duotransit candidates as being due to asteroids crossing the photometric aperture in Cycle 1 or Cycle 3.

We identified monotransit events that were likely caused by eclips-

ing binaries by examining the results from the transit/eclipse fit from `monofind`. We designated any event with a modeled companion radius $> 2 R_J$ ($\sim 22 R_{\oplus}$) to be an eclipsing binary, and removed it from our duotransit candidate list. Such well-separated planets are not expected to be inflated like hot Jupiters (e.g. see figure 2 from [Lopez & Fortney 2016](#)) and thus this limit is justified. We also designated a candidate as an eclipsing binary if there was evidence of a secondary eclipse in the lightcurve. Furthermore, we excluded events which had significant depth differences between Cycles 1 and 3. In total we identified 384 eclipsing binaries from our initial duotransit candidate list.

We searched for blended eclipsing binaries by inspecting the lightcurves of nearby stars, by checking for centroiding offsets during the monotransit events, and by examining the Target Pixel File (TPF) using our custom `spectpf` tool. `spectpf` allows us to make a lightcurve for any pixel or set of pixel in the TPF, and helps determine if the monotransit event is on the star of interest, or is on a nearby star. In total we identified 58 blended eclipsing binaries from our initial duotransit candidate list.

We also examined the pre-normalised and pre-flattened PDCSAP lightcurves, as well as the SAP lightcurves. This was to ensure that neither the PDCSAP algorithm nor our own normalisation and flattening had significantly altered the shape of the detected monotransit events. We also inspected by-eye the full unflagged SPOC lightcurve for additional transit events that may have been excluded from the PDCSAP data.

Finally we checked if any of the duotransit candidates had an associated Data Validation Report (DVR, [Twicken et al. 2019](#)). DVRs are created for potential *TESS* planets candidates processed with the SPOC pipeline and contain initial findings about the system including a detailed model fit and analysis of nearby stars to exclude blend scenarios. If a DVR determined a candidate to be a false-positive, we investigated the target further to assess if it warranted exclusion from our list.

Some candidates displayed depth differences between the Cycle 1 and Cycle 3 events. SPOC lightcurves use unique masks for each individual Sector which may lead to different amounts of dilution from neighboring stars depending on telescope pointing and orientation for each sector ([Bryant et al. 2023](#)). This should be accounted for in the detrended SPOC data (PDCSAP) but might not be perfect, and thus small sector-dependent depth differences are expected ($\lesssim 2$ ppt), particularly in crowded fields. Some candidates had larger depth and width differences, which could be multi-planet systems or blended eclipsing binaries. However, since they do not meet our criteria for selection as duotransit candidates, they were excluded from our list. We also found 39 duotransit candidates that we associate with spacecraft systematics (stray light, momentum dumps etc.³). This number also includes a few events resulting from stellar variability that was poorly matched by our detrending model.

A flowchart illustrating the candidate search and vetting process, and the associated numbers of candidates categorised at each stage, is shown in Figure 3.

In summary, out of the 1,422,473 unique TIC IDs, 9718 duotransit candidates were identified by the `monofind` algorithm, and after visual inspection this was reduced to 736 candidates pre-vetting. As a result of the vetting process, 58 objects were classified as blended eclipsing binaries ('blends'), 170 as Solar System asteroids, 384 as eclipsing binaries, and 39 as events caused by either spacecraft




Figure 3. Flowchart of the `monofind` algorithm candidates and the vetting process that followed, with the number of candidates categorised at each stage as blended eclipsing binaries ('blends'), asteroids, eclipsing binaries, and other systematics and stellar variability features.

systematics or poorly-detrended stellar variability. We list a sample of these events along with their designation in Table A4. The vetting process left us with a final number of 85 duotransit candidates that we present as likely planetary in nature in this paper.

3.3 Modelling Candidate Duotransit Events

In order to derive planetary parameters for the candidates, as well as the most likely period aliases, we must fit the available transits. However, most transit model fitting requires a priori knowledge of the orbital periods, and is certainly not optimised to sample the extremely narrow but widely-separated period regions found for duotransits. We used the `MonoTools` package ([Osborn 2022](#)) which fits transits in a way agnostic of orbital period and then computes the posterior density function of the planet candidate from the orbital velocity implied from the transit impact parameter, radius ratio and duration. This has previously been used to model other duotransiting planet candidates that were subsequently confirmed (e.g. [Osborn et al. 2021, 2022a](#)).

`MonoTools` fits transits using the `exoplanet` python library ([Foreman-Mackey et al. 2021](#)). Stellar parameters from the *TESS* Input Catalogue (TICv8; [Stassun et al. 2019](#)) were used as priors for each of the fits; as was the [Kipping \(2013\)](#) eccentricity prior, the [Espinoza & Jordán \(2016\)](#) impact parameter prior, and quadratic limb-darkening parameters constrained using theoretical predictions for the *TESS* bandpass of [Claret \(2017\)](#). In the majority of cases, a cubic spline was fit to the transit-masked *TESS* PDCSAP flux in order to pre-flatten the lightcurve and a window of 5.5 transit durations was cut around the transit. However, after modelling some of these fits appeared to poorly model the out-of-transit variability. In

³ *TESS* Instrument Handbook; https://archive.stsci.edu/tess/tess_drn.html

these cases we turned to a simple harmonic oscillator (SHO) term celerite Gaussian process (Foreman-Mackey et al. 2017), which we constrained by first sampling out-of-transit data and then using the inferred hyperparameter distributions as priors for a combined model fit. A log scatter parameter was used to encapsulate additional noise for each transit. The transit model was then sampled using the Hamiltonian Monte Carlo of Salvatier et al. (2016) using a burn-in phase of 800 steps and 1500 samples on each of four independent chains, resulting in typical effective sample sizes of 2500-3000 for each parameter for each candidate. The derived parameters are shown in full online but summarised in Table A3. The probabilistic planetary period distributions are shown in Table A3 and in Figure 4 ordered from shortest to longest.

4 RESULTS

4.1 85 Duotransit Planet Candidates

Following the methods described in Section 3, we have found a total of 85 duotransit planet candidates from our `monofind` search and vetting of the SPOC FFI lightcurves for the Southern Ecliptic *TESS* fields (Sectors 1-13, 27-34). The stellar properties of our candidates are set out in Table A1, where N_{sec} denotes the total number of *TESS* sectors in which the target was observed. The transit event parameters are set out in Tables A2 & A3, including the central transit times T_{c1} and T_{c2} of the Cycle 1 and Cycle 3 events respectively, and the separation between the two events in days ΔT_c . We also include the *Gaia* flags of NSS (Non-Single Star), where 0 denotes a good single-star solution, 1 denotes an astrometric binary, 2 denotes a spectroscopic binary and 3 denotes an eclipsing binary; and the *Gaia* robust radial velocity (RV) amplitude in km s^{-1} , calculated using the standard deviations of individual RV measurements as set out in *Gaia* Data Release 3 (DR3, Gaia Collaboration 2022). We do not remove candidates from our target list based on these values, however, since an NSS designation is often consistent with the presence of a planet and it is not clear how robust the *Gaia* DR3 RV amplitudes are (e.g. Seabroke et al. 2021). In Table A3, we also show the most probable value of P_n (P_{marg}) and the corresponding value of n (N_{alias}).

The transit events for all 85 duotransit candidates are plotted in Figures A1-A5. We plot the best fit transit and eclipse model for each event from the `monofind` algorithm (see Section 3.1). We also plot the sky distribution of the candidates in Figure 5.

Figure 6 shows where our duotransit candidate host stars lie on the HR diagram, with most on the main-sequence spanning spectral types A-K, some on the main-sequence turn off, and one in the hot sub-dwarf regime (see Section 4.2). Our candidates span a range of brightness between $8 < T_{\text{mag}} < 14$, with most candidates lying between $10 < T_{\text{mag}} < 13$; see Figure 11. This magnitude range is largely governed by the magnitude distribution of the SPOC FFI lightcurves that cuts off at around $T_{\text{mag}} = 13.5$ due to the number of lightcurves per sector being restricted to approximately 160,000 (Jenkins et al. 2016).

We calculated the total SNRs (Signal-to-Noise Ratios) for each of our candidates as a function of the SNRs of each individual transit event. The SNR of each transit event was estimated using the double box approximation model described by Kipping (2023). Using this square well transit model, the SNR for each transit event with a depth δ in ppm, a duration T_{dur} in hours, light curve noise σ in ppm, and a contamination ratio C was calculated using Equation 2:

$$\text{SNR} = \frac{1}{C} \times \frac{\delta}{\sigma} \sqrt{T_{\text{dur}}} \quad (2)$$

The contamination ratio and noise value were both extracted directly from the headers of the SPOC lightcurves. In some cases these values were missing from the headers of the Cycle 3 lightcurves, in such instances we used the same noise as the Cycle 1 lightcurve. To combine the SNR across both transit events we add the individual SNR for each event in quadrature.

We plot the distribution of the total SNR values for our 85 candidates in Figure 7, and find that most candidates have a SNR between 10-70, with the distribution peaking at SNR values of 25-30. There is one candidate with a total SNR value below 10 and four candidates with a SNR value above 100.

4.2 Candidates of Special Interest

Based on our *Gaia* H-R diagram in Figure 6, we identify three candidates that are distinct from the rest.

4.2.1 AIV type host star, TIC-221915858

TIC-221915858 (candidate 45) is the most luminous star in our sample and resides in the region of the colour-magnitude diagram where the main-sequence approaches the sub-giant branch (Figure 6). This is complemented by *Gaia* DR3 stellar parameters suggesting that TIC-221915858 is a hot ($T_{\text{eff}} = 9000 \text{ K}$) A1 star with a RUWE of 0.814 indicative of a good astrometric solution. The fitted spectral energy distribution in TIC V8, which uses *Gaia* DR2 along with 2MASS colours, suggests TIC-221915858 is a main sequence A1 ($T_{\text{eff}} = 9475 \pm 187 \text{ K}$) star with mass and radius of $2.41 \pm 0.33 M_{\odot}$ and $2.47 \pm 0.09 R_{\odot}$ respectively. In addition to being the most luminous host star in our sample, TIC-221915858 is also the hottest on the main sequence and represents a under-sampled population of exoplanets around hot stars. If a planet is confirmed around TIC-221915858, it would be the hottest planet host star discovered by the *TESS* mission⁴. The 5.74 ppt transit depth is detectable with ground-based photometric facilities and so it is possible to recover the orbital period with ground-based instruments. However, recovering the spectroscopic orbit will be challenging, in-part due to the typical rotation and lack of absorption lines in A-type stars making it difficult to measure a precise CCF (Cross-Correlation Function) centre. Doppler tomography (e.g. Watson et al. 2019) has been used successfully to determine the radial-velocity semi-amplitude of fast-rotating stars with broad CCFs (Temple et al. 2018, 2019) and could be applicable to TIC-221915858 once the orbital period is known. We provide the light curves and transit models for this candidate in Figure A2.

4.2.2 Late K-dwarf host, TOI-4310/TIC-303317324

While the majority of our duotransit candidates orbit F, G and early K type dwarf stars, the candidate TOI-4310 (TIC-303317324, candidate 65) is a late K-dwarf star with $T_{\text{eff}} = 4159 \text{ K}$. This is an outlier amongst our candidates on the *Gaia* colour-magnitude diagram (Fig 6), positioned much further down the main sequence than any of our other candidates. The estimated radius of TOI-4310 is $0.72 R_{\odot}$, and the transit model indicates the transiting planet would have a radius of $2.67 R_{\oplus}$, making it one of the smallest of our duotransit planet candidates. We provide the light curves and transit models for this candidate in Appendix Figure A1.

⁴ exoplanetarchive.ipac.caltech.edu (as of 2023 Oct 10)




Figure 4. Probability distribution of possible periods for each duotransit candidate as computed by MonoTools, normalised to the most likely period alias. The vertical colourbars refer to six steps in the log probability of each alias, and the green triangles represent the minimum eccentricity aliases.




Figure 5. The ecliptic plane (solid red line) with the positions of the 85 duotransit candidates marked with red-inset black points.

4.2.3 Hot subdwarf host, TOI-709/TIC-396720998

TOI-709 (TIC-396720998, candidate 78) has two ~ 6 ppt deep transits with durations of ~ 4.3 hours. It is the only candidate in our list categorised as a hot subdwarf host star (LB 1721; Culpan et al. 2022), residing on the $G_{BP} - G_{RP} < 0$ region of the colour-magnitude diagram in Figure 6 outside of the main population of exoplanet hosts from our candidates. Hot subdwarfs are evolved compact stars, mainly resulting from enhanced mass-loss at the tip of the red giant branch, which likely occurred due to binary interaction (Han et al. 2002, 2003; Maxted et al. 2001; Pelisoli et al. 2020). This makes them targets of interest for probing the survivability of planets to both stellar evolution and binary environments. Most hot subdwarf companions fall into two populations containing either close white dwarf or M-dwarf/brown dwarf companions or wide FGK-type companions (e.g. Schaffenroth et al. 2022), with no planetary companions confirmed to date (e.g. Van Grootel et al. 2021; Thuillier et al. 2022).

Parameters from TICv8 poorly constrain the mass and a radius of the host to $0.5 \pm 0.3 M_{\odot}$ and $0.15 \pm 0.11 R_{\odot}$ ($\log g = 5.8 \pm 0.9$), which are consistent with a hot subdwarf and make it the smallest candidate host star in our list. Jeffery et al. (2021) obtained for this star a T_{eff} of $45,600 \pm 1000$ K and a consistent $\log g$ of 6.08 ± 0.04 and classified it as helium-rich sdO (He-sdO). They reported no radius or mass estimate. To derive a radius consistent with the precise T_{eff} and $\log g$ of Jeffery et al. (2021), we performed a spectral energy distribution (SED) fit using `speedyfit`⁵. We employed spectral models from the Tübingen NLTE Model-Atmosphere Package (TMAP, Werner & Dreizler 1999; Werner et al. 2003; Rauch & Deetjen 2003). Initial attempts revealed the existence of excess flux towards the red/near-infrared compared to the hot subdwarf model. We therefore included a second component in the fit, modelled with ATLAS9 spectra (Castelli & Kurucz 2003). The $\log g$ was kept fixed at the spectroscopic value for the hot subdwarf (as it is poorly constrained by a SED fit), and a Gaussian prior was applied on the temperature. The temperature of the companion star was allowed to vary freely within 3500 and 6000 K (restricted by the lack of contribution to the spectrum in Jeffery et al. 2021, : higher temperatures would lead to visible lines in the spectrum; for lower temperatures, the hot subdwarf would completely dominate and a companion would have no effect) and the $\log g$ was left to vary in the range of 4 to 5. A prior on the *Gaia* parallax was also applied, which enables the determination of precise radii. The MCMC fit converged to radii of $0.18^{+0.06}_{-0.03} R_{\odot}$ and $0.88^{+0.30}_{-0.15} R_{\odot}$ for the hot subdwarf and the companion, respectively.

⁵ <https://speedyfit.readthedocs.io/>

The companion T_{eff} was found to be 5100 ± 120 K, consistent with a K-type main sequence star. FGK-type companions are found with 30 per cent of hot subdwarfs (Stark & Wade 2003), and their periods are in the range of 100-1000 days (e.g. Vos et al. 2019), which could be consistent with the orbital period of the duotransit signal. We note that there is also another much deeper (5.8%) transit/eclipse signal with a 32 day period that is reported in TFOP, and was the reason this candidate was flagged as a TOI. However that signal is distinct from the duotransit signal that we report in this paper. We provide the light curves and transit models for this candidate in Appendix Figure A4.

4.3 Other notable candidates

4.3.1 Continuous Viewing Zone candidates

Three of our candidates lie inside or close to the *TESS* Continuous Viewing Zone (CVZ), where targets are observed in every Sector. TIC-25194908 (candidate 8) was observed in Sectors 1-13 during Cycle 1, and Sectors 27-38 in Cycle 3, for a total of 24 Sectors, with only two transits detected. TIC-294097549 (candidate 62) was observed in multiple Sectors across Cycle 1 and Cycle 3 for a total of 13 Sectors, also with only two transits detected. This additional *TESS* coverage rules out most alias periods, and only 2 and 3 periods are possible respectively for these candidates. These are likely to be among the longest periods from our sample, with the remaining allowed periods in the range 337–675 d and 299–599 d respectively (see Table A3).

TIC-349091983 (candidate 74) was observed in all Sectors in Cycle 1 with the exception of Sector 5, and again in Cycle 3 in all Sectors with the exception of Sector 35, for a total of 24 Sectors. There is only one possible period for this candidate of approximately ~ 549 days, and thus the true orbital period is solved from *TESS* alone.

4.3.2 TOI and CTOI candidates

TESS Objects of Interest (TOIs) are planet candidates that have been vetted by the *TESS* Science Office (TSO), the *TESS* Follow-up Observing Program (TFOP) and designated as promising candidates for follow-up and publication. 11 of our total 85 candidates have previously been flagged as *TESS* Objects of Interest (TOI), and these are labelled in Table A1.

Another 14 of our 85 candidates have previously been flagged as Community *TESS* Objects of Interest (CTOIs), and these are also labelled in Table A1. CTOIs are typically planet candidates submitted by additional community projects outside of the official TOI planet search pipelines, and CTOIs are reviewed by the *TESS* TOI team before being promoted to TOI status and assigned a number.

The majority of CTOIs have been identified by the *WINE* (Warm gIaNTs with *tEss*) collaboration (e.g. Schlecker et al. 2020; Jordán et al. 2020; Hobson et al. 2021), the *PHT* (Planet Hunters *TESS*) project (Eisner et al. 2021), and the *STF* (Single-Transit Finder) project.

4.3.3 Confirmed and published candidates

Two of our duotransit planet candidates have already been confirmed and published. TOI-5153 (candidate 25; TIC-124029677) has been published by Ulmer-Moll et al. (2022). The planet is a large warm Jupiter ($M = 3.26^{+0.18}_{-0.17} M_{\text{J}}$, $R = 1.06^{+0.04}_{-0.04} R_{\text{J}}$) orbiting an F8-type star on a period of 20.33 days. We find that our fitted planetary radius from




Figure 6. *Gaia* DR3 ([Gaia Collaboration 2022](#)) colour-magnitude diagram (corrected for extinction and reddening) showing all stars with *TESS* full frame image SPOC lightcurves from Cycle 1 and Cycle 3 that have measured *Gaia* DR3 distances. Blue circles indicate the positions of our 85 duotransit candidates. The colourbar shows the relative number density of stars at each location in the diagram. For six of the duotransit candidates there was no *Gaia* DR3 distance measurement (see [Table A1](#)) so we estimate the distance from the parallax and do not correct for reddening or extinction.




Figure 7. The distribution of total transit event SNR values (see [Section 4.1](#)) for our 85 duotransit candidates.

[Table A3](#) is in agreement with this published value. [TIC-466206508](#) (candidate 85; [TOI-5542](#)) has also been published by [Grievess et al. \(2022\)](#). The planet is an old warm Jupiter orbiting a relatively metal-poor G-dwarf host star on a period of 75.12 days, with a mass of $M = 1.32^{+0.10}_{-0.10} M_J$ and a radius of $R = 1.01^{+0.04}_{-0.04} R_J$. Other publica-

tions for these solved systems are forthcoming, e.g. [TIC-77437543](#) ([Henderson et al. 2023](#), in preparation), [TIC-333736132](#) ([Kendall et al. 2023](#), in preparation), and [TIC-224279805](#).

4.4 Previous TOIs and CTOIs identified as false positives

Some of the candidates that were detected in our transit search but rejected as false positives have previously been announced as TOIs or CTOIs. In all cases, we found these to be examples of passing asteroids that caused peaks in the SAP background time series and hence false dips in the target light curve (see [Section 3.2](#)). These false positives are [TOI-4312](#) ([TIC-251086776](#)) and the CTOIs [TIC-152070435](#) and [TIC-275180352](#).

5 DISCUSSION

From our initial set of 1,422,473 unique TIC IDs, 9718 were identified as duotransit candidates from our `monofind` algorithm in the first instance, and after a quick visual inspection to reject obvious variable stars this was reduced to 736 candidates pre-vetting (see [Section 3.2](#)). From these candidates, 651 were determined to be caused by false-positive scenarios including blended sources, asteroid crossing events, eclipsing binaries, systematics and stellar variability. This leads to a final list of 85 duotransit candidates presented in this work (see [Section 4](#) and [Tables A1-A3](#)).




Figure 8. The distribution of transit depths for our duotransit candidates (green) and the confirmed *TESS* discovered exoplanets from the Southern Ecliptic Hemisphere (grey).




Figure 10. The distribution of transit durations for our duotransit candidates (green) and the confirmed *TESS* discovered exoplanets from the Southern Ecliptic Hemisphere (grey).




Figure 9. The distribution of planet radii for our duotransit candidates (green) and the confirmed *TESS* discovered exoplanets from the Southern Ecliptic Hemisphere (grey).

We have submitted all of our non-TOI/CTOI duotransit planet candidates to the *TESS* follow-up program (TFOP) (60 candidates). We hope that this will allow the community to follow-up these interesting systems further with the aim of determining their true orbital periods and providing confirmation and full characterisations of new long-period planets.

5.1 Duotransit candidate properties

We plot the transit depths of our 85 duotransit candidates in Figure 8, and for comparison we overlay the *TESS* planets that have been confirmed in the Southern Ecliptic Hemisphere⁶. The transit depths




Figure 11. The distribution of *TESS* magnitudes for our duotransit candidate host stars (green) and the confirmed *TESS* discovered exoplanet host stars from the Southern Ecliptic Hemisphere (grey).

of our duotransit candidates range from approximately 0.1 percent to 1.8 percent, peaking around 0.7 percent. This is significantly deeper than is typical for the confirmed *TESS* exoplanets, the majority of which are less than 0.2 percent. This depth difference is to be expected given that the `monofind` algorithm is designed to detect individual high-SNR transits, which will naturally be deeper than the general TOI population.

In Figure 9 we plot the best-fitting radii of our duotransit planet candidates, again in comparison with the *TESS* confirmed planet sample. We find that our sample peaks at around the radius of Jupiter ($11 R_{\oplus}$), which is also to be expected for a sample on individual high-SNR transits. Our sample extends to radii exceeding $16 R_{\oplus}$, and while the sample of confirmed *TESS* planets also extends to similar radii, these are hot Jupiters with radii inflated by irradiation, which we would not expect for long-period planets (Gu et al. 2003). Nevertheless, rather than applying an arbitrary upper radius cut, we

⁶ NASA Exoplanet Archive; <https://exoplanetarchive.ipac.caltech.edu/>, accessed 2023 Jul 15




Figure 12. The distribution of *Gaia* DR3 distances (in parsecs) for our duotransit candidate host stars.




Figure 13. The distribution of most probable periods from modelling with *MonoTools* for our duotransit candidates.

keep these candidates in our sample for completeness and will rely on follow-up observations to determine the upper radius limit of long-period planets.

Figure 10 shows that most of our candidates have significantly longer transit durations than many of the confirmed *TESS* planets. This is expected as we are probing planets with orbital periods in excess of ~ 17 days, which is significantly longer than the median orbital period of *TESS* planets (~ 5.8 days).

In Figure 11 we plot the *TESS* magnitudes of the host stars of our candidates, again comparing with the sample of confirmed *TESS* planets. Here it can be seen that our long-period planet candidates tend to orbit stars that are significantly fainter than the confirmed planet sample (typically T_{mag} of 10–13). This is for three reasons. First, the confirmed *TESS* sample is biased towards bright stars, partly because brighter stars make it easier to detect shallow transits, and partly because follow-up efforts tend to be focused on bright host stars. Second, the much smaller transit probabilities of wide-

separation planets means that long-period transiting candidates are rare, and so the brightest detected examples will be around fainter stars, which are more numerous (the absence of brighter examples is not a selection effect because such systems would readily be detected). Third, gas giant planets are inherently rarer than the sub-Neptune sized planets that make up the bulk of the confirmed *TESS* sample. The steep cut-off in our detected candidates at $T_{\text{mag}}=13$ simply reflects the selection criteria for the SPOC FFI lightcurves (Caldwell et al. 2020).

In Figure 12 we plot the distances of our duotransit candidate host stars from the *Gaia* DR3 catalog. We find that most candidate host stars are at distances from 200–500 pc, with the furthest object being TIC-218977148 at a distance of ~ 826 pc, and the closest being TIC-303317324/TOI-4310 at a distance of ~ 40 pc.

As described in Section 3.3, we use the shape and width of our detected transit events to determine the most probable orbital period for each candidate (e.g. Osborn et al. 2022b). The probability of each allowed alias is indicated on Figure 4 and the most probable period for each candidate is listed in Table A3. We plot a histogram of these most probable periods in Figure 13. We find that our distribution peaks at periods around 20–30 days. While these values are the most probable periods according to the information and data available, they are not necessarily the correct values, and the true orbital periods must be determined through follow-up observations. We also note the three candidates in our sample with the longest periods: TIC-294097549 with a value of P_{marg} of 200 days, TIC-25194908 with a value of P_{marg} of 338 days, and TIC-349091983 with a value of P_{marg} of 549 days (see Table A3).

5.2 Follow-up programme

5.2.1 Photometric period determination

The limited number of discrete period aliases for duotransit candidates means that precise orbital periods can be determined efficiently with photometric follow-up targeted at predicted transit times for each alias.

Our own team is actively following up the duotransit candidates presented here using the Next Generation Transit Survey (*NGTS*; Wheatley et al. 2018). *NGTS* is a ground-based facility located at the ESO Paranal Observatory in Chile that deploys an array of twelve telescopes optimised for photometric detection of exoplanet transits (e.g. Bryant et al. 2020). Our monotransit programme with *NGTS* confirmed the first planet that was initially identified as a single-transit event with *TESS* (Gill et al. 2020c), and has since confirmed several other long-period planets (e.g. Ulmer-Moll et al. 2022; Grieves et al. 2022, Gill et al., submitted). We have also confirmed three long-period, low-mass eclipsing binary systems (Gill et al. 2020a; Lendl et al. 2020b; Gill et al. 2020b).

We are also searching *TESS* Cycle 5 data as it is released for third transits of our duotransit candidates. In some cases this will significantly reduce the number of period aliases to search, although with a similar two-year gap between Cycles 1, 3 and 5 a third transit usually still leaves multiple aliases to be searched.

5.2.2 Radial velocity detections

Our team has also begun a spectroscopic campaign using the *CORALIE* spectrograph (Queloz et al. 1999) on the Swiss 1.2 m Leonhard Euler Telescope at La Silla Observatory (program ID 500) to help confirm our candidates. With a first *CORALIE* spectrum, we are able to identify and remove double-line spectroscopic binary

(SB2) systems, and detect rapidly rotating stars from the broadening of the cross-correlation function. With a second *CORALIE* spectrum, taken about one week later, we are able to exclude systems for which the change of measured radial velocity is too large to be consistent with a planetary companion ($>1 \text{ km s}^{-1}$). Bright targets hosting relatively massive planetary companions exhibit radial velocity variations detectable with *CORALIE* ($\sim 6 \text{ m s}^{-1}$ for bright stars; e.g. Marmier et al. 2013) and for other targets we use *HARPS*. Previous efforts to confirm these mono- and duotransiting planets using spectroscopic observations have been successful (e.g. Ulmer-Moll et al. 2022; Grieves et al. 2022; Ulmer-Moll et al. 2023).

6 CONCLUSION

We have presented the discovery of 85 duotransit candidates from the *TESS*-SPOC FFI light curves from the Southern Ecliptic Hemisphere. The candidates were found using our custom-built *monofind* algorithm through which all SPOC pipeline data products were passed. We also employed a vetting procedure to rule out false-positive scenarios such as eclipsing binaries and asteroids in the sample. A total of 25 of our candidates have previously been flagged as either TOIs or CTOIs, and we found during our vetting process 3 TOI/CTOI objects outside of our duotransits falsely flagged as such that are asteroids. A dedicated program on the *NGTS* facility is now being used to determine the true orbital periods of these systems, and we welcome other community efforts in this difficult task. We also have programs on instruments such as *CORALIE* and *HARPS* to acquire radial velocity measurements to confirm the planetary nature of these systems and to determine the mass of the exoplanets. Once confirmed, these planets will form the basis for characterisation studies into their atmospheric compositions and orbital obliquities, and hence into the formation, evolution and migration of warm and cool gas giant exoplanets.

ACKNOWLEDGEMENTS

This paper includes data collected by the *TESS* mission. Funding for the *TESS* mission is provided by the NASA Explorer Program. Resources supporting this work were provided by the NASA High-End Computing (HEC) Program through the NASA Advanced Supercomputing (NAS) Division at Ames Research Center for the production of the SPOC data products. The *TESS* team shall assure that the masses of fifty (50) planets with radii less than 4 REarth are determined.

This work makes use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement.

We acknowledge the use of public *TESS* Alert data from pipelines at the *TESS* Science Office and at the *TESS* Science Processing Operations Center.

This research makes use of the Exoplanet Follow-up Observation Program website, which is operated by the California Institute of Technology, under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program.

This paper includes data collected by the *TESS* mission that are publicly available from the Mikulski Archive for Space Telescopes (MAST).

FH, TR and AO are supported by STFC studentships. ML acknowledges support of the Swiss National Science Foundation under grant number PCEFP2_194576. The contribution of MB, ML, HPO, & SUM have been carried out within the framework of the NCCR PlanetS supported by the Swiss National Science Foundation under grants 51NF40_182901 and 51NF40_205606. JSJ acknowledges support by Fondecyt grant 1201371 and by the ANID BASAL project FB210003. IP acknowledges a Warwick Astrophysics prize post-doctoral fellowship made possible thanks to a generous philanthropic donation. The contributions at the University of Warwick by SG, DB, PJW, RGW and DA have been supported by STFC through consolidated grants ST/P000495/1, ST/T000406/1 and ST/X001121/1.

DATA AVAILABILITY

The *TESS* data is accessible via the MAST (Mikulski Archive for Space Telescopes) portal at <https://mast.stsci.edu/portal/Mashup/Clients/Mast/Portal.html>. The associated python scripts for analysis and plotting are available upon reasonable request to the authors.

REFERENCES

- Akeson R. L., et al., 2013, *PASP*, **125**, 989
 Albrecht S., et al., 2012, *ApJ*, **757**, 18
 Borucki W. J., et al., 2010, *Science*, **327**, 977
 Bryant E. M., et al., 2020, *MNRAS*, **499**, 3139
 Bryant E. M., Bayliss D., Van Eylen V., 2023, *MNRAS*, **521**, 3663
 Caldwell D. A., et al., 2020, *Research Notes of the American Astronomical Society*, **4**, 201
 Castelli F., Kurucz R. L., 2003, in Piskunov N., Weiss W. W., Gray D. F., eds, International Astronomical Union 210th Symposium Posters Vol. 210, Modelling of Stellar Atmospheres. p. A20 ([arXiv:astro-ph/0405087](https://arxiv.org/abs/astro-ph/0405087)), doi:10.48550/arXiv.astro-ph/0405087
 Claret A., 2017, *A&A*, **600**, A30
 Cooke B. F., Pollacco D., West R., McCormac J., Wheatley P. J., 2018, *A&A*, **619**, A175
 Cooke B. F., et al., 2021, *MNRAS*, **500**, 5088
 Culpan R., Geier S., Reindl N., Pelisoli I., Gentile Fusillo N., Vorontseva A., 2022, *A&A*, **662**, A40
 Dalba P. A., et al., 2020, *AJ*, **159**, 241
 Eisner N. L., et al., 2021, *MNRAS*, **501**, 4669
 Espinoza N., Jordán A., 2016, *MNRAS*, **457**, 3573
 Foreman-Mackey D., Agol E., Ambikasaran S., Angus R., 2017, *Astrophysics Source Code Library*, pp ascl-1709
 Foreman-Mackey D., et al., 2021, *exoplanet-dev/exoplanet* v0.4.5, doi:10.5281/zenodo.1998447, <https://doi.org/10.5281/zenodo.1998447>
 Gaia Collaboration 2022, *VizieR Online Data Catalog*, p. I/355
 Gill S., et al., 2020a, *MNRAS*, **491**, 1548
 Gill S., et al., 2020b, *MNRAS*, **495**, 2713
 Gill S., et al., 2020c, *ApJ*, **898**, L11
 Grieves N., et al., 2022, *arXiv e-prints*, p. [arXiv:2209.14830](https://arxiv.org/abs/2209.14830)
 Gu P.-G., Lin D. N. C., Bodenheimer P. H., 2003, *ApJ*, **588**, 509
 Han Z., Podsiadlowski P., Maxted P. F. L., Marsh T. R., Ivanova N., 2002, *MNRAS*, **336**, 449
 Han Z., Podsiadlowski P., Maxted P. F. L., Marsh T. R., 2003, *MNRAS*, **341**, 669
 Hattori S., Foreman-Mackey D., Hogg D. W., Montet B. T., Angus R., Pritchard T. A., Curtis J. L., Schölkopf B., 2022, *AJ*, **163**, 284
 Hobson M. J., et al., 2021, *AJ*, **161**, 235
 Hsu D. C., Ford E. B., Ragozzine D., Ashby K., 2019, *AJ*, **158**, 109
 Jeffery C. S., Miszalski B., Snowden E., 2021, *MNRAS*, **501**, 623

- Jenkins J. M., et al., 2016, in *Software and Cyberinfrastructure for Astronomy* IV. p. 99133E, doi:10.1117/12.2233418
- Jordán A., et al., 2020, *AJ*, 159, 145
- Kipping D. M., 2013, *MNRAS*, 434, L51
- Kipping D., 2023, *MNRAS*, 523, 1182
- Lendl M., et al., 2020a, *MNRAS*, 492, 1761
- Lendl M., et al., 2020b, *MNRAS*, 492, 1761
- Lopez E. D., Fortney J. J., 2016, *ApJ*, 818, 4
- Madhusudhan N., Bitsch B., Johansen A., Eriksson L., 2017, *MNRAS*, 469, 4102
- Marmier M., et al., 2013, *A&A*, 551, A90
- Maxted P. F. L., Heber U., Marsh T. R., North R. C., 2001, *MNRAS*, 326, 1391
- Osborn H. P., 2022, MonoTools: Planets of uncertain periods detector and modeler, Astrophysics Source Code Library, record ascl:2204.020 (ascl:2204.020)
- Osborn H. P., et al., 2021, *MNRAS*, 502, 4842
- Osborn H. P., et al., 2022a, *A&A*, 664, A156
- Osborn H. P., et al., 2022b, *A&A*, 664, A156
- Pelisoli I., Vos J., Geier S., Schaffenroth V., Baran A. S., 2020, *A&A*, 642, A180
- Queloz D., et al., 1999, *arXiv e-prints*, pp astro-ph/9910223
- Rauch T., Deetjen J. L., 2003, in Hubeny I., Mihalas D., Werner K., eds, *Astronomical Society of the Pacific Conference Series Vol. 288, Stellar Atmosphere Modeling*. p. 103 (arXiv:astro-ph/0403239), doi:10.48550/arXiv.astro-ph/0403239
- Rice M., et al., 2022, *AJ*, 164, 104
- Ricker G. R., et al., 2015, *Journal of Astronomical Telescopes, Instruments, and Systems*, 1, 014003
- Salvatier J., Wiecki T. V., Fonnesbeck C., 2016, *PeerJ Computer Science*, 2, e55
- Schaffenroth V., Pelisoli I., Barlow B. N., Geier S., Kupfer T., 2022, *A&A*, 666, A182
- Schlecker M., et al., 2020, *AJ*, 160, 275
- Seabroke G. M., et al., 2021, *A&A*, 653, A160
- Smith J. C., et al., 2012, *PASP*, 124, 1000
- Stark M. A., Wade R. A., 2003, *AJ*, 126, 1455
- Stassun K. G., et al., 2019, *AJ*, 158, 138
- Stumpe M. C., et al., 2012, *PASP*, 124, 985
- Stumpe M. C., Smith J. C., Catanzarite J. H., Van Cleve J. E., Jenkins J. M., Twicken J. D., Girouard F. R., 2014, *PASP*, 126, 100
- Temple L. Y., et al., 2018, *MNRAS*, 480, 5307
- Temple L. Y., et al., 2019, *MNRAS*, 490, 2467
- Thuillier A., Van Grootel V., Dévora-Pajares M., Pozuelos F. J., Charpinet S., Siess L., 2022, *arXiv e-prints*, p. arXiv:2209.02437
- Twicken J., et al., 2019, in *American Astronomical Society Meeting Abstracts #233*. p. 140.03
- Ulmer-Moll S., et al., 2022, *A&A*, 666, A46
- Ulmer-Moll S., et al., 2023, *A&A*, 674, A43
- Van Grootel V., et al., 2021, *A&A*, 650, A205
- Villanueva Steven J., Dragomir D., Gaudi B. S., 2019, *AJ*, 157, 84
- Vos J., Vučković M., Chen X., Han Z., Boudreaux T., Barlow B. N., Østensen R., Németh P., 2019, *MNRAS*, 482, 4592
- Wang S., et al., 2021, *AJ*, 162, 50
- Watson C. A., de Mooij E. J. W., Steeghs D., Marsh T. R., Brogi M., Gibson N. P., Matthews S., 2019, *MNRAS*, 490, 1991
- Werner K., Drezler S., 1999, *Journal of Computational and Applied Mathematics*, 109, 65
- Werner K., Deetjen J. L., Drezler S., Nagel T., Rauch T., Schuh S. L., 2003, in Hubeny I., Mihalas D., Werner K., eds, *Astronomical Society of the Pacific Conference Series Vol. 288, Stellar Atmosphere Modeling*. p. 31 (arXiv:astro-ph/0209535), doi:10.48550/arXiv.astro-ph/0209535
- Wheatley P. J., et al., 2018, *MNRAS*, 475, 4476

Table A1: Properties of the host stars of our duotransit candidates.

ID	TIC ID	TOI/CTOI	RA (deg)	Dec (deg)	T _{mag}	R _* (R _☉)	T _{eff} (K)	Distance (pc)	N _{sec}
1	1504460	–	131.9658	-23.9895	11.63	1.5	6218	451.71 ^{+20.73} _{-15.90}	2
2	2669681	–	289.368	-28.5803	11.47	1.88	5160	382.06 ^{+12.79} _{-9.96}	2
3	9844069	–	48.03923	-11.6808	12.96	1.61	5665	777.93 ^{+11.74} _{-11.59}	2
4	13072758	CTOI	76.08837	-29.0342	12.22	1.43	6313	601.42 ^{+2.95} _{-3.19}	2
5	13713700	–	133.6396	-7.94134	12.29	0.97	5864	391.66 ^{+2.67} _{-2.25}	2
6	14445414	–	110.8209	12.66321	11.02	1.31	6425	346.17 ^{+12.45} _{-3.76}	2
7	20904104	–	7.41368	-19.3663	9.76	0.88	5628	105.79 ^{+0.24} _{-0.22}	2
8	25194908	–	63.90318	-66.3519	12.46	1.32	6309	666.61 ^{+32.34} _{-8.91}	24
9	32179255	–	332.0396	-29.6093	10.52	1.75	5989	330.44 ^{+14.95} _{-7.29}	2
10	38138512	–	170.4183	-4.19721	12.94	1.34	6300	769.66 ^{+10.80} _{-9.15}	2
11	39167176	–	88.32738	-3.79767	12.01	1.56	8416	486.84 ^{+3.95} _{-1.46}	2
12	42428568	–	96.29347	-3.76657	10.83	2.16	6686	567.87 ^{+8.12} _{-10.63}	2
13	52195587	–	19.24834	-66.1636	12.25	1.2	5736	450.99 ^{+2.50} _{-2.59}	4
14	66439839	–	14.47851	-30.3231	11.97	1.03	5882	356.25 ^{+2.47} _{-3.35}	3
15	67599025	CTOI	17.69268	-31.5055	11.96	0.78	4816	177.82 ^{+1.19} _{-1.34}	2
16	70561926	–	139.2559	-26.7007	11	1.35	6386	353.55 ^{+5.09} _{-3.76}	2
17	71028120	–	66.12741	-11.9835	10.44	1.37	5596	215.31 ^{+0.82} _{-0.84}	2
18	77437543	TOI-2490	73.12441	-36.2572	11.28	1.17	5459	264.92 ^{+1.04} _{-0.96}	3
19	81089255	CTOI	122.6942	-43.8535	11.84	1.43	7459	547.70 ^{+2.71} _{-2.72}	4
20	100776118	CTOI	88.31784	-34.5185	11.51	1.25	6439		4
21	101824521	–	206.004	-26.4978	12.61	1.23	5553	481.79 ^{+4.21} _{-3.97}	2
22	107113345	–	183.0341	-32.6628	10.34	1.31	6696	267.74 ^{+1.41} _{-1.63}	2
23	118339710	CTOI	129.572	-23.5502	10.35	1.56	6119	285.48 ^{+15.23} _{-4.22}	3
24	123763494	–	90.30269	-21.2359	11.73	1.22	5568	353.10 ^{+29.67} _{-9.36}	2
25	124029677	TOI-5153	91.54028	-19.9533	11.21	1.38	6408	366.92 ^{+2.94} _{-2.71}	2
26	140215502	CTOI	85.15071	-34.7664	11.23	0.93	5528	210.03 ^{+0.62} _{-0.56}	4
27	140750416	–	116.4382	-23.143	11.81	0.87	5331	238.74 ^{+5.45} _{-2.41}	2
28	142278054	–	180.7626	-33.343	10.28	2.39	6400	418.42 ^{+1.64} _{-2.10}	2
29	145006304	–	193.3345	-39.7347	11.3	1.12	6340	309.68 ^{+1.88} _{-1.72}	2
30	145913596	–	123.1298	-30.5555	10.17	1.45	6467	261.13 ^{+1.63} _{-0.99}	4
31	153838604	–	68.07396	-47.6088	12.48	0.91	5649	387.07 ^{+22.34} _{-7.33}	4
32	156716001	–	103.1063	-39.6876	10.9	1.08	6129	240.55 ^{+2.60} _{-11.28}	4
33	157119927	–	175.2155	-31.1374	11.64	1.04	5765	299.69 ^{+8.62} _{-2.98}	2
34	157698565	TOI-2589	107.4883	-37.231	10.72	1.07	5938	198.90 ^{+23.29} _{-1.84}	3
35	159490807	–	220.2349	-39.3757	10.57	1.69	6556	341.73 ^{+3.68} _{-3.50}	2
36	161169240	CTOI	339.3573	-53.3188	12.12	1.19	5952	445.30 ^{+6.33} _{-36.66}	2
37	176518126	–	87.1832	-0.49503	10.48	1.36	7927	281.19 ^{+1.57} _{-1.56}	2
38	188620407	CTOI	350.0511	-13.0494	11.57	1.31	6169	404.71 ^{+3.12} _{-2.81}	3
39	193318850	–	177.7302	-16.7417	12.76	0.79	5560	340.21 ^{+1.78} _{-1.68}	2
40	207783865	–	207.0609	-54.4206	10.54	1.04	5936	180.65 ^{+2.63} _{-0.98}	2
41	211409161	–	321.7768	-30.5273	10.44	0.78	4660		2
42	215402824	–	43.0624	-36.2887	12.97	1.33	6180	761.55 ^{+11.13} _{-8.01}	4
43	218977148	–	349.5331	-33.1369	12.58	1.61	6214	826.08 ^{+12.07} _{-12.51}	2
44	220622886	–	84.89306	-42.6811	12	2.56	5671	796.01 ^{+3.48} _{-16.0}	3
45	221915858	–	249.513	-54.3579	10.07	2.47	9475		2

Table A1: Properties of the host stars of our duotransit candidates.

ID	TIC ID	TOI/CTOI	RA (deg)	Dec (deg)	T _{mag}	R _* (R _☉)	T _{eff} (K)	Distance (pc)	N _{sec}
46	224279805	CTOI	355.7386	-40.7722	12.15	1.2	5591	422.59 ^{+3.22} _{-3.1}	2
47	235058563	–	83.51699	-50.1216	12.43	1.47	6438	706.77 ^{+5.66} _{-5.64}	6
48	237605045	–	103.2896	2.05465	10.29	2.92	6577	471.34 ^{+7.62} _{-7.43}	2
49	242241304	CTOI	211.5851	-43.0447	11.81	1.14	5781	346.35 ^{+1.79} _{-1.77}	2
50	251057075	–	36.17507	-5.85401	11.05	0.84	5225	157.35 ^{+0.52} _{-0.54}	2
51	256912435	–	200.5462	-55.2184	11.8	1.19	5650	333.69 ^{+3.17} _{-3.45}	2
52	265465927	CTOI	320.9166	-62.9238	12.08	1.16	6099	418.77 ^{+65.41} _{-56.26}	2
53	265466589	–	321.0794	-64.4743	12.61	1.03	5834		2
54	268534931	CTOI	15.12618	-23.1795	12.22	0.78	5179	246.72 ^{+7.21} _{-2.25}	2
55	269333648	TOI-2529	118.9944	-52.3549	10.67	1.75	5822	295.40 ^{+2.04} _{-1.85}	6
56	275878706	–	207.9018	-44.3144	12.17	0.93	5038	247.70 ^{+2.42} _{-2.18}	2
57	279727635	–	50.59176	-64.4444	12.5	1.05	5851	447.75 ^{+3.66} _{-3.51}	6
58	287204963	–	30.28722	-11.8835	13.25	1.04	5344	551.78 ^{+10.06} _{-5.87}	2
59	289840544	–	317.5455	-26.3999	10.64	1.75	5989	329.46 ^{+3.72} _{-3.65}	2
60	290165539	–	319.8521	-25.5971	11.99	1.33	5895	453.86 ^{+6.32} _{-15.61}	2
61	292719109	–	107.416	3.56756	10.1	2.08	6216	326.24 ^{+9.07} _{-9.63}	2
62	294097549	–	107.67	-55.7967	10.47	1.15	5872	197.78 ^{+0.46} _{-0.5}	13
63	296737508	CTOI	142.1793	-14.6864	9.2	1.27	5700	117.37 ^{+0.23} _{-0.22}	2
64	300394149	–	170.747	-52.6986	10.13	1.42	5997	218.59 ^{+0.60} _{-0.59}	2
65	303317324	TOI-4310	351.9037	-25.5081	9.5	0.72	4159	40.13 ^{+0.43} _{-0.10}	2
66	304339227	TOI-4629	275.9439	-68.3449	8.66	1.13	6012	91.13 ^{+0.23} _{-0.15}	2
67	306249066	–	166.7285	-54.7388	10.54	1.14	6139	221.63 ^{+1.01} _{-2.35}	2
68	313671132	–	202.1582	-62.1699	10.08	1.22	5878	175.65 ^{+0.41} _{-0.43}	2
69	317923092	–	89.85183	-14.9061	12.26	1.34	7604	668.65 ^{+21.27} _{-10.4}	2
70	323295479	TOI-1861	130.7987	-83.061	9.95	1.06	5684	134.32 ^{+0.24} _{-0.23}	5
71	332697924	–	63.26121	-12.4593	11	1.32	6266	312.19 ^{+3.28} _{-9.2}	2
72	333736132	CTOI	170.2765	-26.0666	12.07	0.76	4722	178.23 ^{+0.58} _{-1.38}	2
73	339399841	–	296.0705	-62.8136	12.35	1.55	5686	596.49 ^{+4.8} _{-5.11}	2
74	349091983	–	107.7051	-61.3524	12.29	1.5	6083		24
75	381553868	–	269.7743	-50.9212	11.85	1.59	6355		2
76	393229954	–	86.74486	-46.7119	11.77	0.9	5571	262.19 ^{+5.47} _{-1.15}	5
77	394340183	–	39.32208	-79.4583	10.95	1.04	6325	253.84 ^{+0.64} _{-0.65}	2
78	396720998	TOI-709	65.65532	-54.1472	13.99	0.15	50000		6
79	404518509	TOI-4320	51.64036	-43.6136	8.6	1.04	5871	79.30 ^{+0.07} _{-0.07}	4
80	412386707	–	110.4871	-23.414	10.92	1.96	6304	425.23 ^{+5.01} _{-4.39}	2
81	437293313	–	191.2234	-22.6177	12.97	0.94	5536	416.28 ^{+7.26} _{-6.35}	2
82	439491923	–	21.38365	-18.1471	12.87	1.26	5944	631.09 ^{+11.04} _{-9.74}	2
83	442893494	–	80.00938	-16.2193	12.41	1.39	6322	622.15 ^{+5.22} _{-5.04}	3
84	457649900	TOI-4958	217.9991	-51.0744	10.75	2.11	6713	479.53 ^{+4.71} _{-3.72}	2
85	466206508	TOI-5542	302.7985	-61.1355	11.85	1.27	5393	342.66 ^{+1.69} _{-1.51}	2

Table A2: The transit event properties of our duotransit candidates.

ID	TIC ID	T_{c1}	T_{c2}	Depth	Duration
—	—	(TBJD, [Sector])	(TBJD, [Sector])	(ppt)	(hours)
1	1504460	1528.39739 [08]	2261.11990 [35]	3.367 [08], 3.415 [35]	3.78
2	2669681	1676.73302 [13]	2044.45589 [27]	7.879 [13], 7.037 [27]	9.69
3	9844069	1431.16511 [04]	2148.92146 [31]	13.452 [04], 12.224 [31]	7.20
4	13072758	1455.59404 [05]	2193.21295 [32]	7.023 [05], 6.755 [32]	9.25
5	13713700	1536.98226 [08]	2245.82903 [34]	7.234 [08], 4.700 [34]	3.26
6	14445414	1497.55590 [07]	2210.97620 [33]	2.317 [07], 2.179 [33]	2.86
7	20904104	1390.11189 [03]	2105.36177 [29]	0.738 [03], 1.065 [29]	4.61
8	25194908	1402.29491 [03]	2077.28343 [28]	6.590 [03], 6.266 [28]	9.16
9	32179255	1345.87922 [01]	2068.45418 [28]	3.619 [01], 4.682 [28]	3.71
10	38138512	1559.95080 [09]	2300.02560 [36]	17.907 [09], 18.148 [36]	6.73
11	39167176	1470.62312 [06]	2211.42630 [33]	7.282 [06], 8.112 [33]	6.25
12	42428568	1475.69109 [06]	2224.32460 [33]	5.380 [06], 4.939 [33]	10.20
13	52195587	1350.27811 [01]	2107.61083 [29]	5.775 [01], 6.138 [29]	9.07
14	66439839	1386.34406 [03]	2137.03203 [30]	6.227 [03], 5.420 [30]	2.16
15	67599025	1404.02101 [03]	2138.82557 [30]	6.684 [03], 7.345 [30]	3.27
16	70561926	1523.06353 [08]	2275.36040 [35]	16.290 [08], 16.053 [35]	5.91
17	71028120	1445.81123 [05]	2189.95412 [32]	1.238 [05], 1.025 [32]	6.11
18	77437543	1456.70241 [05]	2180.69226 [32]	9.825 [05], 9.326 [32]	7.93
19	81089255	1509.94498 [07]	2260.20690 [35]	7.318 [07], 6.372 [35]	5.14
20	100776118	1472.81029 [06]	2219.67770 [33]	13.264 [06], 13.070 [33]	22.34
21	101824521	1620.35423 [11]	2331.52820 [37]	9.396 [11], 9.103 [37]	8.52
22	107113345	1576.32832 [10]	2317.29675 [37]	1.771 [10], 2.035 [37]	1.77
23	118339710	1525.64667 [08]	2249.54641 [34]	4.329 [08], 3.977 [34]	8.68
24	123763494	1486.92891 [06]	2209.76758 [33]	10.645 [06], 11.728 [33]	2.47
25	124029677	1486.12090 [06]	2218.00409 [33]	6.368 [06], 6.105 [33]	4.58
26	140215502	1460.54584 [05]	2194.44033 [32]	2.959 [05], 4.468 [32]	7.14
27	140750416	1504.80151 [07]	2245.39501 [34]	8.912 [07], 9.197 [34]	3.85
28	142278054	1595.02504 [10]	2310.35369 [37]	2.177 [10], 2.022 [37]	4.82
29	145006304	1579.50384 [10]	2318.24443 [37]	2.048 [10], 2.772 [37]	5.19
30	145913596	1539.51696 [08]	2242.69675 [34]	1.728 [08], 2.163 [34]	8.50
31	153838604	1447.10227 [05]	2176.13853 [32]	15.209 [05], 16.237 [32]	2.61
32	156716001	1485.40631 [06]	2224.51833 [33]	2.652 [06], 3.037 [33]	9.11
33	157119927	1591.62175 [10]	2291.00613 [36]	5.445 [10], 6.897 [36]	2.57
34	157698565	1494.57973 [07]	2234.11360 [34]	9.825 [07], 9.338 [34]	6.38
35	159490807	1606.22551 [11]	2355.93796 [38]	5.155 [11], 5.184 [38]	5.38
36	161169240	1328.25945 [01]	2083.31754 [28]	9.707 [01], 10.134 [28]	7.82
37	176518126	1489.05533 [06]	2219.53761 [33]	3.565 [06], 3.244 [33]	5.61
38	188620407	1375.48660 [02]	2461.70715 [42]	5.467 [02], 4.512 [42]	6.08
39	193318850	1553.77031 [09]	2292.51011 [36]	6.916 [09], 7.143 [36]	3.89
40	207783865	1608.93692 [11]	2354.11345 [38]	17.211 [11], 17.729 [38]	1.99
41	211409161	1335.10057 [01]	2070.36583 [28]	13.629 [01], 12.232 [28]	2.14
42	215402824	1413.62655 [04]	2169.04205 [31]	5.589 [04], 9.955 [31]	1.87
43	218977148	1355.47069 [02]	2090.09772 [29]	5.597 [02], 4.944 [29]	8.13
44	220622886	1479.52544 [06]	2204.58502 [33]	4.324 [06], 3.501 [33]	4.43

Table A2: The transit event properties of our duotransit candidates.

ID	TIC ID	T _{e1}	T _{e2}	Depth	Duration
—	—	(TBJD, [Sector])	(TBJD, [Sector])	(ppt)	(hours)
45	221915858	1643.35799 [12]	2372.55118 [39]	5.737 [12], 5.526 [39]	3.05
46	224279805	1369.72941 [02]	2105.21571 [29]	6.392 [02], 6.618 [29]	5.93
47	235058563	1416.61066 [04]	2216.44259 [33]	6.164 [04], 5.630 [33]	10.17
48	237605045	1489.04268 [06]	2210.29724 [33]	6.212 [06], 7.014 [33]	4.42
49	242241304	1623.07095 [11]	2345.37403 [38]	6.158 [11], 6.186 [38]	4.72
50	251057075	1426.31846 [04]	2154.02335 [31]	1.986 [04], 2.348 [31]	4.42
51	256912435	1623.43775 [11]	2338.70979 [38]	20.396 [11], 21.562 [38]	2.72
52	265465927	1342.64749 [01]	2058.60913 [27]	8.678 [01], 8.720 [27]	6.05
53	265466589	1352.52439 [01]	2037.86660 [27]	6.818 [01], 6.664 [27]	4.14
54	268534931	1398.29475 [03]	2139.09941 [30]	5.829 [03], 6.133 [30]	7.82
55	269333648	1538.52419 [08]	2249.05416 [34]	4.162 [08], 4.032 [34]	9.46
56	275878706	1621.08010 [11]	2341.78257 [38]	6.544 [11], 5.320 [38]	4.38
57	279727635	1343.22676 [01]	2124.62930 [30]	7.181 [01], 6.789 [30]	5.71
58	287204963	1402.93640 [03]	2132.33438 [30]	11.498 [03], 10.182 [30]	4.73
59	289840544	1331.28977 [01]	2083.28790 [28]	2.702 [01], 1.997 [28]	2.99
60	290165539	1332.75615 [01]	2067.97094 [28]	13.627 [01], 15.056 [28]	11.50
61	292719109	1499.70159 [07]	2222.31933 [33]	2.859 [07], 2.677 [33]	9.08
62	294097549	1657.74149 [13]	2256.34250 [35]	1.977 [13], 1.426 [35]	7.96
63	296737508	1538.00622 [08]	2271.99796 [35]	2.021 [08], 1.854 [35]	5.65
64	300394149	1576.88642 [10]	2325.42588 [37]	2.051 [10], 2.195 [37]	13.40
65	303317324	1365.17582 [02]	2104.34337 [29]	1.360 [02], 1.323 [29]	5.74
66	304339227	1673.33918 [13]	2388.01911 [39]	0.522 [13], 0.464 [39]	5.79
67	306249066	1593.97611 [10]	2326.51397 [37]	7.873 [10], 7.061 [37]	4.03
68	313671132	1604.35452 [11]	2337.72693 [38]	3.408 [11], 3.506 [38]	5.70
69	317923092	1479.81772 [06]	2209.42617 [33]	4.358 [06], 6.715 [33]	12.96
70	323295479	1622.92483 [11]	2365.44036 [39]	8.727 [11], 8.609 [39]	6.51
71	332697924	1445.74977 [05]	2178.22194 [32]	1.042 [05], 1.423 [32]	6.63
72	333736132	1551.55292 [09]	2283.54681 [36]	15.856 [09], 16.440 [36]	2.96
73	339399841	1662.10984 [13]	2043.59486 [27]	4.773 [13], 6.471 [27]	6.66
74	349091983	1603.10253 [11]	2152.16754 [31]	8.301 [11], 8.133 [31]	9.52
75	381553868	1677.32594 [13]	2367.57869 [39]	5.653 [13], 6.030 [39]	6.49
76	393229954	1446.79953 [05]	2204.55168 [33]	8.172 [05], 6.652 [33]	2.67
77	394340183	1664.84261 [13]	2389.11243 [39]	11.255 [13], 10.820 [39]	8.65
78	396720998	1399.21966 [03]	2176.27178 [30]	6.360 [03], 6.082 [30]	4.33
79	404518509	1431.26615 [04]	2134.88294 [30]	0.842 [04], 0.598 [30]	5.02
80	412386707	1493.95365 [07]	2240.57066 [34]	3.442 [07], 3.587 [34]	5.23
81	437293313	1578.44737 [10]	2328.08914 [37]	9.891 [10], 8.462 [37]	6.58
82	439491923	1405.41115 [03]	2116.79038 [30]	11.608 [03], 11.766 [30]	6.70
83	442893494	1472.65885 [06]	2194.38454 [32]	5.776 [06], 6.458 [32]	7.70
84	457649900	1606.96984 [11]	2355.79522 [38]	2.857 [11], 2.945 [38]	7.61
85	466206508	1679.34576 [13]	2054.97133 [27]	11.214 [13], 9.274 [27]	8.03

Table A3: Properties of our duotransit candidates. N_{alias} is the alias number corresponding to the most probable orbital period P_{marg} .

ID	TIC ID	R_p (R_{\oplus})	$P_{\text{min}}, P_{\text{max}}$ (days)	N_{alias}	P_{marg} (days)	b	Gaia NSS	Gaia RV amp (km s^{-1})	Total SNR
1	1504460	9.6 ± 1.0	19.80333, 732.72251	24	$20.0^{+19.0}_{-1.0}$	$0.78^{+0.12}_{-0.37}$	0	13.84	14.82
2	2669681	17.0 ± 1.2	21.63067, 367.72287	18	$24.5^{+8.9}_{-2.9}$	$0.66^{+0.09}_{-0.24}$	0	36.15	41.8
3	9844069	18.9 ± 1.4	22.42991, 717.75635	33	$25.6^{+7.0}_{-3.2}$	$0.62^{+0.12}_{-0.3}$	0	–	46.12
4	13072758	$11.78^{+0.67}_{-0.65}$	19.93567, 737.61891	38	26.0 ± 10.0	0.31 ± 0.22	0	20.36	35.92
5	13713700	8.3 ± 1.0	19.69018, 708.84677	37	$22.9^{+8.0}_{-3.2}$	$0.79^{+0.11}_{-0.28}$	0	–	19.73
6	14445414	$7.5^{+3.9}_{-1.2}$	19.28163, 713.42031	38	$22.3^{+7.4}_{-3.0}$	$0.93^{+0.06}_{-0.18}$	0	4.78	12.08
7	20904104	2.6 ± 0.19	17.88103, 715.24989	41	$23.0^{+11.0}_{-4.0}$	$0.27^{+0.25}_{-0.19}$	0	2.9	11.84
8	25194908	11.18 ± 0.68	337.49490, 674.98980	2	337.49490	$0.7^{+0.1}_{-0.28}$	0	–	31.43
9	32179255	21.0 ± 8.0	20.64500, 722.57496	36	$22.6^{+6.3}_{-1.9}$	$0.999^{+0.07}_{-0.045}$	2	49.01	36.77
10	38138512	17.8 ± 1.0	18.50185, 740.07480	41	$21.1^{+6.3}_{-2.2}$	0.32 ± 0.2	0	–	49.35
11	39167176	$18.8^{+8.9}_{-2.9}$	20.57787, 740.80317	37	$26.0^{+20.0}_{-5.0}$	$0.947^{+0.081}_{-0.044}$	0	81.59	40.79
12	42428568	$24.0^{+11.0}_{-5.0}$	24.14954, 748.63351	32	$31.0^{+22.0}_{-6.0}$	$0.981^{+0.066}_{-0.038}$	0	16.42	56.65
13	52195587	9.28 ± 0.52	34.42428, 757.33272	18	50.0 ± 17.0	0.29 ± 0.22	0	–	35.19
14	66439839	$9.1^{+2.8}_{-1.0}$	37.53440, 750.68798	17	$38.0^{+20.0}_{-2.0}$	$0.87^{+0.09}_{-0.29}$	0	30.94	19.55
15	67599025	6.86 ± 0.57	24.49349, 734.80456	31	$27.2^{+7.8}_{-2.7}$	$0.62^{+0.13}_{-0.35}$	0	9.86	29.54
16	70561926	17.51 ± 0.82	20.33234, 752.29687	38	$22.1^{+5.7}_{-1.8}$	0.15 ± 0.12	2	18.35	144.8
17	71028120	4.74 ± 0.31	18.60345, 744.14289	41	$21.3^{+6.3}_{-2.7}$	$0.38^{+0.22}_{-0.25}$	0	2.53	14.28
18	77437543	$10.98^{+0.66}_{-0.63}$	22.62465, 723.98986	22	$38.0^{+14.0}_{-15.0}$	0.31 ± 0.22	2	14.02	83.13
19	81089255	12.19 ± 0.69	34.10282, 750.26191	23	$37.5^{+9.4}_{-4.9}$	$0.76^{+0.07}_{-0.26}$	0	24.24	25.14
20	100776118	16.19 ± 0.74	49.79069, 746.86040	16	$62.0^{+31.0}_{-12.0}$	0.842 ± 0.01	3	30.51	177.47
21	101824521	12.17 ± 0.72	24.52324, 711.17397	30	31.0 ± 10.0	$0.18^{+0.18}_{-0.13}$	0	–	34.34
22	107113345	$9.2^{+5.1}_{-2.3}$	20.02618, 740.96843	38	$22.5^{+6.0}_{-2.4}$	0.976 ± 0.048	0	19.1	12.5
23	118339710	$10.64^{+0.65}_{-0.61}$	21.29118, 723.89974	26	33.0 ± 12.0	0.48 ± 0.29	0	4.13	62.67
24	123763494	$19.8^{+7.1}_{-4.5}$	19.53618, 722.83868	38	$21.9^{+5.9}_{-2.4}$	$0.97^{+0.076}_{-0.062}$	0	15.94	37.36
25	124029677	11.35 ± 0.59	18.76624, 731.88319	40	$20.9^{+5.2}_{-2.1}$	$0.43^{+0.22}_{-0.28}$	0	11.35	46.16
26	140215502	$5.78^{+0.38}_{-0.34}$	34.94727, 733.89270	22	$43.0^{+18.0}_{-8.0}$	$0.31^{+0.3}_{-0.21}$	0	9.83	32.98
27	140750416	8.55 ± 0.54	16.83165, 740.59260	45	$19.0^{+4.9}_{-2.2}$	0.36 ± 0.22	0	11.19	30.13
28	142278054	$11.49^{+0.96}_{-0.91}$	24.66649, 715.32865	30	$27.5^{+6.6}_{-2.8}$	$0.79^{+0.11}_{-0.38}$	0	7.72	23.25
29	145006304	$5.33^{+0.47}_{-0.43}$	16.78954, 738.74000	45	$18.9^{+5.7}_{-2.2}$	$0.62^{+0.17}_{-0.35}$	0	6.03	13.84
30	145913596	$6.9^{+2.7}_{-1.1}$	54.09067, 703.17978	14	$59.0^{+29.0}_{-8.0}$	$0.91^{+0.07}_{-0.32}$	0	5.83	25.32

Table A3: Properties of our duotransit candidates. N_{alias} is the alias number corresponding to the most probable orbital period P_{marg} .

ID	TIC ID	R_p (R_{\oplus})	$P_{\text{min}}, P_{\text{max}}$ (days)	N_{alias}	P_{marg} (days)	b	<i>Gaia</i> NSS	<i>Gaia</i> RV amp (km s $^{-1}$)	Total SNR
31	153838604	17.1 $^{+4.7}_{-3.3}$	38.37032, 729.03626	20	41.0 $^{+12.0}_{-4.0}$	0.961 $^{+0.068}_{-0.062}$	0	–	42.87
32	156716001	5.97 \pm 0.3	33.59600, 739.11201	23	43.0 $^{+30.0}_{-10.0}$	0.28 \pm 0.22	0	4.73	33.42
33	157119927	13.4 $^{+5.9}_{-3.1}$	21.19346, 699.38439	34	24.1 $^{+6.3}_{-2.9}$	0.983 $^{+0.071}_{-0.048}$	2	64.82	25.01
34	157698565	11.66 \pm 0.59	35.21588, 739.53387	22	46.0 $^{+28.0}_{-11.0}$	0.827 $^{+0.025}_{-0.041}$	0	3.49	105.68
35	159490807	15.4 $^{+4.7}_{-1.6}$	22.71853, 749.71245	34	29.0 \pm 8.0	0.938 $^{+0.048}_{-0.025}$	0	6.08	45.05
36	161169240	12.19 $^{+0.77}_{-0.73}$	26.03638, 755.05809	30	31.0 $^{+13.0}_{-4.0}$	0.47 $^{+0.21}_{-0.29}$	0	32.37	52.27
37	176518126	8.3 \pm 1.1	21.48477, 730.48210	35	24.3 $^{+7.4}_{-2.9}$	0.3 \pm 0.22	0	9.38	32.89
38	188620407	9.58 $^{+0.57}_{-0.54}$	24.13819, 1086.21870	35	28.0 \pm 10.0	0.39 \pm 0.24	0	11.56	33.51
39	193318850	7.08 \pm 0.47	15.07635, 738.73979	50	17.6 $^{+5.5}_{-2.2}$	0.31 \pm 0.22	0	–	17.05
40	207783865	15.31 \pm 0.76	21.29077, 745.17689	36	23.3 $^{+6.5}_{-2.0}$	0.802 $^{+0.023}_{-0.035}$	2	13.91	95.83
41	211409161	13.1 $^{+4.5}_{-2.6}$	18.85295, 735.26527	40	21.0 $^{+5.3}_{-2.2}$	0.943 $^{+0.077}_{-0.066}$	0	38.64	38.49
42	215402824	16.0 $^{+10.0}_{-3.0}$	41.96752, 755.41550	16	42.0 $^{+21.0}_{-2.0}$	0.93 $^{+0.11}_{-0.22}$	0	–	12.73
43	218977148	11.9 \pm 0.82	27.20845, 734.62703	28	30.6 $^{+8.1}_{-3.4}$	0.4 $^{+0.22}_{-0.26}$	0	–	22.83
44	220622886	28.0 $^{+16.0}_{-8.0}$	42.65054, 725.05958	18	29.0 $^{+23.0}_{-0.0}$	0.994 $^{+0.071}_{-0.048}$	0	65.91	18.89
45	221915858	21.3 $^{+3.7}_{-1.4}$	17.78520, 729.19319	42	19.7 $^{+5.4}_{-1.9}$	0.911 $^{+0.044}_{-0.025}$	0	60.96	33.79
46	224279805	10.55 \pm 0.64	17.51161, 735.48631	43	19.9 $^{+5.5}_{-1.9}$	0.24 \pm 0.18	0	17.77	30.43
47	235058563	11.47 \pm 0.61	88.87019, 799.83193	10	89.0 $^{+25.0}_{-9.0}$	0.38 \pm 0.23	0	–	27.17
48	237605045	44.0 $^{+17.0}_{-11.0}$	21.85618, 721.25400	34	24.0 $^{+6.0}_{-2.2}$	1.01 $^{+0.067}_{-0.051}$	0	36.88	68.84
49	242241304	9.7 \pm 1.0	23.30012, 722.30308	32	28.0 $^{+10.0}_{-4.0}$	0.855 $^{+0.061}_{-0.071}$	0	7.03	28.72
50	251057075	3.88 \pm 0.3	16.53870, 727.70489	45	20.2 $^{+7.8}_{-3.3}$	0.27 \pm 0.22	0	7.82	15.54
51	256912435	25.5 $^{+4.6}_{-3.8}$	24.66455, 715.27204	30	27.5 $^{+6.6}_{-2.8}$	0.95 \pm 0.051	0	46.85	57.7
52	265465927	11.73 \pm 0.63	23.86539, 715.96164	31	27.5 $^{+8.3}_{-3.7}$	0.27 \pm 0.19	0	19.67	38.93
53	265466589	8.57 \pm 0.59	28.55588, 685.34120	25	31.2 $^{+9.2}_{-3.7}$	0.49 $^{+0.21}_{-0.33}$	0	–	13.71
54	268534931	6.12 \pm 0.39	24.69345, 740.80465	31	32.0 $^{+25.0}_{-7.0}$	0.29 \pm 0.23	0	25.23	29.74
55	269333648	11.72 \pm 0.67	64.59377, 710.52997	12	65.0 $^{+24.0}_{-5.0}$	0.64 $^{+0.1}_{-0.25}$	0	3.2	54.46
56	275878706	8.1 $^{+3.1}_{-1.3}$	22.52196, 720.70247	33	27.0 $^{+11.0}_{-4.0}$	0.88 $^{+0.09}_{-0.21}$	1	21.81	20.68
57	279727635	10.5 $^{+4.0}_{-1.2}$	65.11699, 781.40254	13	71.0 $^{+41.0}_{-19.0}$	0.905 \pm 0.078	0	–	28.36
58	287204963	11.3 \pm 2.2	17.79021, 729.39798	42	20.3 $^{+5.8}_{-2.5}$	0.41 $^{+0.25}_{-0.28}$	0	–	17.86
59	289840544	14.0 $^{+9.1}_{-3.6}$	22.78782, 751.99813	34	25.1 $^{+6.3}_{-2.3}$	0.984 $^{+0.063}_{-0.045}$	0	6.43	17.85
60	290165539	23.4 $^{+6.3}_{-3.4}$	21.62396, 735.21479	35	27.0 $^{+19.0}_{-5.0}$	0.95 $^{+0.066}_{-0.046}$	0	45.38	92.47
61	292719109	11.07 \pm 0.59	21.89747, 722.61775	34	24.1 $^{+7.3}_{-2.2}$	0.34 \pm 0.23	0	2.5	45.22

Table A3: Properties of our duotransit candidates. N_{alias} is the alias number corresponding to the most probable orbital period P_{marg} .

ID	TIC ID	R_p (R_{\oplus})	$P_{\text{min}}, P_{\text{max}}$ (days)	N_{alias}	P_{marg} (days)	b	<i>Gaia</i> NSS	<i>Gaia</i> RV amp (km s^{-1})	Total SNR
62	294097549	4.79 ± 0.33	199.535169, 598.60600	3	$199.5^{+99.8}_{-0}$	$0.6^{+0.17}_{-0.37}$	0	4.14	22.63
63	296737508	5.88 ± 0.36	20.97145, 733.99174	36	$23.7^{+6.9}_{-2.7}$	$0.6^{+0.12}_{-0.26}$	0	1.12	41.08
64	300394149	6.46 ± 0.34	19.69843, 748.53947	39	$26.0^{+16.0}_{-6.0}$	0.27 ± 0.22	0	4.92	36.29
65	303317324	2.67 ± 0.26	16.79925, 739.16710	45	$18.5^{+4.6}_{-1.7}$	0.12 ± 0.11	0	1.16	22.71
66	304339227	2.65 ± 0.17	27.48783, 714.68370	27	$31.1^{+8.6}_{-3.6}$	0.34 ± 0.24	0	2.19	13.81
67	306249066	11.09 ± 0.61	23.63027, 732.53786	32	$27.1^{+7.8}_{-3.5}$	0.867 ± 0.021	0	7.34	49.32
68	313671132	$7.78^{+0.67}_{-0.62}$	23.65724, 733.37241	32	$28.0^{+13.0}_{-5.0}$	$0.81^{+0.08}_{-0.22}$	0	7.02	27.41
69	317923092	$12.5^{+4.8}_{-1.4}$	19.19983, 729.59400	39	$27.0^{+19.0}_{-5.0}$	$0.928^{+0.062}_{-0.048}$	0	35.1	30.15
70	323295479	11.28 ± 0.6	67.50129, 742.51554	12	$83.0^{+41.0}_{-15.0}$	0.85 ± 0.013	0	2.43	138.65
71	332697924	4.69 ± 0.33	22.88951, 732.47218	33	$26.2^{+7.1}_{-3.3}$	0.34 ± 0.23	0	5.6	11.96
72	333736132	$10.25^{+0.87}_{-0.82}$	23.61270, 731.99390	32	$26.1^{+7.1}_{-2.5}$	$0.751^{+0.045}_{-0.091}$	0	13.92	55.96
73	339399841	$13.0^{+1.7}_{-1.1}$	22.44020, 381.48503	18	$25.4^{+9.2}_{-4.2}$	$0.73^{+0.15}_{-0.34}$	0	–	21.13
74	349091983	$21.3^{+8.8}_{-4.8}$	549.0649, 549.06501	1	549.0649	$0.975^{+0.075}_{-0.058}$	0	–	44.11
75	381553868	12.41 ± 0.84	20.91681, 690.25274	33	$23.8^{+6.2}_{-2.9}$	$0.39^{+0.22}_{-0.26}$	0	9.94	22
76	393229954	$9.3^{+4.1}_{-1.2}$	75.77521, 757.75214	11	$45.0^{+31.0}_{-0.0}$	$0.907^{+0.082}_{-0.098}$	0	12.97	33.63
77	394340183	10.87 ± 0.61	60.35608, 724.27290	10	$60.0^{+30.0}_{-18.0}$	0.29 ± 0.21	0	6.77	85.43
78	396720998	$1.45^{+0.96}_{-0.85}$	70.64108, 777.05180	12	$78.0^{+19.0}_{-13.0}$	0.46 ± 0.32	0	–	6.44
79	404518509	2.72 ± 0.17	23.45386, 703.61679	18	$23.0^{+12.0}_{-1.0}$	$0.35^{+0.28}_{-0.24}$	0	0.93	22.52
80	412386707	12.37 ± 0.81	23.33181, 746.61701	33	$25.7^{+6.7}_{-2.4}$	$0.65^{+0.15}_{-0.38}$	0	6.18	28.97
81	437293313	$8.57^{+0.56}_{-0.52}$	20.26060, 749.64177	38	27.0 ± 9.0	0.25 ± 0.2	0	–	26.34
82	439491923	13.9 ± 1.3	26.34748, 711.37923	28	$31.0^{+14.0}_{-5.0}$	$0.7^{+0.14}_{-0.3}$	0	–	36.2
83	442893494	$11.55^{+0.67}_{-0.63}$	21.87047, 721.72570	22	36.0 ± 12.0	$0.34^{+0.25}_{-0.23}$	0	–	24.76
84	457649900	$18.0^{+9.8}_{-3.9}$	23.40078, 748.82538	33	$30.0^{+20.0}_{-6.0}$	$0.986^{+0.058}_{-0.033}$	0	13.11	29.26
85	466206508	13.55 ± 0.82	25.04186, 375.62557	16	$31.0^{+16.0}_{-6.0}$	0.37 ± 0.2	0	12.81	64.68

Table A4. A sample of events from `monofind` that were not planetary in nature. This table is available in its entirety online.

TIC ID	Designation	T_{c1}		T_{c2}	
		(TBJD, [Sector])	(TBJD, [Sector])	(TBJD, [Sector])	(TBJD, [Sector])
82286	Asteroid	1606.56693 [11]	2337.34989 [38]		
109979	Asteroid	1601.53746 [11]	2355.66990 [38]		
2490309	Asteroid	1577.58862 [10]	2309.44366 [37]		
3835932	Asteroid	1392.54454 [03]	2133.24553 [30]		
5740937	Asteroid	1378.87982 [02]	2097.43075 [29]		
...




Figure A1. Zoomed-in transit events for duotransit candidates TIC-304339227 to TIC-156716001 ordered by transit depth. The left panel shows the Year 1 transit, and the right panel shows the Year 3 transit. The data in blue is the *TESS* PDCSAP photometry (see Section 2). In cases with < 15 minute cadence blue points represent 30 minute bins with raw photometry shown in grey. The dashed purple lines show the best fit transit model with 1 and 2 standard deviations as shaded purple regions.




Figure A2. Zoomed-in transit events for duotransit candidates TIC-313671132 to TIC-381553868 ordered by transit depth. The left panel shows the Year 1 transit, and the right panel shows the Year 3 transit. The data in blue is the *TESS* PDCSAP photometry (see Section 2). In cases with < 15 minute cadence blue points represent 30 minute bins with raw photometry shown in grey. The dashed purple lines show the best fit transit model with 1 and 2 standard deviations as shaded purple regions.




Figure A3. Zoomed-in transit events for duotransit candidates TIC-235058563 to TIC-306249066. The left panel shows the Year 1 transit, and the right panel shows the Year 3 transit. The data in blue is the *TESS* PDCSAP photometry (see Section 2). The dashed purple line is the best fit transit model with 1 and 2 standard deviations either side.




Figure A4. Zoomed-in transit events for duotransit candidates TIC-396720998 to TIC-439491923. The left panel shows the Year 1 transit, and the right panel shows the Year 3 transit. The data in blue is the *TESS* PDCSAP photometry (see Section 2). The dashed purple line is the best fit transit model with 1 and 2 standard deviations either side.




Figure A5. Zoomed-in transit events for duotransit candidates TIC-287204963 to TIC-256912435. The left panel shows the Year 1 transit, and the right panel shows the Year 3 transit. The data in blue is the *TESS* PDCSAP photometry (see Section 2). The dashed purple line is the best fit transit model with 1 and 2 standard deviations either side.

APPENDIX B: AUTHOR AFFILIATIONS

- ¹ Department of Physics, University of Warwick, Gibbet Hill Road, Coventry CV4 7AL, UK
- ² Centre for Exoplanets and Habitability, University of Warwick, Gibbet Hill Road, Coventry CV4 7AL, UK
- ³ Physikalisches Institut, University of Bern, Gesellschaftstrasse 6, 3012 Bern, Switzerland
- ⁴ Astrophysics Research Centre, School of Mathematics and Physics, Queen's University Belfast, Belfast BT7 1NN, UK
- ⁵ Observatoire de Genève, Université de Genève, 51 Chemin Pegasi, 1290 Versoix, Switzerland
- ⁶ Centre for Exoplanet Research, School of Physics and Astronomy, University of Leicester, University Road, Leicester, LE7 7RH, UK
- ⁷ Institute of Planetary Research, German Aerospace Center, Rutherfordstrasse 2, 12489 Berlin, Germany
- ⁸ European Space Agency (ESA), European Space Research and Technology Centre (ESTEC), Keplerlaan 1, 2201 AZ Noordwijk, The Netherlands
- ⁹ Instituto de Estudios Astrofísicos, Facultad de Ingeniería y Ciencias, Universidad Diego Portales, Av. Ejército 441, Santiago, Chile
- ¹⁰ Centro de Astrofísica y Tecnologías Afines (CATA), Casilla 36-D, Santiago, Chile
- ¹¹ Instituto de Astronomía, Universidad Católica del Norte, Angamos 0610, 1270709, Antofagasta, Chile
- ¹² Armagh Observatory and Planetarium, College Hill, Armagh, BT61 9DG, N Ireland, UK

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.