




UNIVERSITY OF  
OXFORD

# Neutrino Interactions with Transverse Kinematic Imbalance — An *incomplete* Review of MINERvA Results and TKI-Community Efforts

Xianguo LU / 卢显国  
University of Oxford

Fermilab Neutrino Seminar  
1 April 2021



## Neutrino for CP-violation

- ❑ Need to detect accelerator neutrinos at  $O(1)$  GeV


GeV-neutrinos also relevant for

- ❑ Mass hierarchy measurement via atmospheric neutrino oscillations
- ❑ Background to rare event searches



## Outline

- ❑ *Neutrino interactions* with MINERvA
- ❑ Transverse Kinematic Imbalance (TKI)
  - ❖ Ideas and measurements
  - ❖ Further ideas, including
    - Hydrogen-rich high-pressure TPC

T2K / Hyper-K




Super-Kamiokande  
(ICRR, Univ. Tokyo)



Near Detectors to measure  $\nu$  interactions



NOvA






# MINERvA@FNAL


A dedicated  $\nu$ -interaction experiment




- ❑ 5.4 ton active scintillator fiducial volume
- ❑ 10- $\mu$ s beam spill,  $\sim 1$  event in tracker per spill




- ❑ Scintillator bar (CH)
- ❑ 3.3 cm base, 1.7 cm height
- ❑ 3 ns timing resolution




- ❑ 3 views
- ❑ 2.7 mm position resolution per plane



- ❑ Non-magnetized



- ❑ Muon momentum resolution (range + curvature) 8% @ 6 GeV/c
- ❑ Proton threshold 100 MeV K.E., momentum (by range) resolution 2% @ 1 GeV/c
- ❑  $\pi^0$  momentum resolution  $\sim 20\%$
- ❑ High-energy charged  $\pi$  energy resolution by calorimetry  $18\% + 8\% / \sqrt{E_\pi/\text{GeV}}$



### 2013-2019: Medium Energy (NOvA-era) data-taking



- ❖ *Constraint of the MINERvA medium energy neutrino flux using neutrino-electron elastic scattering, Phys.Rev. D100, 092001 (2019)*
- ❖ *High-Statistics Measurement of Neutrino Quasielasticlike Scattering at 6 GeV on a Hydrocarbon Target, Phys.Rev.Lett. 124, 121801 (2020)*



ME: gigantic data sets!

LE: Low Energy, peak at 3 GeV







ME: Medium Energy, peak at 6 GeV







# Neutrino Interactions




# Neutrino-Electron Elastic Scattering

LE: Phys.Rev. D93, 112007 (2016)  
ME: Phys. Rev. D 100, 092001 (2019)



Well-understood SM process

$$\nu e \rightarrow \nu e$$



LE: 135 events  
ME: 810 events

- ❑ Beam flux prediction:
  - GEANT4+hadron production data
- ❑ *in situ* flux constrained by  $\nu e$  scattering
  - ❖ reduced by  $\sim 10\%$
  - ❖ uncertainty near the peak reduced from 8% to 4%




# Inverse Muon Decay

(Muon decay  $\mu^- \rightarrow \nu_\mu + e^- + \bar{\nu}_e$ )

Another well-understood SM process

$$\nu_\mu + e^- \rightarrow \mu^- + \nu_e$$


Inverse muon decay




New flux constraint method—stay tuned!

# Charged-Current Coherent $\pi$ Production


LE: Phys.Rev.Lett. 113, 261802 (2014)  
 Phys.Rev. D97, 032014 (2018)



Pion decay





“Inverse pion decay”  
 $\times$  forbidden  
 $\checkmark$  IF  $\nu_\mu$  is replaced by a heavy neutrino (heavy neutral lepton)



Coherent production  
 à la QED pair prod.  
 $\checkmark$  allowed

$\square$  Intrinsic background to HNL search  
 [T2K, Phys.Rev. D100, 052006 (2019)]




# Kaon Production



LE: Phys.Rev. D94, 012002 (2016), Phys.Rev.Lett. 117, 061802 (2016)  
 Phys.Rev.Lett. 119, 011802 (2017)




“Inverse kaon decay”  
 — Coherent  $K^+$



- ☐  $K^+$  decay-at-rest signature  
 12.4 ns lifetime, kink, energy deposit
- ☐ LE: 6 events, predicted BG 1.77,  $3.0\sigma$






## Protons inside a nucleus

- ❑ Bound nucleons are moving—Fermi motion
- ❑ Interactions while exiting, very often breaking up the nucleus—final state interactions (FSI)


## Bound-proton decay

- ❑  $K^+$  20-200 MeV K.E. (not considering FSI) [JUNO, J.Phys.G 43, 030401 (2016)]
- ✗ background from  $K^+$  production by atmospheric neutrinos
- MINERvA can constrain it!


- ❑ Proton *decay at rest*  $\rightarrow K^+$  105 MeV K.E.
- ✓ Nice kinematic signature with decay chain coincidence
- Or not?



All these art work  
© Cheryl Patrick




*e*-A scattering  
Subedi *et al.*, Science 320, 1476 (2008)




From electron-nucleus scattering

- ❑ Fermi motion
- ❑ FSI breaking up nucleus
- ❑ *2p2h* excitation



All exist in neutrino-nucleus scattering




### Charged-current quasielastic (CCQE)



FSI can *also* modify final-state topology




Difficulty in cross section measurements already at *definition* level!







How well can we measure neutrino energy?  
 (reminder: oscillation very sensitive to baseline and energy,  $L/E$ )





2-body reaction  
 —precise

Fermi motion, FSI  
 —spread



2p2h  
 —large (not well known) fraction of large bias and spread






©Shigeo Fukuda

Let's think about something simpler...


What do you see in this sculpture?

# Transverse Kinematic Imbalance (TKI)

– Precisely identify intranuclear dynamics, or the absence thereof, in interactions between nuclei and GeV-neutrinos from accelerators



Stationary free nucleon target



Nuclear target ( $A > 1$ )

- Fermi motion
- Removal energy
- FSI
- 2p2h

Our collider neighbors have been using something similar since a long time ago

## Missing energy

From Wikipedia, the free encyclopedia



[...]

[neutrinos](#).<sup>[1]</sup> In general, missing energy is used to infer the presence of non-detectable particles and is expected to be a signature of many theories of [physics beyond the Standard Model](#).<sup>[2][3][4]</sup>

[...]

[hadron colliders](#).<sup>[5]</sup> The initial momentum of the colliding [partons](#) along the beam axis is not known —


## TKI

Multi-dimensional observation


- Momentum (magnitude)
- Angle
- Asymmetry

# Transverse Kinematic Imbalance (TKI)

## — Community Effort So Far



# Transverse Boosting Angle $\delta\alpha_T$ [XL *et al.*, Phys.Rev.C 94, 015503 (2016)]



if Fermi motion only



$\delta\vec{p}_T = \vec{p}_T^N$   
 $\delta\alpha_T$  is Fermi motion  
 direction  $\rightarrow$  isotropic


- $\delta\vec{p}_T$
- total transverse momentum
- transverse momentum imbalance
- missing pT
- ...

In full

$$\delta\vec{p}_T = \vec{p}_T^N - \Delta\vec{p}_T$$

— FSI and missing particles

# Transverse Boosting Angle $\delta\alpha_T$ [XL *et al.*, Phys.Rev.C 94, 015503 (2016)]



FSI and momentum sharing with extra particles

- Nucleus break-up
- pion absorption
- 2p2h

# Emulated Nucleon Momentum $p_N$ [Furmanski & Sobczyk, Phys.Rev.C 95, 065501 (2017)]

A more general analysis of kinematic imbalance

Transverse:  $0 = \vec{p}_T^{\ell'} + \vec{p}_T^{N'} - \delta\vec{p}_T$

Longitudinal:  $E_\nu = p_L^{\ell'} + p_L^{N'} - \delta p_L$

New variable:  $p_n \equiv \sqrt{\delta p_T^2 + \delta p_L^2}$

Neutrino energy is unknown (in the first place), equations are not closed.

Assuming exclusive  $\mu$ -p-A' final states

Use energy conservation to close the equations

$$E_\nu + m_A = E_{\ell'} + E_{N'} + E_{A'}$$

$$E_{A'} = \sqrt{m_{A'}^2 + p_n^2}$$


$p_n$ : recoil momentum of the nuclear remnant

final-state

## Dual Interpretation






For CCQE,  $A' = {}^{11}\text{C}^*$   
 No more unknowns  
 $p_n$ : neutron Fermi motion

initial-state



# TKI CCQE-like

LE: Phys.Rev.Lett. 121, 022504 (2018)





Abnormal acceleration

GENIE FSI (v2.8 hA)

*—not dark energy*




Removed in later versions








Partly expected, partly very surprising consistency!

Now compare data to models separately





[XL & Sobczyk, Phys.Rev.C 99, 055504 (2019)]





Initial-state models:

- Relativistic Fermi gas (RFG)**—simple Fermi gas model
- Local Fermi gas (LFG)**—Fermi motion sampling depends on nucleon location (local density)
- Spectral function (SF) and effective spectral function (ESF)**—Fermi motion and removal energy sampling, short range correlation (SRC) leading to momentum exceeding Fermi surface
  - ❖ Decent agreement for  $\nu n \rightarrow \mu p$ , but *not* for  $\nu n \rightarrow \mu p \pi$



GiBUU offers very different predictions



☐ Local Fermi gas embedded in nuclear potential

❖ “More same” implementation of models in different channels

❖ Fermi motion peak location better agree with  $\pi^0$  data

➤ But height *below* data

NB: 2p2h only in QE-like production, not overt pion production  
 FSI and 2p2h start to decouple




T2K measurements:

□ CC  $0\pi$


- ❖ Same-ish signal definition as MINERvA
- ❖ Same observation: prefer SF over LFG

□ CC single  $\pi^+$ :


- ❖ Complementary channel: have additional hydrogen contribution
- ❖ ESF seems OK




# TKI—Final-state effects and 2p2h





Consistent pure QE cross section (no boosting FSI in Nature)



- Smaller pion production and absorption at T2K energy
- Sensitive to 2p2h








DUNE Ar gas TPC near detector, based on GiBUU QE-like predictions

- Small  $\delta\alpha_T$ : very similar between C (fake target) and Ar
  - ✓ Predictable baseline constrained by C data from other experiments
  - ✓ Powerful calibration for new target material
- Large  $\delta\alpha_T$ : target-dependent FSI (including pion absorption) and 2p2h

# Further Ideas





Pion prefers one side: L-R asymmetry

- Resonant and nonresonant interference
- Predicted by Adler and more recent models
- Difficult measurement\* by ANL, MINERvA, T2K, consistent with 0 within error

[Adler, Ann. Phys. (N.Y.) 50, 189 (1968),  
 Sobczyk *et al.*, Phys. Rev. D 98, 073001 (2018),  
 Kabirnezhad, Phys. Rev. D 97, 013002 (2018),  
 Niewczas *et al.*, Phys. Rev. D 103, 053003 (2021)]

\*Caveats!




FIG. 14. Definition of the azimuthal  $\phi$  and polar  $\theta$  angles in the Adler system.  $\vec{k}_1$  and  $\vec{k}_2$  are vectors along the  $\nu$  and  $\mu^-$  directions, respectively, in the  $N\pi$  rest system.






FIG. 16. Distribution of events in the pion azimuthal angle  $\phi$  for the final state  $\mu^- p \pi^+$ , with  $M(p\pi^+) < 1.4$  GeV. The curve is the area-normalized prediction of the Adler model.

ANL data and Adler model [Radecky *et al.*, Phys. Rev. D 25, 1161 (1982)]





# Bird's-eye view

\*Muon pointing up




# Bird's-eye view

\*Muon pointing up




LR asymmetry of (leading) proton  
in QE-like/ $0\pi$

[Cai, XL, Ruterbories, Phys.Rev.D 100, 073010 (2019)]




LE: Phys.Rev.D  
101, 092001 (2020)



$$A_{Tx}(|\delta p_{Tx}|) = \frac{\sigma_+ - \sigma_-}{\sigma_+ + \sigma_-}$$

# Bird's-eye view

\*Muon pointing up



Have searched in MINERvA:

- ❖ Asymmetry is dynamic
- ❖ Asymmetry-flip expected as pion kicking out energetic proton

Assuming LR-asymmetry comes from RES


- ❑ Overall will be smeared or diluted, but it will tell us
- ✓ pion is indeed absorbed
  - Or, how do we actually know pions are absorbed in *ν interactions*?
- ✓ how much 2p2h there is
- ❑ Can be measured more precisely in new detectors?

[Cai, XL, Ruterbories, Phys.Rev.D 100, 073010 (2019)]

$$\nu A \rightarrow \mu p A'$$



LR asymmetry of (leading) proton ?!

# Further Ideas



Consider only charged particle productions  
from  $\nu$ - and  $\bar{\nu}$ -H interactions

➤ Leading channel has 3 final-state particles:  $\mu, \pi, p$




Double-transverse momentum imbalance  $\delta p_{TT}$

- H: 0
- A: *irreducible* broadening  $O(200 \text{ MeV})$  by Fermi motion etc.
- AH compound:  
 $\nu$ - and  $\bar{\nu}$ -H can be extracted
- Given good enough tracking, can work for any targets  
 Examples: plastic scintillator (CH or CH<sub>2</sub>)

[XL *et al.*, Phys.Rev.D 92, 051302 (2015)]

## Hydrogen-rich high-pressure TPC

- ❑ Why gas TPC? Why high pressure?
  - ❖ Acceptance, tracking threshold
  - ❖ Target mass








Raaf, TPC Mini Workshop

<https://indico.cern.ch/event/827540/contributions/3487180/>

## Hydrogen-rich high-pressure TPC

- ❑ Why gas TPC? Why high pressure?
  - ❖ Acceptance, tracking threshold
  - ❖ Target mass
- ❑ Why not pure hydrogen TPC
  - ❖ Bubble chamber: worse tracking
  - ❖ H<sub>2</sub> gas: not hydrogen-rich enough
- ❑ How rich is rich enough?
  - ❖ Element carrying as much hydrogen as possible: Carbon base C<sub>x</sub>H<sub>y</sub>







					2 He
5 B	6 C	7 N	8 O	9 F	10 Ne
13 Al	14 Si	15 P	16 S	17 Cl	18 Ar

[Hamacher-Baumann, XL, Martín-Albo, Phys.Rev.D 102, 033005 (2020)]





Jmol: an open-source Java viewer for chemical structures in 3D.  
<http://www.jmol.org/>

## Hydrogen-rich high-pressure TPC

- ❑ Why gas TPC? Why high pressure?
  - ❖ Acceptance, tracking threshold
  - ❖ Target mass
- ❑ Why not pure hydrogen TPC
  - ❖ Bubble chamber: worse tracking
  - ❖ H<sub>2</sub> gas: not hydrogen-rich enough
- ❑ How rich is rich enough?
  - ❖ Element carrying as much hydrogen as possible: Carbon base C<sub>x</sub>H<sub>y</sub>
    - Saturated, acyclic: Alkane
    - C<sub>n</sub>H<sub>2n+2</sub>
      - ✓ CH<sub>4</sub> most efficient H-carrier, but not the largest one



					2 He
5 B	6 C	7 N	8 O	9 F	10 Ne
13 Al	14 Si	15 P	16 S	17 Cl	18 Ar



[Hamacher-Baumann, XL, Martín-Albo, Phys.Rev.D 102, 033005 (2020)]

Jmol: an open-source Java viewer for chemical structures in 3D.  
<http://www.jmol.org/>

## Hydrogen-rich high-pressure TPC

❑ Why gas TPC? Why high pressure?

- ❖ Acceptance, tracking threshold
- ❖ Target mass

❑ Why not pure hydrogen TPC

- ❖ Bubble chamber: worse tracking
- ❖ H<sub>2</sub> gas: not hydrogen-rich enough

❑ How rich is rich enough?

- ❖ Element carrying as much hydrogen as possible: Carbon base C<sub>x</sub>H<sub>y</sub>

➤ Saturated, acyclic: Alkane








- ✓ CH<sub>4</sub> most efficient H-carrier, but not the largest one

- ❖ Maximal partial pressure limited by vapor pressure




➤ Theoretically hydrogen-richest mix at 10 bar: C<sub>3.93</sub>H<sub>9.86</sub>

= 17% C(CH<sub>3</sub>)<sub>4</sub> (neopentane) + 35% *i*C<sub>4</sub>H<sub>10</sub> (isobutane)  
+ 24% C<sub>4</sub>H<sub>10</sub> (butane) + 24% C<sub>3</sub>H<sub>8</sub> (propane)




[Hamacher-Baumann, XL, Martín-Albo, Phys.Rev.D 102, 033005 (2020)]



					2 He
5 B	6 C	7 N	8 O	9 F	10 Ne
13 Al	14 Si	15 P	16 S	17 Cl	18 Ar





unsaturated



Jmol: an open-source Java viewer for chemical structures in 3D.  
<http://www.jmol.org/>




# Hydrogen-rich high-pressure TPC + TKI




Simulation: “event rates” as a function of “reconstructed”  $\delta p_{TT}$

- ❑ Hydrogen signal sharpens with better tracking resolution  $\Gamma$
- ❑ Background stays wide due to intrinsic nuclear effects
- ❑ More hydrogen purer selection
  - ❖ CH<sub>4</sub> 4 times better than CH in signal/background

[Hamacher-Baumann, XL, Martín-Albo, Phys.Rev.D 102, 033005 (2020)]



helium + alkane



Hydrogen-rich high-pressure TPC

Pure and more hydrogen events

- ✓ Can be purer than CH
- With large volume and powerful beam flux
  - ✓ Can go up to 4t CH-equivalent
    - ❖ H<sub>2</sub>: 1t
  - ✓ 20k events/year
- ✓ Not covered in this talk:
  - ❖ Demonstrated by simulation mixture is workable
- ✓ Further design work on-going—stay tuned!

- Hydrogen is free proton
- Bound protons in nuclei are background

Typical pion production cross section

DUNE proposed flux

[Hamacher-Baumann, XL, Martín-Albo, Phys.Rev.D 102, 033005 (2020)]

# Summary

Yesterday's signal (SM) is today's background (for BSM);  
Yesterday's background (nuclear physics) is still today's background.

– Confucius

GeV neutrinos are fun!  
CP violation! Mass ordering! Proton decay! Heavy neutral leptons!  
Yet we still need to understand the underlying nuclear physics.

像教行子孔師先



# Summary

## □ MINERvA

- ❖ 5.4 t scintillator tracker + nuclear targets + calorimeter + magnetized muon spectrometer
- ❖ LE program was completed, ME analyses in pipeline with more than 10 times statistics, reaching neutrino energy beyond 50 GeV
- ❖ A full scientific program
  - ✓ Filling in the “periodic table” of  $\nu$  interactions

## □ TKI

- ❖ New emerging activities in analysis and detector design
- ❖ Measurements from MINERvA and T2K, and actually more on-going

Sunday, April 18, 2021  
3:57PM - 4:09PM

Live



[L14.00002: Using Transverse Kinematic Imbalance to Probe Intranuclear Dynamics in Pion Scattering on Argon in ProtoDUNE](#)  
Kang Yang



## ❖ $\nu$ interactions

- ✓ Initial state probed by  $p_N$  is a challenge
  - Strong constraint for model consistency in different channels
- ✓ Pure CCQE baseline at small  $\delta\alpha_T$  powerful calibration tool
  - Safe extrapolation between different targets
- ✓ FSI and 2p2h start to decouple when combining QE-like + pion production
- Finally seeing pion absorption (produced and absorbed, without existing outside the nucleus) after having believed in it for many years? (*dynamic LR asymmetry*)
- $\nu$ - and  $\bar{\nu}$ -H scattering can be revived since 1980s? (*Hydrogen-rich high-pressure TPC*)

Thank you!




*Cueva de las Manos, Perito Moreno, Argentina. The art in the cave is dated between 13,000–9,000 BP, stenciled, mostly left hands are shown.*



# BACKUP


# TKI measurements @ MINERvA

– QE-like measurement on C probing  $\nu n \rightarrow \mu p$



2p2h-like enhancement needs to be even stronger to fill the dip









# TKI measurements @ MINERvA

– Inclusive  $\pi^0$  production on C probing  $\nu n \rightarrow \mu p \pi^0$



✗ Fermi motion peak in pion production worse modeled than in QE



✓ Large missing pT region reasonably modeled

Xiangguo Lu, Oxford

15

# TKI measurements @ MINERvA

– Inclusive  $\pi^0$  production on C probing  $\nu n \rightarrow \mu p \pi^0$



Xiangguo Lu, Oxford

16

□ Nominal: version 2.8.4

- ❖ global Fermi Gas (RFG) model with Bodek-Ritchie (BR) tail [Bodek & Ritchie, Phys. Rev. D 23, 1070 (1981)]
- ❖ hA FSI [Dytman & Meyer, AIP Conf.Proc. 1405, 213 (2011)]

□ MnvGENIE-v1: GENIE MINERvA Tune (v1)

- ❖ Added Random Phase Approximation (RPA) [Nieves *et al.*, Phys.Rev. C70, 055503 (2004)]
- ❖ Non-resonance pion production scaled down by 75% [Wilkinson *et al.*, Phys.Rev. D90, 112017 (2014)]
- ❖ Valencia 2p2h [Nieves *et al.*, Phys.Lett. B707, 72 (2012); Sobczyk, Phys. Rev. C 86, 015504 (2012); Gran *et al.*, Phys.Rev. D88, 113007 (2013); Schwehr *et al.*, arXiv:1601.02038]
  - Tuned to MINERvA inclusive data → significant enhancement in small 4-momentum transfer region [MINERvA, Phys.Rev.Lett. 116, 071802 (2016)]

# END