expression
genes
pathogen
gene
susceptibility
defence
time
data
infection
signalling
response
TGA3
plant
SA
JA
ABA
hpi
synthesis
analysis
network
interactions
differentially
overrepresented
regulation
known
Arabidopsis
responses
changes
cleared
models
role
used
mutants
processes
series
GO
GP2S
identified
up-regulated
reduced
lesion
ability
spatial
protein
activation
also
may
biosynthetic
family
important
pathogens
resolve
Although
hormone
motif
interaction
types
around
novel
cell
conditions
modelling
direct
hormones
ANAC055
period
profiles
plants
necrotrophic
PAL
components
NPR1
growth
timing
leaves
key
involved
levels
specific
TFs
down-regulated
core
et
one
leaf
highlighted
host
likely
two
function
TF
roles
high-resolution
due
whole
false
syringae
well
static
differences
biological
many
hypothesis
hence
transcriptional
clear
TGA
compared
expressed
regulatory
increased
small
rate
leads
number
clock
multiple
reflect
tissue
mediated
clusters
suggesting
example
terms
generate
al
shown
and/or
plant
mutants
processes
identified
GO
GP2S
series
TGA3
up-regulated
process
reduced
response
models
genome
generated
points
mm
lesion
ability
spatial
ET
highlight
majority
profiling
Fig
protein
activation
also
may
autophagy
test
experimentally
changes
cleared
help
biosynthetic
family
important
pathogens
resolve
several
early
suggests
single
different
cluster
downstream
negative
Although
hormone
motif
interaction
types
around
novel
cell
conditions
modelling
direct
hormones
known
Arabidopsis
ABA
JA
overrepresented
differentially
regulation
whole
false
syringae
well
static
responses
ANAC055
period
profiles
plants
necrotrophic
PAL
due
high-resolution
roles
TF
function
two
components
et
core
one
leaf
down-regulated
NPR1
growth
likely
host
highlighted
TFs
specific
timing
leaves
key
differences
synthesis
hpi
levels
involved
biological
many
hypothesis
analysis
network
interactions
hence
transcriptional
clear
pathogen
signalling
TGA
compared
gene
susceptibility
expressed
regulatory
increased
small
rate
leads
number
clock