

APTS Statistical Inference, assessment questions, 2017

To students and supervisors. These questions are formative rather than summative. It may be necessary to delete some parts of some questions for a summative assessment, or to provide additional questions.

1. This is a question about Lindley's paradox (Lindley, 1957), P -values, and the conventional 0.05 threshold.

(a) Suppose you are sitting in a bar talking to an experimental psychologist about significance levels. An informal statement of *Lindley's paradox* is that a P -value for H_0 smaller than α can correspond to a likelihood ratio $f_0(y)/f_1(y)$ greater than $1/\alpha$. Provide a proof of this statement which you can sketch on a napkin (by all means include the napkin in your answer.) Hint: see DeGroot and Schervish (2002, sec. 8.9), or work backwards from the next question.

(b) Study Figure 1, and see if you can replicate it, either showing your workings or including your code. What is this graph showing? For example, how would you feel about rejecting H_0 at a P -value of 0.0499, when $n = 20$?

Some hints:

$$\bar{Y} \sim N(\mu, \sigma^2/n) \tag{1a}$$

with σ^2 known (take $\sigma = 1$). The competing hypotheses are

$$H_0 : \mu = 0 \quad \text{versus} \quad H_1 : \mu = 1, \tag{1b}$$

i.e. a separation of σ . In this case, a sensible statistic for forming a P -value for H_0 is $t(\bar{y}) = \bar{y}$, which will be larger under H_1 than H_0 . This gives

$$p(\bar{y}; H_0) = \Pr(\bar{Y} \geq \bar{y}; H_0) = 1 - \Phi(\bar{y}; 0, 1/n)$$

where Φ is the Normal distribution function with specified expectation and variance. You can then find the value of \bar{y} which implies a P -value of 0.05 (use the `qnorm` function), and at this value compute the likelihood ratio $\phi(\bar{y}; H_0)/\phi(\bar{y}; H_1)$ where ϕ is the Normal PDF (use the `dnorm` function).

(c) Consider

$$H_0 : \mu = 0 \quad \text{versus} \quad H_1 : \mu > 0$$




Figure 1: The likelihood ratio corresponding to a P -value for H_0 of 0.05, for the model and hypotheses given in (1).

in the case where σ^2 is known and n is fixed. Produce a graph showing the value of the minimum likelihood ratio over H_1 for a range of P -values for H_0 from 0.001 to 0.1. Check your graph against the minimum shown in Figure 1. Hint: you should be able to compute this graph directly. You might find Edwards *et al.* (1963) or Goodman (1999a,b) helpful.

Comment on whether the conventional choice of 0.05 is a suitable threshold for choosing between hypotheses, or whether some other choice might be better. You may also like to reflect on the origin of the value 0.05, see Cowles and Davis (1982).

2. Here's a question about confidence sets and P -values for the Poisson distribution. Let the model be

$$Y_1, \dots, Y_n \stackrel{\text{iid}}{\sim} \text{Poisson}(\lambda), \quad \lambda > 0.$$

- (a) Consider the general case where $Y_1, \dots, Y_n \stackrel{\text{iid}}{\sim} f(\cdot; \theta)$, for $\theta \in \Omega$, where Ω is a d -dimensional convex subset of Euclidean space. Wilks's theorem states that under some regularity conditions,

$$-2 \log \frac{f(Y; \theta_0)}{f(Y; \hat{\theta}(Y))} \xrightarrow{D} \chi_d^2 \quad \text{under } \theta = \theta_0,$$

where $\hat{\theta}$ is the maximum likelihood estimator (MLE) of θ , and ' \xrightarrow{D} ' denotes convergence in distribution. Show how Wilks's theorem can be used to construct an approximately exact family of confidence procedures (i.e. a family which would be exact in the limit as $n \rightarrow \infty$). Give a simple verbal description of the confidence procedure when $d = 1$ and $\alpha = 0.05$.

- (b) Show that the log-likelihood function has the form

$$\begin{aligned} \ell(\lambda; \mathbf{y}) &= \sum_{i=1}^n \left(-\lambda + y_i \log(\lambda) - \log(y_i!) \right) \\ &= c - n\lambda + n\bar{y} \log(\lambda) \end{aligned}$$

where $\bar{y} := n^{-1}(y_1 + \dots + y_n)$, and c is some additive constant which can be ignored. Check that the Maximum Likelihood (ML) estimator is $\hat{\lambda}(\mathbf{y}) = \bar{y}$ (don't forget to check the second-order condition).

- (c) Now suppose that $\bar{y} = 0.20$ from $n = 20$ observations. Draw the log-likelihood function, and indicate the Wilks's Theorem 95% approximately exact confidence set.
- (d) Compute the P -value for the hypothesis $H_0 : \lambda = 0.1$. Hint: although you could do this by trial and error, you can also do it exactly (i.e. to within computer precision) once you know the value of the log-likelihood at $\lambda = 0.1$.

3. Here is a question on the Neyman-Pearson approach to Hypothesis Testing. Suppose there are two simple hypotheses,

$$H_0 : Y \sim f_0 \quad \text{versus} \quad H_1 : Y \sim f_1.$$

- (a) You must choose between H_0 and H_1 according to the loss function

		‘True’	
		H_0	H_1
Choose	H_0	c_{00}	c_{01}
	H_1	c_{10}	c_{11}

where $c_{00} < c_{10}$ and $c_{11} < c_{01}$. Show that a decision rule for choosing between H_0 and H_1 is admissible if and only if it has the form

$$\frac{f_0(y)}{f_1(y)} \begin{cases} < c & \text{choose } H_1 \\ = c & \text{toss a coin} \\ > c & \text{choose } H_0 \end{cases} \quad (2)$$

for some *critical value* $c > 0$. Hint: find the Bayes Rule for prior probabilities (π_0, π_1) where $\pi_1 = 1 - \pi_0$, and then apply the Complete Class Theorem (CCT).

- (b) Neyman-Pearson proposed a different approach. Let $\mathcal{R} \subset \mathcal{Y}$ be the region in which H_0 is rejected. Define the Type 1 (α) and Type 2 (β) error rates as

$$\alpha := \Pr(Y \in \mathcal{R}; H_0) \quad (\text{Type 1})$$

$$\beta := \Pr(Y \notin \mathcal{R}; H_1) \quad (\text{Type 2})$$

Provide a verbal description of each of these errors. Now show that $\alpha + c \cdot \beta$ is minimised when

$$\mathcal{R} = \{y \in \mathcal{Y} : f_0(y)/f_1(y) < c\} \quad (3)$$

for some $c > 0$. Conclude by explaining why \mathcal{R} defined in (3) minimises the β for any specified α .

- (c) Define

$$r(y) := f_0(y)/f_1(y).$$

Show that $E\{r(Y); H_1\} = 1$ and $E\{r(Y); H_0\} \geq 1$ (hint: Jensen’s inequal-

ity). Sketch, on the same axis, the PMFs of $r(Y)$ under H_0 and H_1 . Add a critical value c , and shade in the areas representing the α and β values.

- (d) The *operating characteristics curve (OCC)* of \mathcal{R} in (3) plots α (x -axis) against β (y -axis) for varying c . Sketch this curve: you will need to show that it goes through $(0, 1)$ and $(1, 0)$, and also that it is convex. Hint for the last part: consider tossing a coin to choose between two different values of c : what would α and β be for this mixed decision rule?
- (e) (Harder.) Explain why the gradient of the OCC is $\Delta\beta/\Delta\alpha \approx -1/c$. Add a few different specific values of c to your sketch: you should make sure that the aspect ratio of your sketch is approximately 1.
- (f) Now consider

$$H_0 : Y_1, \dots, Y_n \stackrel{\text{iid}}{\sim} f_0 \quad \text{versus} \quad H_1 : Y_1, \dots, Y_n \stackrel{\text{iid}}{\sim} f_1.$$

Draw a new sketch showing the two cases where n is small and n is large (i.e. two OCCs on the same sketch). On the sketch draw the line $\alpha = 0.05$ and comment on the value of β in the two cases.

- (g) Finally, explain how a sketch such as the one you have drawn in Q3f can be used to choose the sample size and the critical value for target values such as $\alpha = 0.05$ and $\beta = 0.2$. (These are common values in practice.)

Jonathan Rougier
University of Bristol
Dec 2017

References

- M.S. Bartlett, 1957. A comment on D.V. Lindley's statistical paradox. *Biometrika*, **44**, 533–534.
- M. Cowles and C. Davis, 1982. On the origins of the .05 level of statistical significance. *American Psychologist*, **37**(5), 553–558.
- M.H. DeGroot and M.J. Schervish, 2002. *Probability and Statistics*. Addison-Wesley Publishing Co., Reading MA, 3rd edition.
- W. Edwards, H. Lindman, and L.J. Savage, 1963. Bayesian statistical inference for psychological research. *Psychological Review*, **70**(3), 193–242.

- S. Goodman, 1999a. Toward evidence-based medical statistics. 1: The p -value fallacy. *Annals of Internal Medicine*, **130**, 995–1004.
- S. Goodman, 1999b. Toward evidence-based medical statistics. 2: The Bayes factor. *Annals of Internal Medicine*, **130**, 1005–1013.
- D.V. Lindley, 1957. A statistical paradox. *Biometrika*, **44**, 187–192. See also Bartlett (1957).