

# Statistical Modelling

Anthony Davison and Jon Forster

©2010

<http://stat.epfl.ch>, <http://www.s3ri.soton.ac.uk>

<b>1. Model Selection</b>	<b>2</b>
Overview . . . . .	3
<b>Basic Ideas</b>	<b>4</b>
Why model? . . . . .	5
Criteria for model selection . . . . .	6
Motivation . . . . .	7
Setting . . . . .	10
Logistic regression . . . . .	11
Nodal involvement . . . . .	12
Log likelihood . . . . .	15
Wrong model . . . . .	16
Out-of-sample prediction . . . . .	18
Information criteria . . . . .	19
Nodal involvement . . . . .	21
Theoretical aspects . . . . .	22
Properties of AIC, NIC, BIC . . . . .	23
<b>Linear Model</b>	<b>24</b>
Variable selection . . . . .	25
Stepwise methods . . . . .	26
Nuclear power station data . . . . .	27
Stepwise Methods: Comments . . . . .	29
Prediction error . . . . .	30
Example . . . . .	32
Cross-validation . . . . .	33
Other criteria . . . . .	35
Experiment . . . . .	36
<b>Sparse Variable Selection</b>	<b>40</b>
Motivation . . . . .	41
Desiderata . . . . .	42
Example: Lasso . . . . .	43
Soft thresholding . . . . .	44
Example . . . . .	46

Penalties . . . . .	47
Threshold functions . . . . .	48
Properties of penalties . . . . .	49
Oracle . . . . .	50
<b>Bayesian Inference</b>	<b>51</b>
Thomas Bayes (1702–1761) . . . . .	52
Bayesian inference . . . . .	53
Encompassing model . . . . .	55
Inference . . . . .	56
Lindley’s paradox . . . . .	57
Model averaging . . . . .	58
Cement data . . . . .	59
DIC . . . . .	63
MDL . . . . .	64
<b>Bayesian Variable Selection</b>	<b>65</b>
Variable selection . . . . .	66
Example: NMR data . . . . .	67
Wavelets . . . . .	68
Posterior . . . . .	69
Shrinkage . . . . .	70
Empirical Bayes . . . . .	71
Example: NMR data . . . . .	72
Comments . . . . .	74
<b>2. Beyond the Generalised Linear Model</b>	<b>75</b>
Overview . . . . .	76
<b>Generalised Linear Models</b>	<b>77</b>
GLM recap . . . . .	78
GLM failure . . . . .	79
<b>Overdispersion</b>	<b>80</b>
Example 1 . . . . .	81
Quasi-likelihood . . . . .	85
Reasons . . . . .	87
Direct models . . . . .	89
Random effects . . . . .	91
<b>Dependence</b>	<b>93</b>
Example 1 revisited . . . . .	94
Reasons . . . . .	95
Random effects . . . . .	96
Marginal models . . . . .	97
Clustered data . . . . .	99
Example 2: Rat growth . . . . .	100
<b>Random Effects and Mixed Models</b>	<b>104</b>
Linear mixed models . . . . .	105
Discussion . . . . .	108
LMM fitting . . . . .	110
REML . . . . .	111

Estimating random effects . . . . .	112
Bayesian LMMs . . . . .	113
Example 2 revisited . . . . .	115
GLMMs . . . . .	119
GLMM fitting . . . . .	122
Bayesian GLMMS . . . . .	126
Example 1 revisited . . . . .	127
<b>Conditional independence and graphical representations</b>	<b>129</b>
Conditional independence . . . . .	130
Graphs . . . . .	133
DAGs . . . . .	134
Undirected graphs . . . . .	140
<b>3. Missing Data and Latent Variables</b>	<b>145</b>
Overview . . . . .	146
<b>Missing Data</b>	<b>147</b>
Examples . . . . .	148
Introduction . . . . .	150
Issues . . . . .	151
Models . . . . .	152
Ignorability . . . . .	153
Inference . . . . .	154
Nonignorable models . . . . .	157
<b>Latent Variables</b>	<b>161</b>
Basic idea . . . . .	162
Galaxy data . . . . .	163
Other latent variable models . . . . .	166
<b>EM Algorithm</b>	<b>167</b>
EM algorithm . . . . .	168
Toy example . . . . .	171
Example: Mixture model . . . . .	172
Example: Galaxy data . . . . .	173
Exponential family . . . . .	175
Comments . . . . .	176

## Overview

1. Basic ideas
2. Linear model
3. Sparse variable selection
4. Bayesian inference
5. Bayesian variable selection

APTS: Statistical Modelling

April 2010 – slide 3

## Basic Ideas

### Why model?



George E. P. Box (1919–):

All models are wrong, but some models are useful.

- Some reasons we construct models:
  - to simplify reality (efficient representation);
  - to gain understanding;
  - to compare scientific, economic, ... theories;
  - to predict future events/data;
  - to control a process.
- We (statisticians!) rarely believe in our models, but regard them as temporary constructs subject to improvement.
- Often we have several and must decide which is preferable, if any.

APTS: Statistical Modelling

April 2010 – slide 5

## Criteria for model selection

- Substantive knowledge, from prior studies, theoretical arguments, dimensional or other general considerations (often qualitative)
- Sensitivity to failure of assumptions (prefer models that are robustly valid)
- Quality of fit—residuals, graphical assessment (informal), or goodness-of-fit tests (formal)
- Prior knowledge in Bayesian sense (quantitative)
- Generalisability of conclusions and/or predictions: same/similar models give good fit for many different datasets
- ... but often we have just one dataset ...

APTS: Statistical Modelling

April 2010 – slide 6

## Motivation

Even after applying these criteria (but also before!) we may compare many models:


- linear regression with  $p$  covariates, there are  $2^p$  possible combinations of covariates (each in/out), before allowing for transformations, etc.— if  $p = 20$  then we have a problem;
- choice of bandwidth  $h > 0$  in smoothing problems
- the number of different clusterings of  $n$  individuals is a Bell number (starting from  $n = 1$ ): 1, 2, 5, 15, 52, 203, 877, 4140, 21147, 115975, ...
- we may want to assess which among  $5 \times 10^5$  SNPs on the genome may influence reaction to a new drug;
- ...

For reasons of economy we seek 'simple' models.

APTS: Statistical Modelling

April 2010 – slide 7

## Albert Einstein (1879–1955)



'Everything should be made as simple as possible, **but no simpler.**'

APTS: Statistical Modelling

April 2010 – slide 8

## William of Occam (?1288–?1348)



Occam's razor: **Entia non sunt multiplicanda sine necessitate: entities should not be multiplied beyond necessity.**

APTS: Statistical Modelling

April 2010 – slide 9

## Setting

- To focus and simplify discussion we will consider parametric models, but the ideas generalise to semi-parametric and non-parametric settings
- We shall take generalised linear models (GLMs) as example of moderately complex parametric models:
  - Normal linear model has three key aspects:
    - ▷ *structure for covariates: linear predictor*  $\eta = x^T\beta$ ;
    - ▷ *response distribution:  $y \sim N(\mu, \sigma^2)$* ; and
    - ▷ *relation  $\eta = \mu$  between  $\mu = E(y)$  and  $\eta$ .*
  - GLM extends last two to
    - ▷  *$y$  has density*

$$f(y; \theta, \phi) = \exp \left\{ \frac{y\theta - b(\theta)}{\phi} + c(y; \phi) \right\},$$

where  $\theta$  depends on  $\eta$ ; *dispersion parameter*  $\phi$  is often known; and

- ▷  $\eta = g(\mu)$ , where  $g$  is monotone *link function*.

APTS: Statistical Modelling

April 2010 – slide 10

## Logistic regression

- Commonest choice of link function for binary responses:

$$\Pr(Y = 1) = \pi = \frac{\exp(x^T \beta)}{1 + \exp(x^T \beta)}, \quad \Pr(Y = 0) = \frac{1}{1 + \exp(x^T \beta)},$$

giving linear model for log odds of 'success',

$$\log \left\{ \frac{\Pr(Y = 1)}{\Pr(Y = 0)} \right\} = \log \left( \frac{\pi}{1 - \pi} \right) = x^T \beta.$$

- Log likelihood for  $\beta$  based on independent responses  $y_1, \dots, y_n$  with covariate vectors  $x_1, \dots, x_n$  is

$$\ell(\beta) = \sum_{j=1}^n y_j x_j^T \beta - \sum_{j=1}^n \log \{1 + \exp(x_j^T \beta)\}$$

- Good fit gives small deviance  $D = 2 \left\{ \ell(\tilde{\beta}) - \ell(\hat{\beta}) \right\}$ , where  $\hat{\beta}$  is model fit MLE and  $\tilde{\beta}$  is unrestricted MLE.

APTS: Statistical Modelling

April 2010 – slide 11

## Nodal involvement data

Table 1: Data on nodal involvement: 53 patients with prostate cancer have nodal involvement ( $r$ ), with five binary covariates age etc.

$m$	$r$	age	stage	grade	xray	acid
6	5	0	1	1	1	1
6	1	0	0	0	0	1
4	0	1	1	1	0	0
4	2	1	1	0	0	1
4	0	0	0	0	0	0
3	2	0	1	1	0	1
3	1	1	1	0	0	0
3	0	1	0	0	0	1
3	0	1	0	0	0	0
2	0	1	0	0	1	0
2	1	0	1	0	0	1
2	1	0	0	1	0	0
1	1	1	1	1	1	1
⋮	⋮	⋮	⋮	⋮	⋮	
1	1	0	0	1	0	1
1	0	0	0	0	1	1
1	0	0	0	0	1	0

APTS: Statistical Modelling

April 2010 – slide 12

## Nodal involvement deviances


Deviances  $D$  for 32 logistic regression models for nodal involvement data. + denotes a term included in the model.

age	st	gr	xr	ac	df	$D$	age	st	gr	xr	ac	df	$D$
					52	40.71	+	+	+			49	29.76
+					51	39.32	+	+		+		49	23.67
	+				51	33.01	+	+			+	49	25.54
		+			51	35.13	+		+	+		49	27.50
			+		51	31.39	+		+		+	49	26.70
				+	51	33.17	+			+	+	49	24.92
+	+				50	30.90		+	+	+		49	23.98
+		+			50	34.54		+	+		+	49	23.62
+			+		50	30.48		+		+	+	49	19.64
+				+	50	32.67			+	+	+	49	21.28
	+	+			50	31.00	+	+	+	+		48	23.12
	+		+		50	24.92	+	+	+		+	48	23.38
	+			+	50	26.37	+	+		+	+	48	19.22
		+	+		50	27.91	+		+	+	+	48	21.27
		+		+	50	26.72		+	+	+	+	48	18.22
			+	+	50	25.25	+	+	+	+	+	47	18.07

APTS: Statistical Modelling

April 2010 – slide 13

## Nodal involvement



### Adding terms

- always increases the log likelihood  $\hat{\ell}$  and so reduces  $D$ ,
  - increases the number of parameters,
- so taking the model with highest  $\hat{\ell}$  (lowest  $D$ ) would give the full model

### We need to trade off quality of fit (measured by $D$ ) and model complexity (number of parameters)

APTS: Statistical Modelling

April 2010 – slide 14



## Log likelihood

- Given (unknown) **true model**  $g(y)$ , and **candidate model**  $f(y; \theta)$ , Jensen's inequality implies that

$$\int \log g(y)g(y) dy \geq \int \log f(y; \theta)g(y) dy, \quad (1)$$

with equality if and only if  $f(y; \theta) \equiv g(y)$ .

- If  $\theta_g$  is the value of  $\theta$  that maximizes the expected log likelihood on the right of (1), then it is natural to choose the candidate model that maximises

$$\bar{\ell}(\hat{\theta}) = n^{-1} \sum_{j=1}^n \log f(y_j; \hat{\theta}),$$

which should be an estimate of  $\int \log f(y; \theta)g(y) dy$ . However as  $\bar{\ell}(\hat{\theta}) \geq \bar{\ell}(\theta_g)$ , by definition of  $\hat{\theta}$ , this estimate is biased upwards.

- We need to correct for the bias, but in order to do so, need to understand the properties of likelihood estimators when the assumed model  $f$  is not the true model  $g$ .

APTS: Statistical Modelling

April 2010 – slide 15

## Wrong model

Suppose the true model is  $g$ , that is,  $Y_1, \dots, Y_n \stackrel{\text{iid}}{\sim} g$ , but we assume that  $Y_1, \dots, Y_n \stackrel{\text{iid}}{\sim} f(y; \theta)$ . The log likelihood  $\ell(\theta)$  will be maximised at  $\hat{\theta}$ , and

$$\bar{\ell}(\hat{\theta}) = n^{-1} \ell(\hat{\theta}) \xrightarrow{\text{a.s.}} \int \log f(y; \theta_g)g(y) dy, \quad n \rightarrow \infty,$$

where  $\theta_g$  minimizes the Kullback–Leibler discrepancy

$$KL(f_\theta, g) = \int \log \left\{ \frac{g(y)}{f(y; \theta)} \right\} g(y) dy.$$

$\theta_g$  gives the density  $f(y; \theta_g)$  closest to  $g$  in this sense, and  $\hat{\theta}$  is determined by the finite-sample version of  $\partial KL(f_\theta, g)/\partial \theta$ , i.e.

$$0 = n^{-1} \sum_{j=1}^n \frac{\partial \log f(y_j; \hat{\theta})}{\partial \theta}.$$

APTS: Statistical Modelling

April 2010 – slide 16

## Wrong model II

**Theorem 1** Suppose the true model is  $g$ , that is,  $Y_1, \dots, Y_n \stackrel{\text{iid}}{\sim} g$ , but we assume that  $Y_1, \dots, Y_n \stackrel{\text{iid}}{\sim} f(y; \theta)$ . Then under mild regularity conditions the maximum likelihood estimator  $\hat{\theta}$  satisfies

$$\hat{\theta} \sim N_p \{ \theta_g, I(\theta_g)^{-1} K(\theta_g) I(\theta_g)^{-1} \}, \quad (2)$$

where  $f_{\theta_g}$  is the density minimising the Kullback–Leibler discrepancy between  $f_\theta$  and  $g$ ,  $I$  is the Fisher information for  $f$ , and  $K$  is the variance of the score statistic. The likelihood ratio statistic

$$W(\theta_g) = 2 \left\{ \ell(\hat{\theta}) - \ell(\theta_g) \right\} \sim \sum_{r=1}^p \lambda_r V_r,$$

where  $V_1, \dots, V_p \stackrel{\text{iid}}{\sim} \chi_1^2$ , and the  $\lambda_r$  are eigenvalues of  $K(\theta_g)^{1/2} I_g(\theta_g)^{-1} K(\theta_g)^{1/2}$ . Thus  $E\{W(\theta_g)\} = \text{tr}\{I(\theta_g)^{-1} K(\theta_g)\}$ .

Under the correct model,  $\theta_g$  is the 'true' value of  $\theta$ ,  $K(\theta) = I(\theta)$ ,  $\lambda_1 = \dots = \lambda_p = 1$ , and we recover the usual results.

APTS: Statistical Modelling

April 2010 – slide 17

## Out-of-sample prediction

- We need to fix two problems with using  $\bar{\ell}(\hat{\theta})$  to choose the best candidate model:
  - upward bias, as  $\bar{\ell}(\hat{\theta}) \geq \bar{\ell}(\theta_g)$  because  $\hat{\theta}$  is based on  $Y_1, \dots, Y_n$ ;
  - no penalisation if the dimension of  $\theta$  increases.
- If we had another independent sample  $Y_1^+, \dots, Y_n^+ \stackrel{\text{iid}}{\sim} g$  and computed

$$\bar{\ell}^+(\hat{\theta}) = n^{-1} \sum_{j=1}^n \log f(Y_j^+; \hat{\theta}),$$

then both problems disappear, suggesting that we choose the candidate model that maximises

$$E_g \left[ E_g^+ \left\{ \bar{\ell}^+(\hat{\theta}) \right\} \right],$$

where the inner expectation is over the distribution of the  $Y_j^+$ , and the outer expectation is over the distribution of  $\hat{\theta}$ .

APTS: Statistical Modelling

April 2010 – slide 18

### Information criteria

- Previous results on wrong model give

$$E_g \left[ E_g^+ \left\{ \bar{\ell}^+(\hat{\theta}) \right\} \right] \doteq \int \log f(y; \theta_g) g(y) dy - \frac{1}{2n} \text{tr} \{ I_g(\theta_g)^{-1} K(\theta_g) \},$$

where the second term is a penalty that depends on the model dimension.

- We want to estimate this based on  $Y_1, \dots, Y_n$  only, and get

$$E_g \left\{ \bar{\ell}(\hat{\theta}) \right\} \doteq \int \log f(y; \theta_g) g(y) dy + \frac{1}{2n} \text{tr} \{ I_g(\theta_g)^{-1} K(\theta_g) \},$$

- To remove the bias, we aim to maximise

$$\bar{\ell}(\hat{\theta}) - \frac{1}{n} \text{tr}(\hat{J}^{-1} \hat{K}),$$

where

$$\hat{K} = \sum_{j=1}^n \frac{\partial \log f(y_j; \hat{\theta})}{\partial \theta} \frac{\partial \log f(y_j; \hat{\theta})}{\partial \theta^T}, \quad \hat{J} = - \sum_{j=1}^n \frac{\partial^2 \log f(y_j; \hat{\theta})}{\partial \theta \partial \theta^T};$$

the latter is just the observed information matrix.

APTS: Statistical Modelling

April 2010 – slide 19

### Information criteria


- Let  $p = \dim(\theta)$  be the number of parameters for a model, and  $\hat{\ell}$  the corresponding maximised log likelihood.
- For historical reasons we choose models that **minimise** similar criteria
  - $2(p - \hat{\ell})$  (AIC—Akaike Information Criterion)
  - $2\{\text{tr}(\hat{J}^{-1} \hat{K}) - \hat{\ell}\}$  (NIC—Network Information Criterion)
  - $2(\frac{1}{2}p \log n - \hat{\ell})$  (BIC—Bayes Information Criterion)
  - AIC<sub>c</sub>, AIC<sub>u</sub>, DIC, EIC, FIC, GIC, SIC, TIC, ...
  - Mallows  $C_p = RSS/s^2 + 2p - n$  commonly used in regression problems, where  $RSS$  is residual sum of squares for candidate model, and  $s^2$  is an estimate of the error variance  $\sigma^2$ .

APTS: Statistical Modelling

April 2010 – slide 20

## Nodal involvement data

AIC and BIC for  $2^5$  models for binary logistic regression model fitted to the nodal involvement data. Both criteria pick out the same model, with the three covariates *st*, *xr*, and *ac*, which has deviance  $D = 19.64$ . Note the sharper increase of BIC after the minimum.



APTS: Statistical Modelling

April 2010 – slide 21

## Theoretical aspects

- We may suppose that the true underlying model is of infinite dimension, and that by choosing among our candidate models we hope to get as close as possible to this ideal model, using the data available.
- If so, we need some measure of distance between a candidate and the true model, and we aim to minimise this distance.
- A model selection procedure that selects the candidate closest to the truth for large  $n$  is called **asymptotically efficient**.
- An alternative is to suppose that the true model is among the candidate models.
- If so, then a model selection procedure that selects the true model with probability tending to one as  $n \rightarrow \infty$  is called **consistent**.

APTS: Statistical Modelling

April 2010 – slide 22

## Properties of AIC, NIC, BIC

- We seek to find the correct model by minimising  $IC = c(n, p) - 2\hat{\ell}$ , where the penalty  $c(n, p)$  depends on sample size  $n$  and model dimension  $p$
- Crucial aspect is behaviour of differences of IC.
- We obtain IC for the true model, and  $IC_+$  for a model with one more parameter. Then

$$\begin{aligned} \Pr(IC_+ < IC) &= \Pr\left\{c(n, p+1) - 2\hat{\ell}_+ < c(n, p) - 2\hat{\ell}\right\} \\ &= \Pr\left\{2(\hat{\ell}_+ - \hat{\ell}) > c(n, p+1) - c(n, p)\right\}. \end{aligned}$$

and in large samples

$$\text{for AIC, } c(n, p+1) - c(n, p) = 2$$

$$\text{for NIC, } c(n, p+1) - c(n, p) \sim 2$$

$$\text{for BIC, } c(n, p+1) - c(n, p) = \log n$$

- In a regular case  $2(\hat{\ell}_+ - \hat{\ell}) \sim \chi_1^2$ , so as  $n \rightarrow \infty$ ,

$$\Pr(IC_+ < IC) \rightarrow \begin{cases} 0.16, & \text{AIC, NIC,} \\ 0, & \text{BIC.} \end{cases}$$

Thus AIC and NIC have non-zero probability of over-fitting, even in very large samples, but BIC does not.

## Linear Model

slide 24

### Variable selection

- Consider normal linear model

$$Y_{n \times 1} = X_{n \times p}^\dagger \beta_{p \times 1} + \varepsilon_{n \times 1}, \quad \varepsilon \sim \mathcal{N}_n(0, \sigma^2 I_n),$$

where **design matrix**  $X^\dagger$  has full rank  $p < n$  and columns  $x_r$ , for  $r \in \mathcal{X} = \{1, \dots, p\}$ . Subsets  $\mathcal{S}$  of  $\mathcal{X}$  correspond to subsets of columns.

- Terminology
  - the **true** model corresponds to subset  $\mathcal{T} = \{r : \beta_r \neq 0\}$ , and  $|\mathcal{T}| = q < p$ ;
  - a **correct** model contains  $\mathcal{T}$  but has other columns also, corresponding subset  $\mathcal{S}$  satisfies  $\mathcal{T} \subset \mathcal{S} \subset \mathcal{X}$  and  $\mathcal{T} \neq \mathcal{S}$ ;
  - a **wrong** model has subset  $\mathcal{S}$  lacking some  $x_r$  for which  $\beta_r \neq 0$ , and so  $\mathcal{T} \not\subset \mathcal{S}$ .
- Aim to identify  $\mathcal{T}$ .
- If we choose a wrong model, have bias; if we choose a correct model, increase variance—seek to balance these.

## Stepwise methods

- **Forward selection:** starting from model with constant only,
  1. add each remaining term separately to the current model;
  2. if none of these terms is significant, stop; otherwise
  3. update the current model to include the most significant new term; go to 1
- **Backward elimination:** starting from model with all terms,
  1. if all terms are significant, stop; otherwise
  2. update current model by dropping the term with the smallest  $F$  statistic; go to 1
- **Stepwise:** starting from an arbitrary model,
  1. consider 3 options—add a term, delete a term, swap a term in the model for one not in the model;
  2. if model unchanged, stop; otherwise go to 1

APTS: Statistical Modelling

April 2010 – slide 26

## Nuclear power station data

```
> nuclear
  cost  date t1 t2  cap pr ne ct bw cum.n pt
1 460.05 68.58 14 46  687 0 1 0 0   14 0
2 452.99 67.33 10 73 1065 0 0 1 0    1 0
3 443.22 67.33 10 85 1065 1 0 1 0    1 0
4 652.32 68.00 11 67 1065 0 1 1 0   12 0
5 642.23 68.00 11 78 1065 1 1 1 0   12 0
6 345.39 67.92 13 51  514 0 1 1 0    3 0
7 272.37 68.17 12 50  822 0 0 0 0    5 0
8 317.21 68.42 14 59  457 0 0 0 0    1 0
9 457.12 68.42 15 55  822 1 0 0 0    5 0
10 690.19 68.33 12 71  792 0 1 1 1    2 0
...
32 270.71 67.83  7 80  886 1 0 0 1   11 1
```

APTS: Statistical Modelling

April 2010 – slide 27

## Nuclear power station data

	Full model		Backward		Forward	
	Est (SE)	<i>t</i>	Est (SE)	<i>t</i>	Est (SE)	<i>t</i>
Constant	-14.24 (4.229)	-3.37	-13.26 (3.140)	-4.22	-7.627 (2.875)	-2.66
date	0.209 (0.065)	3.21	0.212 (0.043)	4.91	0.136 (0.040)	3.38
log(T1)	0.092 (0.244)	0.38				
log(T2)	0.290 (0.273)	1.05				
log(cap)	0.694 (0.136)	5.10	0.723 (0.119)	6.09	0.671 (0.141)	4.75
PR	-0.092 (0.077)	-1.20				
NE	0.258 (0.077)	3.35	0.249 (0.074)	3.36		
CT	0.120 (0.066)	1.82	0.140 (0.060)	2.32		
BW	0.033 (0.101)	0.33				
log(N)	-0.080 (0.046)	-1.74	-0.088 (0.042)	-2.11		
PT	-0.224 (0.123)	-1.83	-0.226 (0.114)	-1.99	-0.490 (0.103)	-4.77
<i>s</i> (df)	0.164 (21)		0.159 (25)		0.195 (28)	

Backward selection chooses a model with seven covariates also chosen by minimising AIC.

APTS: Statistical Modelling

April 2010 – slide 28

## Stepwise Methods: Comments

- Systematic search minimising AIC or similar over all possible models is preferable—not always feasible.
- Stepwise methods can fit models to purely random data—main problem is no objective function.
- Sometimes used by replacing  $F$  significance points by (arbitrary!) numbers, e.g.  $F = 4$
- Can be improved by comparing AIC for different models at each step—uses AIC as objective function, but no systematic search.

APTS: Statistical Modelling

April 2010 – slide 29

## Prediction error

- To identify  $\mathcal{T}$ , we fit candidate model

$$Y = X\beta + \varepsilon,$$

where columns of  $X$  are a subset  $\mathcal{S}$  of those of  $X^\dagger$ .

- Fitted value is

$$X\hat{\beta} = X\{(X^T X)^{-1} X^T Y\} = HY = H(\mu + \varepsilon) = H\mu + H\varepsilon,$$

where  $H = X(X^T X)^{-1} X^T$  is the **hat matrix** and  $H\mu = \mu$  if the model is correct.

- Following reasoning for AIC, suppose we also have independent dataset  $Y_+$  from the true model, so  $Y_+ = \mu + \varepsilon_+$
- Apart from constants, previous measure of prediction error is

$$\Delta(X) = n^{-1} \mathbb{E} \mathbb{E}_+ \left\{ (Y_+ - X\hat{\beta})^T (Y_+ - X\hat{\beta}) \right\},$$

with expectations over both  $Y_+$  and  $Y$ .

APTS: Statistical Modelling

April 2010 – slide 30

## Prediction error II

- Can show that

$$\Delta(X) = \begin{cases} n^{-1} \mu^T (I - H) \mu + (1 + p/n) \sigma^2, & \text{wrong model,} \\ (1 + q/n) \sigma^2, & \text{true model,} \\ (1 + p/n) \sigma^2, & \text{correct model;} \end{cases} \quad (6)$$

recall that  $q < p$ .


- **Bias:**  $n^{-1} \mu^T (I - H) \mu > 0$  unless model is correct, and is reduced by including useful terms
- **Variance:**  $(1 + p/n) \sigma^2$  increased by including useless terms
- Ideal would be to choose covariates  $X$  to minimise  $\Delta(X)$ : impossible—depends on unknowns  $\mu, \sigma$ .
- Must estimate  $\Delta(X)$

APTS: Statistical Modelling

April 2010 – slide 31



## Example



$\Delta(X)$  as a function of the number of included variables  $p$  for data with  $n = 20$ ,  $q = 6$ ,  $\sigma^2 = 1$ . The minimum is at  $p = q = 6$ :

- there is a sharp decrease in bias as useful covariates are added;
- there is a slow increase with variance as the number of variables  $p$  increases.

APTS: Statistical Modelling

April 2010 – slide 32

## Cross-validation

- If  $n$  is large, can split data into two parts  $(X', y')$  and  $(X^*, y^*)$ , say, and use one part to estimate model, and the other to compute prediction error; then choose the model that minimises

$$\hat{\Delta} = n'^{-1}(y' - X'\hat{\beta}^*)^T(y' - X'\hat{\beta}^*) = n'^{-1} \sum_{j=1}^{n'} (y'_j - x'_j \hat{\beta}^*)^2.$$

- Usually dataset is too small for this; use **leave-one-out cross-validation** sum of squares

$$n\hat{\Delta}_{CV} = CV = \sum_{j=1}^n (y_j - x_j^T \hat{\beta}_{-j})^2,$$

where  $\hat{\beta}_{-j}$  is estimate computed without  $(x_j, y_j)$ .

- Seems to require  $n$  fits of model, but in fact

$$CV = \sum_{j=1}^n \frac{(y_j - x_j^T \hat{\beta})^2}{(1 - h_{jj})^2},$$

where  $h_{11}, \dots, h_{nn}$  are diagonal elements of  $H$ , and so can be obtained from one fit.

APTS: Statistical Modelling

April 2010 – slide 33

## Cross-validation II

- Simpler (more stable?) version uses **generalised cross-validation** sum of squares

$$\text{GCV} = \sum_{j=1}^n \frac{(y_j - x_j^T \hat{\beta})^2}{\{1 - \text{tr}(H)/n\}^2}.$$

- Can show that

$$E(\text{GCV}) = \mu^T(I - H)\mu/(1 - p/n)^2 + n\sigma^2/(1 - p/n) \approx n\Delta(X) \quad (7)$$

so try and minimise GCV or CV.

- Many variants of cross-validation exist. Typically find that model chosen based on CV is somewhat unstable, and that GCV or  $k$ -fold cross-validation works better. Standard strategy is to split data into 10 roughly equal parts, predict for each part based on the other nine-tenths of the data, and find model that minimises this estimate of prediction error.

APTS: Statistical Modelling

April 2010 – slide 34

## Other selection criteria

- Corrected version of AIC for models with normal responses:

$$\text{AIC}_c \equiv n \log \hat{\sigma}^2 + n \frac{1 + p/n}{1 - (p + 2)/n},$$

where  $\hat{\sigma}^2 = \text{RSS}/n$ . Related (unbiased)  $\text{AIC}_u$  replaces  $\hat{\sigma}^2$  by  $S^2 = \text{RSS}/(n - p)$ .

- Mallows suggested

$$C_p = \frac{SS_p}{s^2} + 2p - n,$$

where  $SS_p$  is RSS for fitted model and  $s^2$  estimates  $\sigma^2$ .

- Comments:

- AIC tends to choose models that are too complicated;  $\text{AIC}_c$  cures this somewhat
- BIC chooses true model with probability  $\rightarrow 1$  as  $n \rightarrow \infty$ , if the true model is fitted.

APTS: Statistical Modelling

April 2010 – slide 35

### Simulation experiment

Number of times models were selected using various model selection criteria in 50 repetitions using simulated normal data for each of 20 design matrices. The true model has  $p = 3$ .


$n$		Number of covariates						
		1	2	3	4	5	6	7
10	$C_p$		131	504	91	63	83	128
	BIC		72	373	97	83	109	266
	AIC		52	329	97	91	125	306
	$AIC_c$	15	398	565	18	4		
20	$C_p$		4	673	121	88	61	53
	BIC		6	781	104	52	30	27
	AIC		2	577	144	104	76	97
	$AIC_c$		8	859	94	30	8	1
40	$C_p$			712	107	73	66	42
	BIC			904	56	20	15	5
	AIC			673	114	90	69	54
	$AIC_c$			786	105	52	41	16

APTS: Statistical Modelling

April 2010 – slide 36

### Simulation experiment

Twenty replicate traces of AIC, BIC, and  $AIC_c$ , for data simulated with  $n = 20$ ,  $p = 1, \dots, 16$ , and  $q = 6$ .




APTS: Statistical Modelling

April 2010 – slide 37

### Simulation experiment

Twenty replicate traces of AIC, BIC, and AIC<sub>c</sub>, for data simulated with  $n = 40$ ,  $p = 1, \dots, 16$ , and  $q = 6$ .




APTS: Statistical Modelling

April 2010 – slide 38

### Simulation experiment

Twenty replicate traces of AIC, BIC, and AIC<sub>c</sub>, for data simulated with  $n = 80$ ,  $p = 1, \dots, 16$ , and  $q = 6$ .



As  $n$  increases, note how

- AIC and AIC<sub>c</sub> still allow some over-fitting, but BIC does not, and
- AIC<sub>c</sub> approaches AIC.

APTS: Statistical Modelling

April 2010 – slide 39

**Motivation**

- 'Traditional' analysis methods presuppose that  $p < n$ , so the number of observations exceeds the number of covariates: tall thin design matrices
- Many modern datasets have design matrices that are short and fat:  $p \gg n$ , so the number of covariates (far) exceeds the number of observations—e.g., survival data ( $n$  a few hundred) with genetic information on individuals ( $p$  many thousands)
- Need approaches to deal with this
- Only possibility is to drop most of the covariates from the analysis, so the model has many fewer active covariates
  - usually impracticable in fitting to have  $p > n$
  - anyway impossible to interpret when  $p$  too large
- Seek **sparse** solutions, in which coefficients of most covariates are set to zero, and only covariates with large coefficients are retained. One way to do this is by thresholding: kill small coefficients, and keep the rest.

APTS: Statistical Modelling

April 2010 – slide 41

**Desiderata**

Would like variable selection procedures that satisfy:

- **sparsity**—small estimates are reduced to zero by a threshold procedure; and
- **near unbiasedness**—the estimators almost provide the true parameters, when these are large and  $n \rightarrow \infty$ ;
- **continuity**—the estimator is continuous in the data, to avoid instability in prediction.

None of the previous approaches is sparse, and stepwise selection (for example) is known to be highly unstable. To overcome this, we consider a **regularised** (or penalised) log likelihood of the form

$$\frac{1}{2} \sum_{j=1}^n \ell_j(x_j^T \beta; y_j) - n \sum_{r=1}^p p_\lambda(|\beta_r|),$$

where  $p_\lambda(|\beta|)$  is a penalty discussed below.

APTS: Statistical Modelling

April 2010 – slide 42

### Example: Lasso

- The **lasso** (least absolute selection and shrinkage operator) chooses  $\beta$  to minimise

$$(y - X\beta)^T(y - X\beta) \text{ such that } \sum_{r=1}^p |\beta_r| \leq \lambda,$$

for some  $\lambda > 0$ ; call resulting estimator  $\tilde{\beta}_\lambda$ .

- $\lambda \rightarrow 0$  implies  $\tilde{\beta}_\lambda \rightarrow 0$ , and  $\lambda \rightarrow \infty$  implies  $\tilde{\beta}_\lambda \rightarrow \hat{\beta} = (X^T X)^{-1} X^T y$ .
- Simple case: orthogonal design matrix  $X^T X = I_p$ , gives


$$\tilde{\beta}_{\lambda,r} = \begin{cases} 0, & |\hat{\beta}_r| < \gamma, \\ \text{sign}(\hat{\beta}_r)(|\hat{\beta}_r| - \gamma), & \text{otherwise,} \end{cases} \quad r = 1, \dots, p. \quad (8)$$

- Call this **soft thresholding**.
- Computed using **least angle regression** algorithm (Efron et al., 2004, Annals of Statistics).

APTS: Statistical Modelling

April 2010 – slide 43

### Soft thresholding




APTS: Statistical Modelling

April 2010 – slide 44

### Graphical explanation

In each case aim to minimise the quadratic function subject to remaining inside the shaded region.




APTS: Statistical Modelling

April 2010 – slide 45

### Lasso: Nuclear power data

Left: traces of coefficient estimates  $\hat{\beta}_\lambda$  as constraint  $\lambda$  is relaxed, showing points at which the different covariates enter the model. Right: behaviour of Mallows'  $C_p$  as  $\lambda$  increases.



APTS: Statistical Modelling

April 2010 – slide 46

## Penalties

Some (of many) possible penalty functions  $p_\lambda(|\beta|)$ , all with  $\lambda > 0$ :

- ridge regression** takes  $\lambda|\beta|^2$ ;
- lasso** takes  $\lambda|\beta|$ ;
- elastic net** takes  $\lambda\{(1 - \alpha)|\beta| + \alpha|\beta|^2\}$ , with  $0 \leq \alpha < 1$ ;
- bridge regression** takes  $\lambda|\beta|^q$ , for  $q > 0$ ;
- hard threshold** takes  $\lambda^2 - (|\beta| - \lambda)^2 I(|\beta| < \lambda)$ ;
- smoothly clipped absolute deviation (SCAD)** takes

$$\begin{cases} \lambda|\beta|, & |\beta| < \lambda, \\ -(\beta^2 - 2a\lambda|\beta| + \lambda^2)/\{2(a - 1)\}, & \lambda < |\beta| < a\lambda, \\ (a + 1)\lambda^2/2, & |\beta| > a\lambda, \end{cases}$$

for some  $a > 2$ .

In least squares case with a single observation seek to minimise  $\frac{1}{2}(z - \beta)^2 + p_\lambda(|\beta|)$ , whose derivative

$$\text{sign}(\beta)\{|\beta| + \partial p_\lambda(|\beta|)/\partial\beta\} - z$$


determines the properties of the estimator.

APTS: Statistical Modelling

April 2010 – slide 47

## Some threshold functions

- Ridge—shrinkage but no selection; hard threshold—subset selection, unstable; soft threshold—lasso, biased; SCAD—continuous, selection, unbiased for large  $\beta$ , but non-monotone.



APTS: Statistical Modelling

April 2010 – slide 48



## Properties of penalties

It turns out that to achieve

- sparsity**, the minimum of the function  $|\beta| + \partial p_\lambda(|\beta|)/\partial\beta$  must be positive;
- near unbiasedness**, the penalty must satisfy  $\partial p_\lambda(|\beta|)/\partial\beta \rightarrow 0$  when  $|\beta|$  is large, so then the estimating function approaches  $\beta - z$ ; and
- continuity**, the minimum of  $|\beta| + \partial p_\lambda(|\beta|)/\partial\beta$  must be attained at  $\beta = 0$ .

The SCAD is constructed to have these properties, but there is no unique minimum to the resulting objective function, so numerically it is awkward.

APTS: Statistical Modelling

April 2010 – slide 49

## Oracle

- Oracle:

A person or thing regarded as an infallible authority or guide.

- A **statistical oracle** says how to choose the model or bandwidth that will give us optimal estimation of the true parameter or function, but not the truth itself.
- In the context of variable selection, an oracle tells us which variables we should select, but not their coefficients.
- It turns out that under mild conditions on the model, and provided  $\lambda \equiv \lambda_n \rightarrow 0$  and  $\sqrt{n}\lambda_n \rightarrow \infty$  as  $n \rightarrow \infty$ , variable selection using the hard and SCAD penalties has an oracle property: the estimators of  $\beta$  work as well as *if* we had known in advance which covariates should be excluded.
- Same ideas extend to generalised linear models, survival analysis, and many other regression settings (Fan and Li, 2001, JASA).
- Harder: what happens when  $p \rightarrow \infty$  also?

APTS: Statistical Modelling

April 2010 – slide 50

**Thomas Bayes (1702–1761)**



Bayes (1763/4) *Essay towards solving a problem in the doctrine of chances*. Philosophical Transactions of the Royal Society of London.

APTS: Statistical Modelling

April 2010 – slide 52

**Bayesian inference**

Parametric model for data  $y$  assumed to be realisation of  $Y \sim f(y; \theta)$ , where  $\theta \in \Omega_\theta$ .

**Frequentist viewpoint** (cartoon version):

- there is a true value of  $\theta$  that generated the data;
- this 'true' value of  $\theta$  is to be treated as an unknown constant;
- probability statements concern randomness in hypothetical replications of the data (possibly conditioned on an ancillary statistic).

**Bayesian viewpoint** (cartoon version):

- all ignorance may be expressed in terms of probability statements;
- a joint probability distribution for data and all unknowns can be constructed;
- Bayes' theorem should be used to convert prior beliefs  $\pi(\theta)$  about unknown  $\theta$  into posterior beliefs  $\pi(\theta | y)$ , conditioned on data;
- probability statements concern randomness of unknowns, conditioned on all known quantities.

APTS: Statistical Modelling

April 2010 – slide 53

## Mechanics

- Separate from data, we have prior information about parameter  $\theta$  summarised in density  $\pi(\theta)$
- Data model  $f(y | \theta) \equiv f(y; \theta)$
- Posterior density given by Bayes' theorem:

$$\pi(\theta | y) = \frac{\pi(\theta)f(y | \theta)}{\int \pi(\theta)f(y | \theta) d\theta}.$$

- $\pi(\theta | y)$  contains all information about  $\theta$ , conditional on observed data  $y$
- If  $\theta = (\psi, \lambda)$ , then inference for  $\psi$  is based on **marginal posterior density**

$$\pi(\psi | y) = \int \pi(\theta | y) d\lambda$$

APTS: Statistical Modelling

April 2010 – slide 54

## Encompassing model

- Suppose we have  $M$  alternative models for the data, with respective parameters  $\theta_1 \in \Omega_{\theta_1}, \dots, \theta_m \in \Omega_{\theta_m}$ . Typically dimensions of  $\Omega_{\theta_m}$  are different.
- We enlarge the parameter space to give an **encompassing model** with parameter

$$\theta = (m, \theta_m) \in \Omega = \bigcup_{m=1}^M \{m\} \times \Omega_{\theta_m}.$$

- Thus need priors  $\pi_m(\theta_m | m)$  for the parameters of each model, plus a prior  $\pi(m)$  giving pre-data probabilities for each of the models; overall

$$\pi(m, \theta_m) = \pi(\theta_m | m)\pi(m) = \pi_m(\theta_m)\pi_m,$$

say.

- Inference about model choice is based on marginal posterior density

$$\pi(m | y) = \frac{\int f(y | \theta_m)\pi_m(\theta_m)\pi_m d\theta_m}{\sum_{m'=1}^M \int f(y | \theta_{m'})\pi_{m'}(\theta_{m'})\pi_{m'} d\theta_{m'}} = \frac{\pi_m f(y | m)}{\sum_{m'=1}^M \pi_{m'} f(y | m')}.$$

APTS: Statistical Modelling

April 2010 – slide 55

## Inference

- Can write

$$\pi(m, \theta_m | y) = \pi(\theta_m | y, m)\pi(m | y),$$

so Bayesian updating corresponds to

$$\pi(\theta_m | m)\pi(m) \mapsto \pi(\theta_m | y, m)\pi(m | y)$$

and for each model  $m = 1, \dots, M$  we need

- posterior probability  $\pi(m | y)$ , which involves the marginal likelihood  $f(y | m) = \int f(y | \theta_m, m)\pi(\theta_m | m) d\theta_m$ ; and
- the posterior density  $f(\theta_m | y, m)$ .

- If there are just two models, can write

$$\frac{\pi(1 | y)}{\pi(2 | y)} = \frac{\pi_1 f(y | 1)}{\pi_2 f(y | 2)},$$

so the posterior odds on model 1 equal the prior odds on model 1 multiplied by the **Bayes factor**  $B_{12} = f(y | 1)/f(y | 2)$ .

APTS: Statistical Modelling

April 2010 – slide 56

## Sensitivity of the marginal likelihood

Suppose the prior for each  $\theta_m$  is  $\mathcal{N}(0, \sigma^2 I_{d_m})$ , where  $d_m = \dim(\theta_m)$ . Then, dropping the  $m$  subscript for clarity,

$$\begin{aligned} f(y | m) &= \sigma^{-d/2} (2\pi)^{-d/2} \int f(y | m, \theta) \prod_r \exp\{-\theta_r^2 / (2\sigma^2)\} d\theta_r \\ &\approx \sigma^{-d/2} (2\pi)^{-d/2} \int f(y | m, \theta) \prod_r d\theta_r, \end{aligned}$$

for a highly diffuse prior distribution (large  $\sigma^2$ ). The Bayes factor for comparing the models is approximately

$$\frac{f(y | 1)}{f(y | 2)} \approx \sigma^{(d_2 - d_1)/2} g(y),$$

where  $g(y)$  depends on the two likelihoods but is independent of  $\sigma^2$ . Hence, *whatever the data tell us about the relative merits of the two models*, the Bayes factor in favour of the simpler model can be made arbitrarily large by increasing  $\sigma$ .

This illustrates **Lindley's paradox**, and implies that we must be careful when specifying prior dispersion parameters to compare models.

APTS: Statistical Modelling

April 2010 – slide 57

## Model averaging

- If a quantity  $Z$  has the same interpretation for all models, it may be necessary to allow for model uncertainty:
  - in prediction, each model may be just a vehicle that provides a future value, not of interest *per se*;
  - physical parameters (means, variances, etc.) may be suitable for averaging, but care is needed.
- The predictive distribution for  $Z$  may be written

$$f(z | y) = \sum_{m=1}^M f(z | y, m) \Pr(m | y)$$

where


$$\Pr(m | y) = \frac{f(y | m) \Pr(m)}{\sum_{m'=1}^M f(y | m') \Pr(m')}$$

APTS: Statistical Modelling

April 2010 – slide 58

## Example: Cement data

Percentage weights in clinkers of 4 constituents of cement ( $x_1, \dots, x_4$ ) and heat evolved  $y$  in calories, in  $n = 13$  samples.



APTS: Statistical Modelling

April 2010 – slide 59

### Example: Cement data

```
> cement
  x1 x2 x3 x4    y
1   7 26  6 60  78.5
2   1 29 15 52  74.3
3  11 56  8 20 104.3
4  11 31  8 47  87.6
5   7 52  6 33  95.9
6  11 55  9 22 109.2
7   3 71 17  6 102.7
8   1 31 22 44  72.5
9   2 54 18 22  93.1
10 21 47  4 26 115.9
11  1 40 23 34  83.8
12 11 66  9 12 113.3
13 10 68  8 12 109.4
```

APTS: Statistical Modelling

April 2010 – slide 60

### Example: Cement data

Bayesian model choice and prediction using model averaging for the cement data ( $n = 13, p = 4$ ). For each of the 16 possible subsets of covariates, the table shows the log Bayes factor in favour of that subset compared to the model with no covariates and gives the posterior probability of each model. The values of the posterior mean and scale parameters  $a$  and  $b$  are also shown for the six most plausible models;  $(y_+ - a)/b$  has a posterior  $t$  density. For comparison, the residual sums of squares are also given.


Model	RSS	$2 \log B_{10}$	$\Pr(M   y)$	$a$	$b$
----	2715.8	0.0	0.0000		
1---	1265.7	7.1	0.0000		
-2--	906.3	12.2	0.0000		
--3-	1939.4	0.6	0.0000		
---4	883.9	12.6	0.0000		
12--	57.9	45.7	0.2027	93.77	2.31
1-3-	1227.1	4.0	0.0000		
1--4	74.8	42.8	0.0480	99.05	2.58
-23-	415.4	19.3	0.0000		
-2-4	868.9	11.0	0.0000		
--34	175.7	31.3	0.0002		
123-	48.11	43.6	0.0716	95.96	2.80
12-4	47.97	47.2	0.4344	95.88	2.45
1-34	50.84	44.2	0.0986	94.66	2.89
-234	73.81	33.2	0.0004		
1234	47.86	45.0	0.1441	95.20	2.97

APTS: Statistical Modelling

April 2010 – slide 61

### Example: Cement data

Posterior predictive densities for cement data. Predictive densities for a future observation  $y_+$  with covariate values  $x_+$  based on individual models are given as dotted curves. The heavy curve is the average density from all 16 models.



APTS: Statistical Modelling

April 2010 – slide 62

### DIC

- How to compare complex models (e.g. hierarchical models, mixed models, Bayesian settings), in which the 'number of parameters' may:
  - outnumber the number of observations?
  - be unclear because of the regularisation provided by a prior density?
- Suppose model has 'Bayesian deviance'

$$D(\theta) = -2 \log f(y | \theta) + 2 \log f(y)$$

for some normalising function  $f(y)$ , and suppose that samples from the posterior density of  $\theta$  are available and give  $\bar{\theta} = E(\theta | y)$ .

- One possibility is the **deviance information criterion (DIC)**

$$D(\bar{\theta}) + 2p_D,$$

where the number of associated parameters is

$$p_D = \overline{D(\theta)} - D(\bar{\theta}).$$

- This involves only (MCMC) samples from the posterior, no analytical computations, and reproduces AIC for some classes of models.

APTS: Statistical Modelling

April 2010 – slide 63

## Minimum description length

Model selection can also be based on related ideas of **minimum description length (MDL)** or **minimum message length (MML)**, which use ideas from computer science—coding and information theory:

- idea is to choose encoding of data that minimises length of equivalent binary sequence, regarding all data as discrete;
- minimum message includes parameter estimates, data using optimal code based on parameter estimates, (and prior information);
- close links to AIC, BIC, etc.;
- see <http://www.mdl-research.org/> or tutorial on <http://homepages.cwi.nl/~pdg/ftp/mdlintro.pdf> to learn more.

APTS: Statistical Modelling

April 2010 – slide 64

## Bayesian Variable Selection

slide 65

### Variable selection

- In Bayesian context, must determine prior probability for the inclusion (or not) of each variable in the model.
- Common to use 'spike and slab' prior for coefficient  $\theta$ :

$$\theta = \begin{cases} 0, & \text{with probability } 1 - p \\ \mathcal{N}(0, \tau^2), & \text{with probability } p, \end{cases}$$

corresponding to prior 'density'

$$\pi(\theta) = (1 - p)\delta(\theta) + p\tau^{-1}\phi(\theta/\tau), \quad \theta \in \mathbb{R},$$

where  $\delta(\theta)$  is the delta function putting unit mass at  $\theta = 0$ .


- Now find posterior for  $\beta$  based on data.
- Usually independent priors for each covariate, and typically need clever (dimension-jumping) MCMC.

APTS: Statistical Modelling

April 2010 – slide 66



## Example: NMR data



Left: original data, with  $n = 1024$

Right: orthogonal transformation into  $n = 1024$  coefficients at different resolutions

APTS: Statistical Modelling

April 2010 – slide 67

## Orthogonal transformation

- Model:** original data  $X \sim \mathcal{N}_n(\mu, \sigma^2 I_n)$ , where signal  $\mu_{n \times 1}$  is perturbed by normal noise, giving noisy data  $X_{n \times 1}$
- set  $Y_{n \times 1} = W_{n \times n} X_{n \times 1}$ , where  $W^T W = W W^T = I_n$  is orthogonal
- choose  $W$  so that  $\theta = W \mu$  should be 'sparse' (i.e. most elements of  $\theta$  are zero)—good choice is wavelet coefficients (mathematical compression properties)
- 'kill' small coefficients of  $Y$ , which correspond to noise, giving  $\tilde{\theta}_{n \times 1} = \text{kill}(Y) = \text{kill}(WX)$ , say, then
- estimate signal  $\mu$  by

$$\tilde{\mu} = W^T \tilde{\theta} = W^T (\text{kill}(WX)).$$

APTS: Statistical Modelling

April 2010 – slide 68

## Posterior

If given  $\theta$ ,  $Y \sim \mathcal{N}(\theta, \sigma^2)$ , then the posterior 'density' is of form

$$\pi(\theta | y) = (1 - p_y) \delta(\theta) + p_y b^{-1} \phi\left(\frac{\theta - ay}{b}\right), \quad \theta \in \mathbb{R},$$

where

$$a = \tau^2 / (\tau^2 + \sigma^2), \quad b^2 = 1 / (1/\sigma^2 + 1/\tau^2),$$

and

$$p_y = \frac{p(\sigma^2 + \tau^2)^{-1/2} \phi\{y / (\sigma^2 + \tau^2)^{1/2}\}}{(1 - p)\sigma^{-1} \phi(y/\sigma) + p(\sigma^2 + \tau^2)^{-1/2} \phi\{y / (\sigma^2 + \tau^2)^{1/2}\}}$$

is the posterior probability that  $\theta \neq 0$ .

Summary statistic: **posterior median**  $\tilde{\theta}$ , for which  $\Pr(\theta \leq \tilde{\theta} | y) = 0.5$ . For small  $|y|$ , this gives  $\tilde{\theta} = 0$ . (Next slide)

APTS: Statistical Modelling

April 2010 – slide 69


## Shrinkage

Prior CDF of  $\theta$  (left), and posterior CDFs when  $p = 0.5, \sigma = \tau = 1$ , and  $y = -2.5$  (centre), and  $y = -1$  (right).

Red horizontal line: cumulative probability=0.5

Blue vertical line: data  $y$

Green vertical line: posterior median  $\tilde{\theta}$



APTS: Statistical Modelling

April 2010 – slide 70

## Empirical Bayes

The parameters  $p, \sigma, \tau$  are unknown. We estimate them by empirical Bayes:

- we note that the marginal density of  $y$  is

$$f(y) = (1-p)\sigma^{-1}\phi(y/\sigma) + p(\sigma^2 + \tau^2)^{-1/2}\phi\{y/(\sigma^2 + \tau^2)^{1/2}\}, \quad y \in \mathbb{R},$$

so if we have  $y_1, \dots, y_n \stackrel{\text{iid}}{\sim} f$  we estimate  $p, \sigma, \tau$  by maximising the log likelihood


$$\ell(p, \sigma, \tau) = \sum_{j=1}^n \log f(y_j; p, \sigma, \tau).$$

- Here we obtain  $\tilde{p} = 0.04$ ,  $\tilde{\sigma} = 2.1$ , and  $\tilde{\tau} = 52.1$ .
- Now compute the posterior medians  $\tilde{\theta}_j$  corresponding to each  $y_j$ .

APTS: Statistical Modelling

April 2010 – slide 71


### Example: NMR data



APTS: Statistical Modelling

April 2010 – slide 72

### Example: NMR data



APTS: Statistical Modelling

April 2010 – slide 73

### Comments

- Large and rapidly-growing literature on Bayesian 'variable' selection, now particularly focused on 'large  $p$ , small  $n$ ' paradigm
- Close relation to classical 'super-efficient' estimation: James–Stein theorem, Hodges–Lehmann estimator, biased (but lower loss) estimation

APTS: Statistical Modelling

April 2010 – slide 74

## 2. Beyond the Generalised Linear Model

slide 75

### Overview

1. Generalised linear models
2. Overdispersion
3. Correlation
4. Random effects models
5. Conditional independence and graphical representations

APTS: Statistical Modelling

April 2010 – slide 76

## Generalised Linear Models

slide 77

### GLM recap

$y_1, \dots, y_n$  are observations of response variables  $Y_1, \dots, Y_n$  assumed to be independently generated by a distribution of the same exponential family form, with means  $\mu_i \equiv E(Y_i)$  linked to explanatory variables  $X_1, X_2, \dots, X_p$  through

$$g(\mu_i) = \eta_i \equiv \beta_0 + \sum_{r=1}^p \beta_r x_{ir} \equiv x_i^T \beta$$

GLMs have proved remarkably effective at modelling real world variation in a wide range of application areas.

APTS: Statistical Modelling

April 2010 – slide 78

### GLM failure

However, situations frequently arise where GLMs do not adequately describe observed data. This can be due to a number of reasons including:

- The mean model cannot be appropriately specified as there is dependence on an unobserved (or unobservable) explanatory variable.
- There is excess variability between experimental units beyond that implied by the mean/variance relationship of the chosen response distribution.
- The assumption of independence is not appropriate.
- Complex multivariate structure in the data requires a more flexible model class

APTS: Statistical Modelling


April 2010 – slide 79

**Example 1: toxoplasmosis**

The table below gives data on the relationship between rainfall ( $x$ ) and the proportions of people with toxoplasmosis ( $y/m$ ) for 34 cities in El Salvador.

City	$y$	$x$	City	$y$	$x$	City	$y$	$x$
1	5/18	1620	12	3/5	1800	23	3/10	1973
2	15/30	1650	13	8/10	1800	24	1/6	1976
3	0/1	1650	14	0/1	1830	25	1/5	2000
4	2/4	1735	15	53/75	1834	26	0/1	2000
5	2/2	1750	16	7/16	1871	27	7/24	2050
6	2/8	1750	17	24/51	1890	28	46/82	2063
7	2/12	1756	18	3/10	1900	29	7/19	2077
8	6/11	1770	19	23/43	1918	30	9/13	2100
9	33/54	1770	20	3/6	1920	31	4/22	2200
10	8/13	1780	21	0/1	1920	32	4/9	2240
11	41/77	1796	22	3/10	1936	33	8/11	2250
						34	23/37	2292

**Example**



Toxoplasmosis data and fitted models

### Example

Fitting various binomial logistic regression models relating toxoplasmosis incidence to rainfall:

Model	df	deviance
Constant	33	74.21
Linear	32	74.09
Quadratic	31	74.09
Cubic	30	62.62

So evidence in favour of the cubic over other models, but a poor fit ( $\chi^2 = 58.21$  on 30df).


This is an example of **overdispersion** where residual variability is greater than would be predicted by the specified mean/variance relationship

$$\text{var}(Y) = \frac{\mu(1 - \mu)}{m}.$$

APTS: Statistical Modelling

April 2010 – slide 83

### Example



Toxoplasmosis residual plot

APTS: Statistical Modelling

April 2010 – slide 84

## Quasi-likelihood

A quasi-likelihood approach to accounting for overdispersion models the mean and variance, but stops short of a full probability model for  $Y$ .

For a model specified by the mean relationship  $g(\mu_i) = \eta_i = x_i^T \beta$ , and variance  $\text{var}(Y_i) = \sigma^2 V(\mu_i)/m_i$ , the quasi-likelihood equations are

$$\sum_{i=1}^n x_i \frac{y_i - \mu_i}{\sigma^2 V(\mu_i) g'(\mu_i)/m_i} = 0$$

If  $V(\mu_i)/m_i$  represents  $\text{var}(Y_i)$  for a standard distribution from the exponential family, then these equations can be solved for  $\beta$  using standard GLM software.

Provided the mean and variance functions are correctly specified, asymptotic normality for  $\hat{\beta}$  still holds. The dispersion parameter  $\sigma^2$  can be estimated using

$$\hat{\sigma}^2 \equiv \frac{1}{n - p - 1} \sum_{i=1}^n \frac{m_i (y_i - \hat{\mu}_i)^2}{V(\hat{\mu}_i)}$$

APTS: Statistical Modelling

April 2010 – slide 85

## Quasi-likelihood for toxoplasmosis data

Assuming the same mean model as before, but  $\text{var}(Y_i) = \sigma^2 \mu_i (1 - \mu_i)/m_i$ , we obtain  $\hat{\sigma}^2 = 1.94$  with  $\hat{\beta}$  (and corresponded fitted mean curves) as before.

Comparing cubic with constant model, one now obtains

$$F = \frac{(74.21 - 62.62)/3}{1.94} = 1.99$$

which provides much less compelling evidence in favour of an effect of rainfall on toxoplasmosis incidence.

APTS: Statistical Modelling

April 2010 – slide 86

## Reasons

To construct a full probability model in the presence of overdispersion, it is necessary to consider **why** overdispersion might be present.

Possible reasons include:

- There may be an important explanatory variable, other than rainfall, which we haven't observed.
- Or there may be many other features of the cities, possibly unobservable, all having a small individual effect on incidence, but a larger effect in combination. Such effects may be individually undetectable – sometimes described as *natural excess variability between units*.

APTS: Statistical Modelling

April 2010 – slide 87

### Reasons: unobserved heterogeneity

When part of the linear predictor is 'missing' from the model,

$$\eta_i^{\text{true}} = \eta_i^{\text{model}} + \eta_i^{\text{diff}}$$

We can compensate for this, in modelling, by assuming that the missing  $\eta_i^{\text{diff}} \sim F$  in the population. Hence, given  $\eta_i^{\text{model}}$

$$\mu_i \equiv g^{-1}(\eta_i^{\text{model}} + \eta_i^{\text{diff}}) \sim G$$

where  $G$  is the distribution induced by  $F$ . Then

$$E(Y_i) = E_G[E(Y_i | \mu_i)] = E_G(\mu_i)$$

$$\text{var}(Y_i) = E_G(V(\mu_i)/m_i) + \text{var}_G(\mu_i)$$

APTS: Statistical Modelling

April 2010 – slide 88

### Direct models

One approach is to model the  $Y_i$  directly, by specifying an appropriate form for  $G$ .

For example, for the toxoplasmosis data, we might specify a **beta-binomial** model, where

$$\mu_i \sim \text{Beta}(k\mu_i^*, k[1 - \mu_i^*])$$

leading to

$$E(Y_i) = \mu_i^*, \quad \text{var}(Y_i) = \frac{\mu_i^*(1 - \mu_i^*)}{m_i} \left( 1 + \frac{m_i - 1}{k + 1} \right)$$

with  $(m_i - 1)/(k + 1)$  representing an overdispersion factor.

APTS: Statistical Modelling

April 2010 – slide 89

### Direct models: fitting

Models which explicitly account for overdispersion can, in principle, be fitted using your preferred approach, e.g. the beta-binomial model has likelihood

$$f(y | \mu^*, k) \propto \prod_{i=1}^n \frac{\Gamma(k\mu_i^* + m_i y_i) \Gamma\{k(1 - \mu_i^*) + m_i(1 - y_i)\} \Gamma(k)}{\Gamma(k\mu_i^*) \Gamma\{k(1 - \mu_i^*)\} \Gamma(k + m_i)}.$$

Similarly the corresponding model for count data specifies a gamma distribution for the Poisson mean, leading to a *negative binomial* marginal distribution for  $Y_i$ .

However, these models have limited flexibility and can be difficult to fit, so an alternative approach is usually preferred.

APTS: Statistical Modelling

April 2010 – slide 90



## A random effects model for overdispersion

A more flexible, and extensible approach models the excess variability by including an extra term in the linear predictor

$$\eta_i = x_i^T \beta + u_i \quad (9)$$

where the  $u_i$  can be thought of as representing the 'extra' variability between units, and are called **random effects**.

The model is completed by specifying a distribution  $F$  for  $u_i$  in the population – almost always, we use

$$u_i \sim N(0, \sigma^2)$$

for some unknown  $\sigma^2$ .

We set  $E(u_i) = 0$ , as an unknown mean for  $u_i$  would be unidentifiable in the presence of the intercept parameter  $\beta_0$ .

APTS: Statistical Modelling

April 2010 – slide 91

## Random effects: likelihood

The parameters of this random effects model are usually considered to be  $(\beta, \sigma^2)$  and therefore the likelihood is given by

$$\begin{aligned} f(\mathbf{y} \mid \beta, \sigma^2) &= \int f(\mathbf{y} \mid \beta, \mathbf{u}, \sigma^2) f(\mathbf{u} \mid \beta, \sigma^2) d\mathbf{u} \\ &= \int f(\mathbf{y} \mid \beta, \mathbf{u}) f(\mathbf{u} \mid \sigma^2) d\mathbf{u} \\ &= \int \prod_{i=1}^n f(y_i \mid \beta, u_i) f(u_i \mid \sigma^2) du_i \end{aligned} \quad (10)$$

where  $f(y_i \mid \beta, u_i)$  arises from our chosen exponential family, with linear predictor (9) and  $f(u_i \mid \sigma^2)$  is a univariate normal p.d.f.

Often no further simplification of (10) is possible, so computation needs careful consideration – we will come back to this later.

APTS: Statistical Modelling

April 2010 – slide 92

**Toxoplasmosis example revisited**

We can think of the toxoplasmosis proportions  $Y_i$  in each city ( $i$ ) as arising from the sum of binary variables,  $Y_{ij}$ , representing the toxoplasmosis status of individuals ( $j$ ), so  $m_i Y_i = \sum_{j=1}^{m_i} Y_{ij}$ . Then

$$\begin{aligned} \text{var}(Y_i) &= \frac{1}{m_i^2} \sum_{j=1}^{m_i} \text{var}(Y_{ij}) + \frac{1}{m_i^2} \sum_{j \neq k} \text{cov}(Y_{ij}, Y_{ik}) \\ &= \frac{\mu_i(1 - \mu_i)}{m_i} + \frac{1}{m_i^2} \sum_{j \neq k} \text{cov}(Y_{ij}, Y_{ik}) \end{aligned}$$

So any positive correlation between individuals induces overdispersion in the counts.

**Dependence: reasons**

There may be a number of plausible reasons why the responses corresponding to units within a given **cluster** are dependent (in the toxoplasmosis example, cluster = city)

One compelling reason is the unobserved heterogeneity discussed previously.

In the 'correct' model (corresponding to  $\eta_i^{\text{true}}$ ), the toxoplasmosis status of individuals,  $Y_{ij}$ , are independent, so

$$Y_{ij} \perp\!\!\!\perp Y_{ik} \mid \eta_i^{\text{true}} \Leftrightarrow Y_{ij} \perp\!\!\!\perp Y_{ik} \mid \eta_i^{\text{model}}, \eta_i^{\text{diff}}$$

However, in the absence of knowledge of  $\eta_i^{\text{diff}}$

$$Y_{ij} \not\perp\!\!\!\perp Y_{ik} \mid \eta_i^{\text{model}}$$

Hence conditional (given  $\eta_i^{\text{diff}}$ ) independence between units in a common cluster  $i$  becomes marginal dependence, when marginalised over the population distribution  $F$  of unobserved  $\eta_i^{\text{diff}}$ .

**Random effects and dependence**

The correspondence between positive intra-cluster correlation and unobserved heterogeneity suggests that intra-cluster dependence might be modelled using random effects, For example, for the individual-level toxoplasmosis data

$$Y_{ij} \overset{\text{ind}}{\sim} \text{Bernoulli}(\mu_{ij}), \quad \log \frac{\mu_{ij}}{1 - \mu_{ij}} = x_{ij}^T \beta + u_i, \quad u_i \sim N(0, \sigma^2)$$

which implies

$$Y_{ij} \not\perp\!\!\!\perp Y_{ik} \mid \beta, \sigma^2$$

Intra-cluster dependence arises in many applications, and random effects provide an effective way of modelling it.

## Marginal models

Random effects modelling is not the only way of accounting for intra-cluster dependence.

A **marginal model** models  $\mu_{ij} \equiv E(Y_{ij})$  as a function of explanatory variables, through  $g(\mu_{ij}) = x_{ij}^T \beta$ , and also specifies a variance relationship  $\text{var}(Y_{ij}) = \sigma^2 V(\mu_{ij})/m_{ij}$  and a model for  $\text{corr}(Y_{ij}, Y_{ik})$ , as a function of  $\mu$  and possibly additional parameters.

It is important to note that the parameters  $\beta$  in a marginal model have a different interpretation from those in a random effects model, because for the latter

$$E(Y_{ij}) = E(g^{-1}[x_{ij}^T \beta + u_i]) \neq g^{-1}(x_{ij}^T \beta) \quad (\text{unless } g \text{ is linear}).$$

- A random effects model describes the mean response at the subject level ('subject specific')
- A marginal model describes the mean response across the population ('population averaged')

APTS: Statistical Modelling

April 2010 – slide 97

## GEEs

As with the quasi-likelihood approach above, marginal models do not generally provide a full probability model for  $Y$ . Nevertheless,  $\beta$  can be estimated using **generalised estimating equations (GEEs)**.

The GEE for estimating  $\beta$  in a marginal model is of the form

$$\sum_i \left( \frac{\partial \mu_i}{\partial \beta} \right)^T \text{var}(Y_i)^{-1} (Y_i - \mu_i) = 0$$

where  $Y_i = (Y_{ij})$  and  $\mu_i = (\mu_{ij})$

Consistent covariance estimates are available for GEE estimators.

Furthermore, the approach is generally robust to mis-specification of the correlation structure.

For the rest of this module, we focus on fully specified probability models.

APTS: Statistical Modelling

April 2010 – slide 98

## Clustered data

Examples where data are collected in clusters include:

- Studies in biometry where **repeated measures** are made on experimental units. Such studies can effectively mitigate the effect of between-unit variability on important inferences.
- Agricultural field trials, or similar studies, for example in engineering, where experimental units are arranged within **blocks**
- Sample surveys where collecting data within clusters or **small areas** can save costs

Of course, other forms of dependence exist, for example spatial or serial dependence induced by arrangement in space or time of units of observation. This will be a focus of APTS: Spatial and Longitudinal Data Analysis.

APTS: Statistical Modelling

April 2010 – slide 99

## Example 2: Rat growth


The table below is extracted from a data set giving the weekly weights of 30 young rats.

Rat	Week				
	1	2	3	4	5
1	151	199	246	283	320
2	145	199	249	293	354
3	147	214	263	312	328
4	155	200	237	272	297
5	135	188	230	280	323
6	159	210	252	298	331
7	141	189	231	275	305
8	159	201	248	297	338
...	...	...	...	...	...
30	153	200	244	286	324

APTS: Statistical Modelling

April 2010 – slide 100

## Example



Rat growth data

APTS: Statistical Modelling

April 2010 – slide 101


## A simple model

Letting  $Y$  represent weight, and  $X$  represent week, we can fit the simple linear regression

$$y_{ij} = \beta_0 + \beta_1 x_{ij} + \epsilon_{ij}$$

with resulting estimates  $\hat{\beta}_0 = 156.1$  (2.25) and  $\hat{\beta}_1 = 43.3$  (0.92)

Residuals show clear evidence of an unexplained difference between rats



APTS: Statistical Modelling


April 2010 – slide 102

## Model elaboration

Naively adding a (fixed) effect for animal gives

$$y_{ij} = \beta_0 + \beta_1 x_{ij} + u_i + \epsilon_{ij}.$$

Residuals show evidence of a further unexplained difference between rats in terms of dependence on  $x$ .



More complex cluster dependence required.

APTS: Statistical Modelling

April 2010 – slide 103

**Linear mixed models**

A linear mixed model (LMM) for observations  $y = (y_1, \dots, y_n)$  has the general form

$$Y \sim N(\mu, \Sigma), \quad \mu = X\beta + Zb, \quad b \sim N(0, \Sigma_b), \tag{11}$$

where  $X$  and  $Z$  are matrices containing values of explanatory variables. Usually,  $\Sigma = \sigma^2 I_n$ .

A typical example for clustered data might be

$$Y_{ij} \stackrel{\text{ind}}{\sim} N(\mu_{ij}, \sigma^2), \quad \mu_{ij} = x_{ij}^T \beta + z_{ij}^T b_i, \quad b_i \stackrel{\text{ind}}{\sim} N(0, \Sigma_b^*), \tag{12}$$

where  $x_{ij}$  contain the explanatory data for cluster  $i$ , observation  $j$  and (normally)  $z_{ij}$  contains that sub-vector of  $x_{ij}$  which is allowed to exhibit extra between cluster variation in its relationship with  $Y$ . In the simplest (random intercept) case,  $z_{ij} = (1)$ , as in equation (9).

**LMM example**

A plausible LMM for  $k$  clusters with  $n_1, \dots, n_k$  observations per cluster, and a single explanatory variable  $x$  (e.g. the rat growth data) is

$$y_{ij} = \beta_0 + b_{0i} + (\beta_1 + b_{1i})x_{ij} + \epsilon_{ij}, \quad (b_{0i}, b_{1i})^T \stackrel{\text{ind}}{\sim} N(0, \Sigma_b^*).$$

This fits into the general LMM framework (11) with  $\Sigma = \sigma^2 I_n$  and

$$X = \begin{pmatrix} 1 & x_{11} \\ \vdots & \vdots \\ 1 & x_{kn_k} \end{pmatrix}, \quad Z = \begin{pmatrix} Z_1 & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & Z_k \end{pmatrix}, \quad Z_i = \begin{pmatrix} 1 & x_{i1} \\ \vdots & \vdots \\ 1 & x_{in_i} \end{pmatrix},$$

$$\beta = \begin{pmatrix} \beta_0 \\ \beta_1 \end{pmatrix}, \quad b = \begin{pmatrix} b_1 \\ \vdots \\ b_k \end{pmatrix}, \quad b_i = \begin{pmatrix} b_{0i} \\ b_{1i} \end{pmatrix}, \quad \Sigma_b = \begin{pmatrix} \Sigma_b^* & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & \Sigma_b^* \end{pmatrix}$$

where  $\Sigma_b^*$  is an unspecified  $2 \times 2$  positive definite matrix.

## Variance components

The term **mixed model** refers to the fact that the linear predictor  $X\beta + Zb$  contains both fixed effects  $\beta$  and random effects  $b$ .

Under an LMM, we can write the marginal distribution of  $Y$  directly as

$$Y \sim N(X\beta, \Sigma + Z\Sigma_b Z^T) \quad (13)$$

where  $X$  and  $Z$  are matrices containing values of explanatory variables.

Hence  $\text{var}(Y)$  is comprised of two **variance components**.

Other ways of describing LMMs for clustered data, such as (12) (and their generalised linear model counterparts) are as **hierarchical** models or **multilevel** models. This reflects the two-stage structure of the model, a conditional model for  $Y_{ij} | b_i$ , followed by a marginal model for the random effects  $b_i$ .

Sometimes the hierarchy can have further levels, corresponding to clusters nested within clusters, for example, patients within wards within hospitals, or pupils within classes within schools.

APTS: Statistical Modelling

April 2010 – slide 107

## Discussion: Why random effects?

It would be perfectly possible to take a model such as (12) and ignore the final component, leading to fixed cluster effects (as we did for the rat growth data).

The main issue with such an approach is that inferences, particularly predictive inferences can then only be made about those clusters present in the observed data.

Random effects models, on the other hand, allow inferences to be extended to a wider population (at the expense of a further modelling assumption).

It also can be the case, as in (9) with only one observation per 'cluster', that fixed effects are not identifiable, whereas random effects can still be estimated. Similarly, some treatment variables must be applied at the cluster level, so fixed treatment and cluster effects are aliased.

Finally, random effects allow 'borrowing strength' across clusters by shrinking fixed effects towards a common mean.

APTS: Statistical Modelling

April 2010 – slide 108

## Discussion: A Bayesian perspective

A Bayesian LMM supplements (11) with prior distributions for  $\beta$ ,  $\Sigma$  and  $\Sigma_b$ .

In one sense the distinction between fixed and random effects is much less significant, as in the full Bayesian probability specification, both  $\beta$  and  $b$ , as unknowns have probability distributions,  $f(\beta)$  and  $f(b) = \int f(b | \Sigma_b) f(\Sigma_b) d\Sigma_b$

Indeed, prior distributions for 'fixed' effects are sometimes constructed in a hierarchical fashion, for convenience (for example, heavy-tailed priors are often constructed this way).

The main difference is the possibility that random effects for which we have no relevant data (for example cluster effects for unobserved clusters) might need to be predicted.

APTS: Statistical Modelling

April 2010 – slide 109

## LMM fitting

The likelihood for  $(\beta, \Sigma, \Sigma_b)$  is available directly from (13) as

$$f(y | \beta, \Sigma, \Sigma_b) \propto |V|^{-1/2} \exp\left(\frac{1}{2}(y - X\beta)^T V^{-1}(y - X\beta)\right) \quad (14)$$

where  $V = \Sigma + Z\Sigma_b Z^T$ . This likelihood can be maximised directly (usually numerically).

However, mles for variance parameters of LMMs can have large downward bias (particularly in cluster models with a small number of observed clusters).

Hence estimation by **REML** – *REstricted* (or *REsidual*) Maximum Likelihood is usually preferred.

REML proceeds by estimating the variance parameters  $(\Sigma, \Sigma_b)$  using a *marginal likelihood* based on the residuals from a (generalised) least squares fit of the model  $E(Y) = X\beta$ .

APTS: Statistical Modelling

April 2010 – slide 110

## REML

In effect, REML maximizes the likelihood of any linearly independent sub-vector of  $(I_n - H)y$  where  $H = X(X^T X)^{-1} X^T$  is the usual hat matrix. As

$$(I_n - H)y \sim N(0, (I_n - H)V(I_n - H))$$

this likelihood will be free of  $\beta$ . It can be written in terms of the full likelihood (14) as

$$f(r | \Sigma, \Sigma_b) \propto f(y | \hat{\beta}, \Sigma, \Sigma_b) |X^T V X|^{1/2} \quad (15)$$

where

$$\hat{\beta} = (X^T V^{-1} X)^{-1} X^T V^{-1} y \quad (16)$$

is the usual generalised least squares estimator given known  $V$ .

Having first obtained  $(\hat{\Sigma}, \hat{\Sigma}_b)$  by maximising (15),  $\hat{\beta}$  is obtained by plugging the resulting  $\hat{V}$  into (16).

Note that REML maximised likelihoods cannot be used to compare different fixed effects specifications, due to the dependence of 'data'  $r$  in  $f(r | \Sigma, \Sigma_b)$  on  $X$ .

APTS: Statistical Modelling

April 2010 – slide 111



## Estimating random effects

A natural predictor  $\tilde{b}$  of the random effect vector  $b$  is obtained by minimising the mean squared prediction error  $E[(\tilde{b} - b)^T(\tilde{b} - b)]$  where the expectation is over both  $b$  and  $y$ .

This is achieved by

$$\tilde{b} = E(b | y) = (Z^T \Sigma^{-1} Z + \Sigma_b^{-1})^{-1} Z^T \Sigma^{-1} (y - X\beta) \quad (17)$$

giving the **Best Linear Unbiased Predictor** (BLUP) for  $b$ , with corresponding variance

$$\text{var}(b | y) = (Z^T \Sigma^{-1} Z + \Sigma_b^{-1})^{-1}$$

The estimates are obtained by plugging in  $(\hat{\beta}, \hat{\Sigma}, \hat{\Sigma}_b)$ , and are **shrunk** towards 0, in comparison with equivalent fixed effects estimators.

Any component,  $b_k$  of  $b$  with no relevant data (for example a cluster effect for an as yet unobserved cluster) corresponds to a null column of  $Z$ , and then  $\tilde{b}_k = 0$  and  $\text{var}(b_k | y) = [\Sigma_b]_{kk}$ , which may be estimated if, as is usual,  $b_k$  shares a variance with other random effects.

APTS: Statistical Modelling

April 2010 – slide 112

## Bayesian estimation: the Gibbs sampler

Bayesian estimation in LMMs (and their generalised linear model counterparts) generally proceeds using **Markov Chain Monte Carlo (MCMC)** methods, in particular approaches based on the **Gibbs sampler**. Such methods have proved very effective.

MCMC computation provides posterior summaries, by **generating a dependent** sample from the posterior distribution of interest. Then, any posterior expectation can be estimated by the corresponding Monte Carlo sample mean, densities can be estimated from samples etc.

MCMC will be covered in detail in APTS: Computer Intensive Statistics. Here we simply describe the (most basic) Gibbs sampler.

*To generate from  $f(y_1, \dots, y_n)$ , (where the component  $y_i$ s are allowed to be multivariate) the Gibbs sampler starts from an arbitrary value of  $y$  and updates components (sequentially or otherwise) by generating from the conditional distributions  $f(y_i | y_{\setminus i})$  where  $y_{\setminus i}$  are all the variables other than  $y_i$ , set at their currently generated values.*

Hence, to apply the Gibbs sampler, we require conditional distributions which are available for sampling.

APTS: Statistical Modelling

April 2010 – slide 113

## Bayesian estimation for LMMs

For the LMM

$$Y \sim N(\mu, \Sigma), \quad \mu = X\beta + Zb, \quad b \sim N(0, \Sigma_b)$$

with corresponding prior densities  $f(\beta)$ ,  $f(\Sigma)$ ,  $f(\Sigma_b)$ , we obtain the *conditional* posterior distributions

$$\begin{aligned} f(\beta \mid y, \text{rest}) &\propto \phi(y - Zb; X\beta, \Sigma) f(\beta) \\ f(b \mid y, \text{rest}) &\propto \phi(y - X\beta; Zb, \Sigma) \phi(b; 0, \Sigma_b) \\ f(\Sigma \mid y, \text{rest}) &\propto \phi(y - X\beta - Zb; 0, \Sigma) f(\Sigma) \\ f(\Sigma_b \mid y, \text{rest}) &\propto \phi(b; 0, \Sigma_b) f(\Sigma_b) \end{aligned}$$

where  $\phi(y; \mu, \Sigma)$  is a  $N(\mu, \Sigma)$  p.d.f. evaluated at  $y$ .

We can exploit **conditional conjugacy** in the choices of  $f(\beta)$ ,  $f(\Sigma)$ ,  $f(\Sigma_b)$  making the conditionals above of known form and hence straightforward to sample from. The conditional independence  $(\beta, \Sigma) \perp\!\!\!\perp \Sigma_b \mid b$  is also helpful.

See Practical 3 for further details.

APTS: Statistical Modelling

April 2010 – slide 114

## Example: Rat growth revisited

Here, we consider the model

$$y_{ij} = \beta_0 + b_{0i} + (\beta_1 + b_{1i})x_{ij} + \epsilon_{ij}, \quad (b_{0i}, b_{1i})^T \stackrel{\text{iid}}{\sim} N(0, \Sigma_b)$$

where  $\epsilon_{ij} \stackrel{\text{iid}}{\sim} \mathcal{N}(0, \sigma^2)$  and  $\Sigma_b$  is an unspecified covariance matrix. This model allows for random (cluster specific) slope and intercept.

Estimates obtained by REML (ML in brackets) are

Parameter	Estimate	Standard error
$\beta_0$	156.05	2.16 (2.13)
$\beta_1$	43.27	0.73 (0.72)
$\Sigma_{00}^{1/2} = s.d.(b_0)$	10.93 (10.71)	
$\Sigma_{11}^{1/2} = s.d.(b_1)$	3.53 (3.46)	
$Corr(b_0, b_1)$	0.18 (0.19)	


As expected ML variances are smaller, but not by much.

APTS: Statistical Modelling

April 2010 – slide 115

### Example: Fixed v. random effect estimates

The shrinkage of random effect estimates towards a common mean is clearly illustrated.



Random effects estimates 'borrow strength' across clusters, due to the  $\Sigma_b^{-1}$  term in (17). Extent of this is determined by cluster similarity. This is usually considered to be a desirable behaviour.

APTS: Statistical Modelling

April 2010 – slide 116

### Random effect shrinkage

The following simple example illustrates (from a Bayesian perspective) why and how random effects are shrunk to a common value.

Suppose that  $y_1, \dots, y_n$  satisfy

$$y_j | \theta_j \stackrel{\text{iid}}{\sim} N(\theta_j, v_j), \quad \theta_1, \dots, \theta_n | \mu \stackrel{\text{iid}}{\sim} N(\mu, \sigma^2), \quad \mu \sim N(\mu_0, \tau^2),$$

where  $v_1, \dots, v_n, \sigma^2, \mu_0$  and  $\tau^2$  are assumed known here. Then, the usual posterior calculations give us

$$E(\mu | y) = \frac{\mu_0/\tau^2 + \sum y_j/(\sigma^2 + v_j)}{1/\tau^2 + \sum 1/(\sigma^2 + v_j)}, \quad \text{var}(\mu | y) = \frac{1}{1/\tau^2 + \sum 1/(\sigma^2 + v_j)},$$

and

$$E(\theta_j | y) = (1 - w)E(\mu | y) + wy_j,$$

where


$$w = \frac{\sigma^2}{\sigma^2 + v_j}.$$

APTS: Statistical Modelling

April 2010 – slide 117

### Example: Diagnostics

Normal Q-Q plots of intercept (panel 1) and slope (panel 2) random effects and residuals v. week (panel 3)



Evidence of a common quadratic effect, confirmed by AIC (1036 v. 1099) and BIC (1054 v. 1114) based on full ML fits. AIC would also include a cluster quadratic effect (BIC equivocal).

APTS: Statistical Modelling

April 2010 – slide 118

### Generalised linear mixed models

Generalised linear mixed models (GLMMs) generalise LMMs to non-normal data, in the obvious way:

$$Y_i \stackrel{\text{ind}}{\sim} F(\cdot | \mu_i, \sigma^2), \quad g(\mu) \equiv \begin{pmatrix} g(\mu_1) \\ \vdots \\ g(\mu_n) \end{pmatrix} = X\beta + Zb, \quad b \sim N(0, \Sigma_b) \quad (18)$$

where  $F(\cdot | \mu_i, \sigma^2)$  is an exponential family distribution with  $E(Y) = \mu$  and  $\text{var}(Y) = \sigma^2 V(\mu)/m$  for known  $m$ . Commonly (e.g. Binomial, Poisson)  $\sigma^2 = 1$ , and we shall assume this from here on.

It is not necessary that the distribution for the random effects  $b$  is normal, but this usually fits. It is possible (but beyond the scope of this module) to relax this.

APTS: Statistical Modelling

April 2010 – slide 119

### GLMM example

A plausible GLMM for binary data in  $k$  clusters with  $n_1, \dots, n_k$  observations per cluster, and a single explanatory variable  $x$  (e.g. the toxoplasmosis data at individual level) is

$$Y_{ij} \stackrel{\text{ind}}{\sim} \text{Bernoulli}(\mu_i), \quad \log \frac{\mu_i}{1 - \mu_i} = \beta_0 + b_{0i} + \beta_1 x_{ij}, \quad b_{0i} \stackrel{\text{ind}}{\sim} N(0, \sigma_b^2) \quad (19)$$

[note: no random slope here]. This fits into the general GLMM framework (18) with

$$X = \begin{pmatrix} 1 & x_{11} \\ \vdots & \vdots \\ 1 & x_{kn_k} \end{pmatrix}, \quad Z = \begin{pmatrix} Z_1 & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & Z_k \end{pmatrix}, \quad Z_i = \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix},$$

$$\beta = (\beta_0, \beta_1)^T, \quad b = (b_{01}, \dots, b_{0k})^T, \quad \Sigma_b = \sigma_b^2 I_k$$

[or equivalent binomial representation for city data, with clusters of size 1.]

APTS: Statistical Modelling

April 2010 – slide 120

## GLMM likelihood

The marginal distribution for the observed  $Y$  in a GLMM does not usually have a convenient closed-form representation.

$$\begin{aligned} f(y | \beta, \Sigma_b) &= \int f(y | \beta, b, \Sigma_b) f(b | \beta, \Sigma_b) db \\ &= \int f(y | \beta, b) f(b | \Sigma_b) db \\ &= \int \prod_{i=1}^n f(y_i | g^{-1}([X\beta + Zb]_i)) f(b | \Sigma_b) db. \end{aligned} \quad (20)$$

For **nested** random effects structures, some simplification is possible. For example, for (19)

$$f(y | \beta, \sigma_b^2) \propto \prod_{i=1}^n \int \frac{\exp(\sum_j y_{ij}(\beta_0 + b_{0i} + \beta_1 x_{ij}))}{\{1 + \exp(\sum_j y_{ij}(\beta_0 + b_{0i} + \beta_1 x_{ij}))\}^{n_k}} \phi(b_{0i}; 0, \sigma_b^2) db_{0i}$$

a product of one-dimensional integrals.

APTS: Statistical Modelling

April 2010 – slide 121

## GLMM fitting: quadrature

Fitting a GLMM by likelihood methods requires some method for approximating the integrals involved.

The most reliable when the integrals are of low dimension is to use Gaussian quadrature (see APTS: Statistical computing). For example, for a one-dimensional cluster-level random intercept  $b_i$  we might use

$$\begin{aligned} \int \prod_j f(y_{ij} | g^{-1}(x_i^T \beta + b_i)) \phi(b_i | 0, \sigma_b^2) db_i \\ \approx \sum_{q=1}^Q w_q \prod_j f(y_{ij} | g^{-1}(x_i^T \beta + b_{iq})) \end{aligned}$$

for suitably chosen weights ( $w_q, q = 1, \dots, Q$ ) and quadrature points ( $b_{iq}, q = 1, \dots, Q$ )

Effective quadrature approaches use information about the mode and dispersion of the integrand (can be done adaptively).

For multi-dimensional  $b_i$ , quadrature rules can be applied recursively, but performance (in fixed time) diminishes rapidly with dimension.

APTS: Statistical Modelling

April 2010 – slide 122

### GLMM fitting: Penalised quasi-likelihood

An alternative approach to fitting a GLMM uses penalised quasi-likelihood (PQL).

The most straightforward way of thinking about PQL is to consider the adjusted dependent variable  $v$  constructed when computing mles for a GLM using Fisher scoring

$$v_i = (y_i - \mu_i)g'(\mu_i) + \eta_i$$

Now, for a GLMM,

$$E(v | b) = \eta = X\beta + Zb$$

and

$$\text{var}(v | b) = W^{-1} = \text{diag}(\text{var}(y_i)g'(\mu_i)^2),$$

where  $W$  is the weight matrix used in Fisher scoring.

APTS: Statistical Modelling

April 2010 – slide 123

### GLMM fitting: PQL continued

Hence, approximating the conditional distribution of  $v$  by a normal distribution, we have

$$v \sim N(X\beta + Zb, W^{-1}), \quad b \sim N(0, \Sigma_b) \quad (21)$$

where  $v$  and  $W$  also depend on  $\beta$  and  $b$ .

PQL proceeds by iteratively estimating  $\beta$ ,  $b$  and  $\Sigma_b$  for the linear mixed model (21) for  $v$ , updating  $v$  and  $W$  at each stage, based on the current estimates of  $\beta$  and  $b$ .

An alternative justification for PQL is as using a Laplace-type approximation to the integral in the GLMM likelihood.

A full Laplace approximation (expanding the complete log-integrand, and evaluating the Hessian matrix at the mode) is an alternative, equivalent to one-point Gaussian quadrature.

APTS: Statistical Modelling

April 2010 – slide 124

### GLMM fitting: discussion

Using PQL, estimates of random effects  $b$  come 'for free'. With Gaussian quadrature, some extra effort is required to compute  $E(b | y)$  – further quadrature is an obvious possibility.

There are drawbacks with PQL, and the best advice is to use it with caution.

- It can fail badly when the normal approximation that justifies it is invalid (for example for binary observations)
- As it does not use a full likelihood, model comparison should not be performed using PQL maximised 'likelihoods'

Likelihood inference for GLMMs remains an area of active research and vigorous debate. Recent approaches include HGLMs (hierarchical GLMs) where inference is based on the h-likelihood  $f(y | \beta, b)f(b | \Sigma)$ .

APTS: Statistical Modelling

April 2010 – slide 125

## Bayesian estimation for GLMMs

Bayesian estimation in GLMMs, as in LMMs, is generally based on the Gibbs sampler. For the GLMM

$$Y_i \stackrel{\text{ind}}{\sim} F(\cdot | \mu), \quad g(\mu) = X\beta + Zb, \quad b \sim N(0, \Sigma_b)$$

with corresponding prior densities  $f(\beta)$  and  $f(\Sigma_b)$ , we obtain the *conditional* posterior distributions

$$f(\beta | y, \text{rest}) \propto f(\beta) \prod_i f(y_i | g^{-1}(X\beta + Zb))$$

$$f(b | y, \text{rest}) \propto \phi(b; 0, \Sigma_b) \prod_i f(y_i | g^{-1}(X\beta + Zb))$$

$$f(\Sigma_b | y, \text{rest}) \propto \phi(b; 0, \Sigma_b) f(\Sigma_b)$$

For a conditionally conjugate choice of  $f(\Sigma_b)$ ,  $f(\Sigma_b | y, \text{rest})$  is straightforward to sample from. The conditionals for  $\beta$  and  $b$  are not generally available for direct sampling, but there are a number of ways of modifying the basic approach to account for this.

APTS: Statistical Modelling

April 2010 – slide 126

## Toxoplasmosis revisited

Estimates and standard errors obtained by ML (quadrature), Laplace and PQL for the individual-level model

$$Y_{ij} \stackrel{\text{ind}}{\sim} \text{Bernoulli}(\mu_i), \quad \log \frac{\mu_i}{1 - \mu_i} = \beta_0 + b_{0i} + \beta_1 x_{ij}, \quad b_{0i} \stackrel{\text{ind}}{\sim} N(0, \sigma_b^2)$$

Parameter	Estimate (s.e.)		
	ML	Laplace	PQL
$\beta_0$	-0.1343 (1.440)	-0.1384 (1.488)	-0.150 (1.392)
$\beta_1 (\times 10^6)$	5.930 (745.7)	7.215 (770.2)	-5.711 (721.7)
$\sigma_b$	0.5132	0.5209	0.4911
AIC	65.75	65.96	'65.98'

APTS: Statistical Modelling

April 2010 – slide 127

## Toxoplasmosis continued

Estimates and standard errors obtained by ML (quadrature), Laplace and PQL for the extended model

$$\log \frac{\mu_i}{1 - \mu_i} = \beta_0 + b_{0i} + \beta_1 x_{ij} + \beta_2 x_{ij}^2 + \beta_3 x_{ij}^3.$$

Parameter	Estimate (s.e.)		
	ML	Laplace	PQL
$\beta_0$	-335.5 (136.6)	-335.0 (136.3)	-330.8 (140.7)
$\beta_1$	0.5238 (0.2118)	0.5231 (0.2112)	0.5166 (0.2180)
$\beta_2 (\times 10^4)$	-2.710 (1.089)	-2.706 (1.086)	-2.674 (1.121)
$\beta_3 (\times 10^8)$	4.463 (1.857)	4.636 (1.852)	4.583 (1.910)
$\sigma_b$	0.4232	0.4171	0.4508
AIC	63.84	63.97	'64.03'

So for this example, a good agreement between the different computational methods. Some evidence for the cubic model over the linear model.

APTS: Statistical Modelling

April 2010 – slide 128

## Conditional independence and graphical representations

slide 129

### The role of conditional independence

In modelling clustered data, the requirement is often (as in the toxoplasmosis example above) to construct a model to incorporate both non-normality and dependence. There are rather few 'off-the-shelf' models for dependent observations (and those that do exist, such as the multivariate normal, often require strong assumptions which may be hard to justify in practice).

The 'trick' with GLMMs was to model dependence via a series of **conditionally independent** sub-models for the observations  $y$  given the random effects  $b$ , with dependence induced by marginalising over the distribution of  $b$ .

**De Finetti's theorem** provides some theoretical justification for modelling dependent random variables as conditionally independent given some unknown parameter (which we here denote by  $\phi$ ).

APTS: Statistical Modelling

April 2010 – slide 130



## De Finetti's theorem

De Finetti's theorem states (approximately) that any  $y_1, \dots, y_n$  which can be thought of as a finite subset of an **exchangeable** infinite sequence of random variables  $y_1, y_2, \dots$ , has a joint density which can be written as

$$f(y) = \int f(\phi) \prod_{i=1}^n f(y_i | \phi) d\phi$$

for some  $f(\phi)$ ,  $f(y_i | \phi)$ . Hence the  $y_i$  can be modelled as conditionally independent given  $\phi$ .

An *exchangeable* infinite sequence is one for which any finite subsequence has a distribution which is invariant under permutation of the labels of its components.

We can invoke this as an argument for treating as conditionally independent any set of variables about which our prior belief is symmetric.

APTS: Statistical Modelling

April 2010 – slide 131

## Complex stochastic models

In many applications we want to model a multivariate response and/or to incorporate a complex (crossed or hierarchically nested) cluster structure amongst the observations.

The same general approach, splitting the model up into small components, with a potentially rich conditional independence structure linking them facilitates both model construction and understanding, and (potentially) computation.

APTS: Statistical Modelling

April 2010 – slide 132

## Conditional independence graphs

An extremely useful tool, for model description, model interpretation, and to assist identifying efficient methods for computation is the **directed acyclic graph (DAG)** representing the model.

Denote by  $Y = (Y_1, \dots, Y_\ell)$  the collection of elements of the model which are considered random (given a probability distribution). Then the model is a (parametric) description of the joint distribution  $f(y)$ , which we can decompose as

$$f(y) = f(y_1) f(y_2 | y_1) \cdots f(y_\ell | y_1, \dots, y_{\ell-1}) = \prod_i f(y_i | y_{<i})$$

where  $y_{<i} = \{y_1, \dots, y_{i-1}\}$ . Now, for certain orderings of the variables in  $Y$ , the model may admit conditional independences, exhibited through  $f(y_i | y_1, \dots, y_{i-1})$  being functionally free of  $y_j$  for one or more  $j < i$ . This is expressed as

$$Y_i \perp\!\!\!\perp Y_j | Y_{<i \setminus j}$$

where  $Y_{<i \setminus j} = \{Y_1, \dots, Y_{j-1}, Y_{j+1}, \dots, Y_{i-1}\}$ .

APTS: Statistical Modelling

April 2010 – slide 133

## DAGs


The directed acyclic graph (DAG) representing the probability model, decomposed as

$$f(y) = \prod_i f(y_i | y_{<i})$$

consists of a vertex (or node) for each  $Y_i$ , together with an directed edge (arrow) **to** each  $Y_j$  **from** each  $Y_i, i < j$  such that  $f(y_j | y_{<j})$  depends on  $y_i$ . For example, the model

$$f(y_1, y_2, y_3) = f(y_1)f(y_2 | y_1)f(y_3 | y_2)$$

is represented by the DAG



The conditional independence of  $Y_1$  and  $Y_3$  given  $Y_2$  is represented by the absence of a directed edge from  $Y_1$  to  $Y_3$ .

APTS: Statistical Modelling

April 2010 – slide 134

## DAG for a GLMM


The DAG for the general GLMM

$$Y_i \stackrel{\text{ind}}{\sim} F(\cdot | \mu_i, \sigma^2), \quad g(\mu) = X\beta + Zb, \quad b \sim N(0, \Sigma_b)$$

consists, in its most basic form of two nodes:



It can be informative to include parameters and explanatory data in the DAG. Such fixed (non-stochastic) quantities are often denoted by a different style of vertex.



It may also be helpful to consider the components of  $Y$  as separate vertices.


APTS: Statistical Modelling

April 2010 – slide 135

## DAG for a Bayesian GLMM

A Bayesian model is a full joint probability model, across both the variables treated as stochastic in a classical approach, and any unspecified model parameters. The marginal probability distribution for the parameters represents the prior (to observing data) uncertainty about these quantities.

The appropriate DAG for a Bayesian GLMM reflects this, augmenting the DAG on the previous slide to:



where  $\phi_\sigma$ ,  $\phi_\Sigma$  and  $\phi_\beta$  are *hyperparameters* – fixed inputs into the prior distributions for  $\sigma^2$ ,  $\sigma_b$  and  $\beta$  respectively.

APTS: Statistical Modelling

April 2010 – slide 136

## DAG properties

Suppose we have a DAG representing our model for a collection of random variables  $Y = (Y_1, \dots, Y_\ell)$  where the ordering of the  $Y_i$ s is chosen such that all edges in the DAG are from lower to higher numbered vertices. [This must be possible for an acyclic graph, but there will generally be more than one possible ordering]. Then the joint distribution for  $Y$  factorises as

$$f(y) = \prod_i f(y_i \mid pa[y_i])$$

where  $pa[y_i]$  represents the subset of  $\{y_j, j < i\}$  with edges **to**  $y_i$ . Such variables are called the **parents** of  $y_i$ .

APTS: Statistical Modelling


April 2010 – slide 137

## The local Markov property

A natural consequence of the DAG factorisation of the joint distribution of  $Y$  is the **local Markov property for DAGS**. *This states that any variable  $Y_i$  is conditionally independent of its non-descendants, given its parents.*

A descendent of  $Y_i$  is any variable in  $\{Y_j, j > i\}$  which can be reached in the graph by following a sequence of edges **from**  $Y_i$  (respecting the direction of the edges).

For example, for the simple DAG above




the conditional independence of  $Y_3$  and  $Y_1$  given  $Y_2$  is an immediate consequence of the local Markov property.

APTS: Statistical Modelling

April 2010 – slide 138

## The local Markov property – limitations

Not all useful conditional independence properties of DAG models follow immediately from the local Markov property. For example, for the Bayesian GLMM



the posterior distribution is conditional on observed  $Y$ , for which the local Markov property is unhelpful, as  $Y$  is not a parent of any other variable.

To learn more about conditional independences arising from a DAG, it is necessary to construct the corresponding **undirected conditional independence graph**.

APTS: Statistical Modelling

April 2010 – slide 139

## Undirected graphs

An undirected conditional independence graph for  $Y$  consists of a vertex for each  $Y_i$ , together with a set of undirected edges (lines) between vertices such that **absence** of an edge between two vertices  $Y_i$  and  $Y_j$  implies the conditional independence

$$Y_i \perp\!\!\!\perp Y_j \mid Y_{\setminus\{i,j\}}$$

where  $Y_{\setminus\{i,j\}}$  is the set of variables excluding  $Y_i$  and  $Y_j$ .


From a DAG, we can obtain the corresponding undirected conditional independence graph via a two stage process

- First we *moralise* the graph by adding an (undirected) edge between ('marrying') any two vertices which have a *child* in common, and which are not already joined by an edge.
- Then we replace all directed edges by undirected edges.

APTS: Statistical Modelling

April 2010 – slide 140

### Undirected graphs: examples



APTS: Statistical Modelling

April 2010 – slide 141

### Global Markov property

For an undirected conditional independence graph, the **global Markov property** states that any two variables,  $Y_i$  and  $Y_j$  say, are conditionally independent given any subset  $Y_{\text{sep}}$  of the other variables which *separate*  $Y_i$  and  $Y_j$  in the graph.


We say that  $Y_{\text{sep}}$  separates  $Y_i$  and  $Y_j$  in an undirected graph if any path from  $Y_i$  to  $Y_j$  via edges in the graph must pass through a variable in  $Y_{\text{sep}}$ .

APTS: Statistical Modelling


April 2010 – slide 142

### Undirected graph for Bayesian GLMM

The DAG for the Bayesian GLMM



has corresponding undirected graph (for the stochastic vertices)



The conditional independence of  $(\beta, \sigma^2)$  and  $\Sigma_b$  given  $b$  (and  $Y$ ) is immediately obvious.

APTS: Statistical Modelling

April 2010 – slide 143

## Markov equivalence

Any moral DAG (one which has no 'unmarried' parents) is **Markov equivalent** to its corresponding undirected graph (i.e. it encodes exactly the same conditional independence structure).

Conversely, the vertices of any **decomposable** undirected graph (one with no chordless cycles of four or more vertices) can be numbered so that replacing the undirected edges by directed edges from lower to higher numbered vertices produces a Markov equivalent DAG.

Such a numbering is called a **perfect** numbering for the graph, and is not unique.

It immediately follows that the Markov equivalence classes for DAGs can have (many) more than one member, each of which implies the same model for the data (in terms of conditional independence structure)

The class of DAGs is clearly much larger than the class of undirected graphs, and encompasses a richer range of conditional independence structures.

### 3. Missing Data and Latent Variables

slide 145

#### Overview

1. Missing data
2. Latent variables
3. EM algorithm

APTS: Statistical Modelling

April 2010 – slide 146

### Missing Data

slide 147

#### Example 1: Birthweight and smoking

Data from the Collaborative Perinatal Project

		Birth weight (known)		Birth weight (unknown)	
		< 2500	≥ 2500	< 2500	≥ 2500
Mother smokes? (known)	Y	4512	21009		1049
	N	3394	24132		1135
<hr/>					
Mother smokes? (unknown)	Y				
	N	142	464		1224

APTS: Statistical Modelling

April 2010 – slide 148

#### Example 2: Political opinions

Data extracted from the British General Election Panel Survey

Sex	Social class	Intention known				Intention unknown			
		Con.	Lab.	Lib.	Other	Con.	Lab.	Lib.	Other
M	1	26	8	7	0				11
	2	87	37	30	6				64
	3	66	77	23	8				77
	4	14	25	15	1				12
	5	6	6	2	0				7
F	1	1	1	0	1				2
	2	63	34	32	2				68
	3	102	52	22	4				77
	4	10	32	10	2				38
	5	20	25	8	2				19

1=Professional, 2=Managerial and technical, 3= Skilled, 4=Semi-skilled or unskilled, 5=Never worked.

APTS: Statistical Modelling

April 2010 – slide 149

## Introduction

Missing data arises in many practical applications. Typically, our data might appear (with missing observations indicated by \*) as

Unit ( $i$ )	Variable ( $j$ )				
	1	2	3	...	$p$
1	$y_{11}$	$y_{12}$	$y_{13}$	...	$y_{1p}$
2	$y_{21}$	*	$y_{23}$	...	$y_{2p}$
3	*	$y_{32}$	*	...	*
4	*	$y_{42}$	$y_{43}$	...	$y_{4p}$
...	...	...	...	...	...
$n$	$y_{n1}$	*	$y_{n3}$	...	$y_{np}$

Variables which have missing observations in our data frame are said to be subject to **item nonresponse**. When there are units with no available data whatsoever, that is referred to as **unit nonresponse**.

If the variables can be ordered so that, for any unit  $i$ , ( $y_{ij}$  is missing)  $\Rightarrow$  ( $y_{ik}$  is missing for all  $k > j$ ), the missing data pattern is said to be **monotone** (e.g. longitudinal dropout, special cases like Example 2).

APTS: Statistical Modelling

April 2010 – slide 150

## Issues

Missing data creates two major problems for analysis.

- Suppose that we have a model  $f(y | \theta)$ , for which the likelihood is tractable. When certain  $y_{ij}$  are missing, the likelihood for inference must be based on the observed data distribution

$$f(y_{\text{obs}} | \theta) = \int f(y_{\text{obs}}, y_{\text{mis}} | \theta) dy_{\text{mis}} \quad (22)$$

where the subscripts obs and mis refer to observed and missing components, respectively. It is typically much more difficult to compute (22) than  $f(y | \theta)$  for fully observed data.

- Even when it can be computed, the likelihood (22) is only valid for inference about  $\theta$  under the assumption that the fact that certain observations are missing provides no information about  $\theta$ .

APTS: Statistical Modelling

April 2010 – slide 151



## Models

To formalise this, it is helpful to introduce a series of binary response indicator variables  $r_1, \dots, r_p$ , where

$$r_{ij} = 1 \Leftrightarrow y_{ij} \text{ is observed, } i = 1, \dots, n; j = 1, \dots, p.$$

We factorise the joint distribution of  $(y, r)$  into a data model for  $y$  and a (conditional) response model for  $r$

$$f(y, r | \theta, \phi) = f(y | \theta)f(r | y, \phi).$$

Then the likelihood for the observed data,  $(y_{\text{obs}}, r)$  is

$$f(y_{\text{obs}}, r | \theta, \phi) = \int f(y_{\text{obs}}, y_{\text{mis}} | \theta)f(r | y_{\text{obs}}, y_{\text{mis}}, \phi)dy_{\text{mis}}. \quad (23)$$

In this set-up inference for  $\theta$  should be based on (23), but there are situations when it is valid to **ignore** the missing data mechanism (and the corresponding variable  $r$ ) and base inference for  $\theta$  on the simpler  $f(y_{\text{obs}} | \theta)$ .

APTS: Statistical Modelling

April 2010 – slide 152

## Ignorability

If  $R \perp\!\!\!\perp Y_{\text{mis}} | Y_{\text{obs}}, \phi$  then  $f(r | y_{\text{obs}}, y_{\text{mis}}, \phi)$  in (23) can be replaced by  $f(r | y_{\text{obs}}, \phi)$ , and (23) is simplified to:

$$f(y_{\text{obs}}, r | \theta, \phi) = f(y_{\text{obs}} | \theta)f(r | y_{\text{obs}}, \phi). \quad (24)$$

Hence, the likelihood for  $(\theta, \phi)$  factorises and *provided that  $\theta$  and  $\phi$  are independent* (in a functional sense for likelihood analysis, and in the usual stochastic sense for Bayesian analysis) inference for  $\theta$  can be based on  $f(y_{\text{obs}} | \theta)$ .

- Any missing data model which satisfies the two requirements above, namely  $[R \perp\!\!\!\perp Y_{\text{mis}} | Y_{\text{obs}}, \phi]$  and  $[\phi \text{ independent of } \theta]$  is said to be **ignorable**. Otherwise, it is nonignorable.
- Missing data which satisfies  $R \perp\!\!\!\perp Y_{\text{mis}} | Y_{\text{obs}}, \phi$  is said to be **missing at random (MAR)**
- Missing data which satisfies the stronger condition  $R \perp\!\!\!\perp Y_{\text{mis}}, Y_{\text{obs}} | \phi$  is said to be **missing completely at random (MCAR)**. For MCAR data, correct (but potentially highly sub-optimal) inferences can be obtained by *complete case analysis*.

APTS: Statistical Modelling

April 2010 – slide 153

## Inference under ignorability

For monotone missing data patterns, it may be possible to deal with  $f(y_{\text{obs}} | \theta)$  directly. For example, suppose that the  $Y_i = (Y_{i1}, \dots, Y_{ip})$  are conditionally independent given  $\theta$ , and furthermore that

$$f(y_i | \theta) = \prod_j f(y_{ij} | y_{i,<j}, \theta_j)$$

where  $\theta = (\theta_1, \dots, \theta_p)$  is a partition into distinct components. Then

$$f(y_{\text{obs}} | \theta) = \prod_i f(y_{i,\text{obs}} | \theta) = \prod_i \prod_{j=1}^{k_i} f(y_{ij} | y_{i,<j}, \theta_j)$$

where  $k_i$  is the 'last' observed variable for unit  $i$ . Hence the likelihood for  $\theta$  factorises into individual components.

Otherwise, methods for inference in the presence of an ignorable missing data mechanism typically exploit the fact the full data analysis, based on  $f(y | \theta)$  is tractable (assuming that it is!)

APTS: Statistical Modelling

April 2010 – slide 154

## Gibbs sampler

For Bayesian analysis, this typically involves generating a sequence of values  $\{\theta^t, y_{\text{mis}}^t, t = 1, \dots\}$  from the joint posterior distribution  $f(\theta, y_{\text{mis}} | y_{\text{obs}})$  using a Gibbs sampler iteratively sampling

- the model-based conditional for  $Y_{\text{mis}} | \theta, y_{\text{obs}}$ .
- the complete data posterior conditional for  $\theta | Y_{\text{mis}}, y_{\text{obs}}$

Often, both of these are convenient for sampling.

The subsample  $\{\theta^t, t = 1, \dots\}$  may then be considered as being drawn from the marginal posterior for  $\theta | y_{\text{obs}}$ , as required.

This is sometimes referred to as **data augmentation**.

APTS: Statistical Modelling

April 2010 – slide 155

## EM algorithm

For maximum likelihood, it is often the case that a corresponding iterative algorithm can be constructed by taking the Gibbs sampler steps above and replacing generation from conditionals with (i) taking expectation (for  $Y_{\text{mis}}$ ) and (ii) likelihood maximisation (for  $\theta$ ), respectively.

- for the current  $\theta^t$  construct the expected log-likelihood  $E[\log f(Y_{\text{mis}}, y_{\text{obs}} | \theta) | y_{\text{obs}}, \theta^t]$
- maximise this expected log-likelihood w.r.t.  $\theta$  to obtain  $\theta^{t+1}$

This is the EM algorithm, of which more details will be presented shortly. The maximisation (M) step is generally straightforward, and for many models, so is the expectation (E) step.

APTS: Statistical Modelling

April 2010 – slide 156

## Nonignorable models

If considered appropriate, then a nonignorable missing data mechanism can be incorporated in  $f(y, r | \theta, \phi)$ . A **selection model** utilises the decomposition

$$f(y, r | \theta, \phi) = f(y | \theta)f(r | y, \phi).$$

where a nonignorable model incorporates dependence of  $R$  on  $Y_{\text{mis}}$ .

Alternatively, a **pattern mixture model** decomposes  $f(y, r | \theta, \phi)$  as

$$f(y, r | \theta, \phi) = f(y | r, \theta)f(r | \phi).$$

Pattern mixture models tend to be less intuitively appealing, but may be easier to analyse (particularly for monotone missing data patterns).

Under either specification, inference must be based on the observed data likelihood

$$f(y_{\text{obs}}, r | \theta, \phi) = \int f(y_{\text{obs}}, y_{\text{mis}}, r | \theta, \phi) dy_{\text{mis}}.$$

Gibbs sampling or EM can be used for computation, but convergence may be slow.

APTS: Statistical Modelling


April 2010 – slide 157

## A simple selection model

Consider the selection model

$$Y \sim N(\theta_1, \theta_2), \quad \Pr(R = 1 | Y = y) = \frac{\exp(\phi_0 + \phi_1 y)}{1 + \exp(\phi_0 + \phi_1 y)}$$

An example of  $f(y | r = 1)$ , the marginal density for  $y_{\text{obs}}$  is



for  $(\theta_1, \theta_2, \phi_0, \phi_1) = (0, 1, 0, 2)$ .

The **selection effect** is quite subtle and will clearly be hard to estimate accurately.

APTS: Statistical Modelling

April 2010 – slide 158

## Nonignorable model issues

In the previous example, it will be impossible to distinguish, on the basis of observed data only, between the proposed selection model, and an ignorable model where the population distribution of  $y$  is naturally slightly skewed.

Generally, nonignorable model inferences are sensitive to model assumptions, and there exist alternative models which cannot be effectively compared on the basis of fit to observed data alone.

Furthermore, inferences from alternative, equally well-fitting models may be very different, as the following (artificial) example illustrates.

$y_1$	$y_2$ (Observed)		$y_2$ (Missing)		
	A	B	A	B	
1	6	18			16
2	3	9			8
3	3	27			10

APTS: Statistical Modelling

April 2010 – slide 159

## Sensitivity example

Missing data estimates based on the ignorable model  $R_2 \perp\!\!\!\perp Y_2 \mid Y_1$

$y_1$	$y_2$ (Observed)		$y_2$ (Missing)		
	A	B	A	B	
1	6	18	<b>4</b>	<b>12</b>	
2	3	9	<b>2</b>	<b>6</b>	
3	3	27	<b>1</b>	<b>9</b>	

Missing data estimates based on the nonignorable model  $R_2 \perp\!\!\!\perp Y_1 \mid Y_2$

$y_1$	$y_2$ (Observed)		$y_2$ (Missing)		
	A	B	A	B	
1	6	18	<b>14</b>	<b>2</b>	
2	3	9	<b>7</b>	<b>1</b>	
3	3	27	<b>7</b>	<b>3</b>	

Potentially very different inferences for the marginal distribution of  $y_2$ .

Pragmatic approaches are based on investigating sensitivity to a range of missing data assumptions.

APTS: Statistical Modelling

April 2010 – slide 160

**Basic idea**


- Many statistical models simplify when written in terms of unobserved **latent variable**  $U$  in addition to the observed data  $Y$ . The latent variable
  - may really exist, for example, when  $Y = I(U > c)$  for some continuous  $U$  ('do you earn less than £ $c$  per year?');
  - may be imaginary—something called IQ is said to underlie scores on intelligence tests, but is IQ just a cultural construct? ("Mismeasure of man" debate ...);
  - may just be a mathematical/computational device (e.g. in MCMC or EM algorithms).
- Examples include random effects models, use of hidden variables in probit regression, mixture models.

**Galaxy data**

Velocities (km/second) of 82 galaxies in a survey of the Corona Borealis region. The error is thought to be less than 50 km/second.

9172	9350	9483	9558	9775	10227	10406	16084	16170	18419
18552	18600	18927	19052	19070	19330	19343	19349	19440	19473
19529	19541	19547	19663	19846	19856	19863	19914	19918	19973
19989	20166	20175	20179	20196	20215	20221	20415	20629	20795
20821	20846	20875	20986	21137	21492	21701	21814	21921	21960
22185	22209	22242	22249	22314	22374	22495	22746	22747	22888
22914	23206	23241	23263	23484	23538	23542	23666	23706	23711
24129	24285	24289	24366	24717	24990	25633	26960	26995	32065
32789	34279								

**Galaxy data**



## Mixture density

- Natural model for such data is a  $p$ -component mixture density

$$f(y; \theta) = \sum_{r=1}^p \pi_r f_r(y; \theta), \quad 0 \leq \pi_r \leq 1, \quad \sum_{r=1}^p \pi_r = 1,$$

where  $\pi_r$  is the probability that  $Y$  comes from the  $r$ th component and  $f_r(y; \theta)$  is its density conditional on this event.

- Can represent this using indicator variables  $U$  taking a value in  $1, \dots, p$  with probabilities  $\pi_1, \dots, \pi_p$  and indicating from which component  $Y$  is drawn.
- Widely used class of models, often with number of components  $p$  unknown.
- Aside: such models are non-regular for likelihood inference:
  - non-identifiable under permutation of components;
  - setting  $\pi_r = 0$  eliminates parameters of  $f_r$ ;
  - maximum of likelihood can be  $+\infty$ , achieved for several  $\theta$

APTS: Statistical Modelling

April 2010 – slide 165

## Other latent variable models

- Let  $[U], D$  denote discrete random variables, and  $(U), X$  continuous ones. Then in notation for graphical models:
  - $[U] \rightarrow X$  or  $[U] \rightarrow D$  denotes finite mixture models, hidden Markov models, changepoint models, etc.;
  - $(U) \rightarrow D$  denotes data coarsening (censoring, truncation, ...);
  - $(U) \rightarrow X$  or  $(U) \rightarrow D$  denotes variance components and other hierarchical models.
- Binary regression:  $U \sim \mathcal{N}(x^T \beta, 1)$  and observed response  $Y = I(U \geq 0)$ , gives probit regression model, log likelihood contribution

$$Y \log \Phi(x^T \beta) + (1 - Y) \log \{1 - \Phi(x^T \beta)\},$$

and similarly if different continuous distribution is chosen for  $U$  (logistic, extreme-value, ...).

APTS: Statistical Modelling

April 2010 – slide 166

**EM algorithm**

- Aim to use observed value  $y$  of  $Y$  for inference on  $\theta$  when we cannot easily compute

$$f(y; \theta) = \int f(y | u; \theta) f(u; \theta) du$$

- The **complete-data log likelihood**

$$\log f(y, u; \theta) = \log f(y; \theta) + \log f(u | y; \theta), \quad (25)$$

is based on  $(U, Y)$ , whereas the **observed-data log likelihood** is

$$\ell(\theta) = \log f(y; \theta).$$

- Take expectation in (25) with respect to  $f(u | y; \theta')$  to get

$$E \{ \log f(Y, U; \theta) | Y = y; \theta' \} = \ell(\theta) + E \{ \log f(U | Y; \theta) | Y = y; \theta' \}, \quad (26)$$

or equivalently  $Q(\theta; \theta') = \ell(\theta) + C(\theta; \theta')$ .

APTS: Statistical Modelling

April 2010 – slide 168

**EM algorithm II**

Fix  $\theta'$  and consider how  $Q(\theta; \theta')$  and  $C(\theta; \theta')$  depend on  $\theta$ .

- Note that  $C(\theta'; \theta') \geq C(\theta; \theta')$ , with equality only when  $\theta = \theta'$  (Jensen's inequality).
- Thus

$$Q(\theta; \theta') \geq Q(\theta'; \theta') \text{ implies } \ell(\theta) - \ell(\theta') \geq C(\theta'; \theta') - C(\theta; \theta') \geq 0. \quad (27)$$

- Under mild smoothness conditions,  $C(\theta; \theta')$  has a stationary point at  $\theta = \theta'$ , so if  $Q(\theta; \theta')$  is stationary at  $\theta = \theta'$ , so too is  $\ell(\theta)$ .

- Hence **EM algorithm**: starting from an initial value  $\theta'$  of  $\theta$ ,

1. compute  $Q(\theta; \theta') = E \{ \log f(Y, U; \theta) | Y = y; \theta' \}$ ; then
2. with  $\theta'$  fixed, maximize  $Q(\theta; \theta')$  over  $\theta$ , giving  $\theta^\dagger$ , say; and
3. check if the algorithm has converged, using  $\ell(\theta^\dagger) - \ell(\theta')$  if available, or  $|\theta^\dagger - \theta'|$ , or both. If not, set  $\theta' = \theta^\dagger$  and go to 1.

Steps 1 and 2 are the **expectation (E)** and **maximization (M)** steps.

- The M-step ensures that  $Q(\theta^\dagger; \theta') \geq Q(\theta'; \theta')$ , so (27) implies that  $\ell(\theta^\dagger) \geq \ell(\theta')$ : the log likelihood never decreases.

APTS: Statistical Modelling

April 2010 – slide 169

## Convergence

- If  $\ell(\theta)$  has
  - only one stationary point, and if  $Q(\theta; \theta')$  eventually reaches a stationary value at  $\hat{\theta}$ , then  $\hat{\theta}$  must maximize  $\ell(\theta)$ ;
  - otherwise the algorithm may converge to a local maximum of the log likelihood or to a turning point.
- The EM algorithm never decreases the log likelihood so is more stable than Newton–Raphson-type algorithms.
- Rate of convergence depends on closeness of  $Q(\theta; \theta')$  and  $\ell(\theta)$ :

$$-\frac{\partial^2 \ell(\theta)}{\partial \theta \partial \theta^T} = E \left\{ -\frac{\partial^2 \log f(y, U; \theta)}{\partial \theta \partial \theta^T} \middle| Y = y; \theta \right\} - E \left\{ -\frac{\partial^2 \log f(U | y; \theta)}{\partial \theta \partial \theta^T} \middle| Y = y; \theta \right\},$$

or  $J(\theta) = I_c(\theta; y) - I_m(\theta; y)$ , giving the **missing information principle**: the observed information equals the complete-data information minus the missing information.


- Rate of convergence slow if largest eigenvalue of  $I_c(\theta; y)^{-1} I_m(\theta; y) \approx 1$ ; this occurs if the missing information is a high proportion of the total.

APTS: Statistical Modelling

April 2010 – slide 170

## (Toy) Example: Negative binomial model

Conditional on  $U = u$ ,  $Y \sim \text{Pois}(u)$  and  $U$  is gamma with mean  $\theta$  and variance  $\theta^2/\nu$ . Suppose  $\nu > 0$  known and make inference for  $\theta$ .



EM algorithm for negative binomial example. Left panel: observed-data log likelihood  $\ell(\theta)$  (solid) and functions  $Q(\theta; \theta')$  for  $\theta' = 1.5, 1.347$  and  $1.028$  (dots, from right). The blobs show the values of  $\theta$  that maximize these functions, which correspond to the first, fifth and fortieth iterations of the EM algorithm. Right: convergence of EM algorithm (dots) and Newton–Raphson algorithm (solid). The panel shows how successive EM iterations update  $\theta'$  and  $\hat{\theta}$ . Notice that the EM iterates always increase  $\ell(\theta)$ , while the Newton–Raphson steps do not.

APTS: Statistical Modelling

April 2010 – slide 171



### Example: Mixture model

- Consider earlier  $p$ -component mixture density  $f(y; \theta) = \sum_{r=1}^p \pi_r f_r(y; \theta)$ , for which likelihood contribution from  $(y, u)$  would be  $\prod_r \{f_r(y; \theta) \pi_r\}^{I(u=r)}$ , giving contribution

$$\log f(y, u; \theta) = \sum_{r=1}^p I(u = r) \{\log \pi_r + \log f_r(y; \theta)\}$$

to the complete-data log likelihood.

- Must compute the expectation of  $\log f(y, u; \theta)$  over

$$w_r(y; \theta') = \Pr(U = r \mid Y = y; \theta') = \frac{\pi_r f_r(y; \theta')}{\sum_{s=1}^p \pi_s f_s(y; \theta')}, \quad r = 1, \dots, p, \quad (28)$$

the weight attributable to component  $r$  if  $y$  has been observed.


- The expected value of  $I(U = r)$  with respect to (28) is  $w_r(y; \theta')$ , so the expected value of the log likelihood based on a random sample  $(y_1, u_1), \dots, (y_n, u_n)$  is

$$\begin{aligned} Q(\theta; \theta') &= \sum_{j=1}^n \sum_{r=1}^p w_r(y_j; \theta') \{\log \pi_r + \log f_r(y_j; \theta)\} \\ &= \sum_{r=1}^p \left\{ \sum_{j=1}^n w_r(y_j; \theta') \right\} \log \pi_r + \sum_{r=1}^p \sum_{j=1}^n w_r(y_j; \theta') \log f_r(y_j; \theta). \end{aligned}$$


### Example: Galaxy data

$p$	1	2	3	4	5
$\hat{\ell}$	-240.42	-220.19	-203.48	-202.52	-192.42

Fitted mixture model with  $p = 4$  normal components:



## Galaxy data



AIC and BIC for the normal mixture models fitted to the galaxy data. BIC is minimised for  $p = 3$  components, and AIC for  $p = 5$  components.

APTS: Statistical Modelling

April 2010 – slide 174

## Exponential family

- Suppose the complete-data log likelihood is from an exponential family:

$$\log f(y, u; \theta) = s(y, u)^T \theta - \kappa(\theta) + c(y, u).$$

- For EM algorithm, need expected value of  $\log f(y, u; \theta)$  with respect to  $f(u | y; \theta')$ . Final term can be ignored, so M-step involves maximizing

$$Q(\theta; \theta') = E \{s(y, U)^T \theta | Y = y; \theta'\} - \kappa(\theta),$$

or equivalently solving for  $\theta$  the equation

$$E \{s(y, U) | Y = y; \theta'\} = \frac{d\kappa(\theta)}{d\theta}.$$

- Likelihood equation for  $\theta$  based on the complete data is  $s(y, u) = d\kappa(\theta)/d\theta$ , so the EM algorithm replaces  $s(y, u)$  by its conditional expectation  $E \{s(y, U) | Y = y; \theta'\}$  and solves the likelihood equation. Thus a routine to fit the complete-data model can readily be adapted for missing data if the conditional expectations are available.

APTS: Statistical Modelling

April 2010 – slide 175

## Comments

- Often E-step requires numerical approximation:
  - simulation from conditional distribution of  $U$  given  $Y$ ;
  - importance sampling;
  - Markov chain algorithm;
- M-step can be performed using Newton–Raphson or similar algorithm, using first and second loglikelihood derivatives (exercise)—may need to be performed in parts, rather than overall
- Can obtain standard errors using these derivatives (exercise)
- In Bayesian analysis, may often be helpful to include latent variables, either
  - because they have useful interpretation in terms of model—*all* parameters are hidden variables, because unobservable in practice
  - to simplify MCMC algorithm—Gibbs sampler is 'Bayesian equivalent' of EM algorithm (exercise)