

1. INVERSE ROOTS

Suppose that $Q(x) = ax^2 + bx + c$ satisfies $ac \neq 0$ and has roots (i.e. solutions of $Q(x) = 0$) α and β .

Show that the quadratic $\tilde{Q}(x) = cx^2 + bx + a$ has roots $\frac{1}{\alpha}$ and $\frac{1}{\beta}$.

Extensions

(1) Show that if $\alpha_1, \dots, \alpha_n$ are the roots of the polynomial P , where

$$P(x) = a_0x^n + \dots + a_{n-1}x + a_n \text{ with } a_0a_n \neq 0,$$

then the roots of \tilde{P} given by

$$\tilde{P}(x) = a_nx^n + \dots + a_0$$

are $\frac{1}{\alpha_1}, \dots, \frac{1}{\alpha_n}$.

(2) Show that the roots of

$$P_e(x) = a_0x^{2n} + a_1x^{2n-1} + \dots + a_{n-1}x^{n+1} + a_nx^n + a_{n-1}x^{n-1} + a_{n-2}x^{n-2} + \dots + a_0 \quad (a_0 \neq 0)$$

are of the form $\alpha_1, \dots, \alpha_n, \frac{1}{\alpha_1}, \dots, \frac{1}{\alpha_n}$.

(3) What can you say about the roots of

$$P_+(x) = a_0x^{2n+1} + a_1x^{2n} + \dots + a_nx^{n+1} + a_nx^n + a_{n-1}x^{n-1} + \dots + a_0$$

and the roots of

$$P_-(x) = a_0x^{2n+1} + a_1x^{2n} + \dots + a_nx^{n+1} - a_nx^n - a_{n-1}x^{n-1} - \dots - a_0?$$

2. TRIGONOMETRIC POLYNOMIALS

The angle sum formula tells us that

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta = 2 \cos^2 \theta - 1.$$

Find a similar expression involving powers of $\cos \theta$ for $\cos 3\theta$.

Extensions

(1) Find the roots of $4\sqrt{2}x^3 - 3\sqrt{2}x = 1$.

(2) What is $\cos \frac{\pi}{12}$?