

# Lecture Notes 7: Dynamic Equations

## Part B: Second and Higher-Order Linear Difference Equations in One Variable

Peter J. Hammond

Latest revision 2020 September 24th,  
typeset from `dynEqLects23B.tex`

# Lecture Outline

## Solving Second-Order Equations

Inhomogeneous Equations

Particular Solutions in Two Special Cases

Method of Undetermined Coefficients

Higher-Order Linear Equations with Constant Coefficients

Stationary States and Stability for Second-Order Equations

## Second-Order Equations

A general **second-order difference equation** specifies the state  $x_t$  at each time  $t$  as a function  $x_t = F_t(x_{t-1}, x_{t-2})$  of the state at **two** previous times.

Suppose we define a new variable defined by  $y_t := x_{t-1}$ . Then the equation  $x_t = F_t(x_{t-1}, x_{t-2})$  can be converted into the coupled pair

$$\begin{aligned}x_t &= F_t(x_{t-1}, y_{t-1}) \\y_t &= x_{t-1}\end{aligned}$$

of **first**-order equations that express the vector  $(x_t, y_t)^\top \in \mathbb{R}^2$  as a function of the vector  $(x_{t-1}, y_{t-1})^\top \in \mathbb{R}^2$ .

## The Linear Case

We focus on linear equations in one variable with constant coefficients, which take the form

$$x_{t+1} + ax_t + bx_{t-1} = f_t$$

Here  $a, b$  are scalars, and  $f_t$  is the forcing term.

We assume that  $b \neq 0$  because otherwise we have the first-order equation  $x_{t+1} + ax_t = f_t$ .

If we define  $y_t = x_{t-1}$ , the equation becomes the coupled pair

$$x_{t+1} = -ax_t - by_t + f_t; \quad y_{t+1} = x_t$$

In matrix form, these can be written as

$$\begin{pmatrix} x_{t+1} \\ y_{t+1} \end{pmatrix} = \begin{pmatrix} -a & -b \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x_t \\ y_t \end{pmatrix} + \begin{pmatrix} f_t \\ 0 \end{pmatrix}$$

Such **vector difference equations** are the subject of part C.

## The Homogeneous Case

Nevertheless, consider the homogeneous case when the vector equation is

$$\begin{pmatrix} x_{t+1} \\ y_{t+1} \end{pmatrix} = \begin{pmatrix} -a & -b \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x_t \\ y_t \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

The solution in matrix form is evidently

$$\begin{pmatrix} x_t \\ y_t \end{pmatrix} = \begin{pmatrix} -a & -b \\ 1 & 0 \end{pmatrix}^t \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$$

for an arbitrary initial state  $\begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$ .

Inspired by our earlier discussion of matrix powers, consider the case when  $(\lambda, (x_0, y_0)^\top)$  is an eigenpair, that is

$$\begin{pmatrix} -a & -b \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \lambda \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \quad \text{where} \quad \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \neq \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

## Solving the Homogeneous Case

In case  $\begin{pmatrix} -a & -b \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \lambda \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$ , the solution takes the form

$$\begin{pmatrix} x_t \\ y_t \end{pmatrix} = \begin{pmatrix} -a & -b \\ 1 & 0 \end{pmatrix}^t \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \lambda^t \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$$

For this to work, the initial vector  $\begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$  must solve

the matrix equation  $\begin{pmatrix} -a - \lambda & -b \\ 1 & -\lambda \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ .

For a non-trivial solution to exist,

the matrix  $\begin{pmatrix} -a - \lambda & -b \\ 1 & -\lambda \end{pmatrix}$  must be singular, implying that

$$\begin{vmatrix} -a - \lambda & -b \\ 1 & -\lambda \end{vmatrix} = \lambda^2 + a\lambda + b = 0$$

## The Auxiliary Equation

Instead of treating the second-order equation as a coupled pair, consider directly the homogeneous second-order equation

$$x_{t+1} + ax_t + bx_{t-1} = 0$$

Inspired by our previous analysis using eigenvalues of a suitable matrix, we look for a solution of the form  $x_t = \lambda^t x_0$ , for suitable constants  $\lambda$  and  $x_0$ .

It is a solution provided that  $\lambda^{t+1}x_0 + a\lambda^t x_0 + b\lambda^{t-1}x_0 = 0$ .

Ignoring the trivial solutions when  $x_0 = 0$  or  $\lambda = 0$ , cancel  $\lambda^{t-1}x_0$  to obtain the **auxiliary** or **characteristic equation**

$$\lambda^2 + a\lambda + b = 0$$

This, of course, is the condition for  $\lambda$  to be an eigenvalue.

## The Auxiliary Equation and Its Roots

The auxiliary equation  $\lambda^2 + a\lambda + b = 0$  is quadratic.

It therefore has two roots  $\lambda_1, \lambda_2$   
satisfying  $\lambda^2 + a\lambda + b = (\lambda - \lambda_1)(\lambda - \lambda_2)$ .

In particular  $\lambda_1 + \lambda_2 = -a$  and  $\lambda_1\lambda_2 = b$ .

The assumption that  $b \neq 0$  implies  
that the two roots  $\lambda_1, \lambda_2$  are both non-zero.

This leaves three cases:

1. two distinct real roots  $\lambda_1, \lambda_2 \in \mathbb{R}$ ,  
which is true iff  $a^2 > 4b$ ;
2. two complex conjugate roots  $\lambda_1, \lambda_2 = re^{\pm i\theta} \in \mathbb{C}$ ,  
which is true iff  $a^2 < 4b$ ;
3. two coincident real roots  $\lambda = \lambda_1 = \lambda_2 \in \mathbb{R}$ ,  
which is true iff  $a^2 = 4b$ .



## Case 1: Two Distinct Real Roots

In this case  $\lambda^2 + a\lambda + b = (\lambda - \lambda_1)(\lambda - \lambda_2)$ ,

where  $\lambda_1, \lambda_2 = -\frac{1}{2}a \pm \frac{1}{2}\sqrt{a^2 - 4b}$ .

Note that  $a = \lambda_1 + \lambda_2$  and  $b = \lambda_1\lambda_2$

with  $a^2 - 4b = (\lambda_1 + \lambda_2)^2 - 4\lambda_1\lambda_2 = (\lambda_1 - \lambda_2)^2 > 0$ .

There are two degrees of freedom in the difference equation, so we look for two **linearly independent** solutions  $x_t^{H(1)}$  and  $x_t^{H(2)}$  of the homogeneous difference equation  $x_{t+1} + ax_t + bx_{t-1} = 0$ .

— that is two solutions for which  $Ax_t^{H(1)} + Bx_t^{H(2)} \equiv 0$  implies that the two scalars  $A$  and  $B$  satisfy  $A = B = 0$ .

## Two Linearly Independent Solutions

Note that  $A\lambda_1^t + B\lambda_2^t = 0$  for both  $t = 0$  and  $t = 1$  if and only if

$$\begin{pmatrix} 1 & 1 \\ \lambda_1 & \lambda_2 \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

This has a non-trivial solution in the two constants  $A$  and  $B$  iff  $0 = \begin{vmatrix} 1 & 1 \\ \lambda_1 & \lambda_2 \end{vmatrix}$ , or if and only if  $0 = \lambda_2 - \lambda_1$ .

So when  $\lambda_1 \neq \lambda_2$ , the only solution is trivial, with  $A = B = 0$ .

Hence, the two functions  $x_t^{(1)} = x_0\lambda_1^t$  and  $x_t^{(2)} = x_0\lambda_2^t$  with  $x_0 \neq 0$  are linearly independent solutions of  $x_{t+1} + ax_t + bx_{t-1} = 0$ .

There are two degrees of freedom in the difference equation.

Therefore, its general solution with these two degrees of freedom is  $x_t = A\lambda_1^t + B\lambda_2^t$  for arbitrary real constants  $A$  and  $B$ .

## Example: The Fibonacci Sequence

The **Fibonacci sequence** is

$$(x_t)_{t=0}^{\infty} = (0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, 233, \dots)$$

It is the unique solution with  $x_0 = 0$  and  $x_1 = 1$  of the **Fibonacci difference equation**  $x_{t+1} - x_t - x_{t-1} = 0$ .

The characteristic equation is  $\lambda^2 - \lambda - 1 = 0$ , with characteristic roots  $\lambda_{1,2} = -\frac{1}{2}(-1 \pm \sqrt{5})$ .

Its two roots are:

(i) the **golden ratio**  $\varphi := \lambda_1 = \frac{1}{2}(1 + \sqrt{5}) \approx 1.61803398875$ ;  
and (ii)  $\lambda_2 = 1 - \lambda_1 = \frac{1}{2}(1 - \sqrt{5}) \approx -0.61803398875$ .

The general solution of the Fibonacci difference equation is  $x_t = A\lambda_1^t + B\lambda_2^t$  for arbitrary constants  $A$  and  $B$ .

To obtain the Fibonacci sequence with  $x_0 = 0$  and  $x_1 = 1$  requires  $B = -A$  and  $1 = A(\lambda_1 - \lambda_2) = A\sqrt{5}$ , so  $B = -A = -\frac{1}{5}\sqrt{5}$ .

Hence  $x_t = \frac{1}{5}\sqrt{5} \cdot 2^{-t} [(1 + \sqrt{5})^t - (1 - \sqrt{5})^t]$ , so  $x_t \in \mathbb{N}$ .

## Case 2: Two Complex Conjugate Roots

Consider next the case where the equation  $\lambda^2 + a\lambda + b = 0$  has two complex conjugate roots that we write as

$$\lambda = re^{\pm i\theta} = r(\cos \theta \pm i \sin \theta) \quad \text{where } \sin \theta \neq 0$$

In this case  $\lambda^2 + a\lambda + b = (\lambda - re^{i\theta})(\lambda - re^{-i\theta})$  where

$$a = re^{i\theta} + re^{-i\theta} = r(\cos \theta + i \sin \theta) + r(\cos \theta - i \sin \theta) = 2r \cos \theta$$

and  $b = (re^{i\theta})(re^{-i\theta}) = r^2$  with  $\sin \theta \neq 0$ .

It follows that  $a^2 - 4b = 4r^2 \cos^2 \theta - 4r^2 = -4r^2 \sin^2 \theta < 0$ .

Note that  $r = \sqrt{|b|}$  and  $\theta = \arccos\left(\frac{a}{2r}\right) = \arccos\left(\frac{1}{2}a|b|^{-\frac{1}{2}}\right)$ .

## Case 2: Oscillating Solutions

In the complex plane  $\mathbb{C}$ , two possible solutions of the difference equation  $x_{t+1} + ax_t + bx_{t-1} = 0$  with  $x_0 \neq 0$  are

$$x_t^{(1)} = x_0(re^{i\theta})^t = x_0r^te^{i\theta t} = x_0r^t(\cos \theta t + i \sin \theta t)$$

$$\text{and } x_t^{(2)} = x_0(re^{-i\theta})^t = x_0r^te^{-i\theta t} = x_0r^t(\cos \theta t - i \sin \theta t)$$

In the real line  $\mathbb{R}$ , two possible solutions are

$$x_t^{(1)} = r^t \cos \theta t \quad \text{and} \quad x_t^{(2)} = r^t \sin \theta t$$

These are linearly independent because

$$\begin{vmatrix} x_0^{(1)} & x_0^{(2)} \\ x_1^{(1)} & x_1^{(2)} \end{vmatrix} = \begin{vmatrix} 1 & 0 \\ r \cos \theta & r \sin \theta \end{vmatrix} = r \sin \theta \neq 0$$

The general solution is therefore  $x_t = r^t(A \cos \theta t + B \sin \theta t)$  for arbitrary real constants  $A$  and  $B$ , where  $A = x_0$ .

## Case 3: Two Coincident Roots

In this case  $\lambda^2 + a\lambda + b = (\lambda - \bar{\lambda})^2$ ,  
where  $a = -2\bar{\lambda}$  and  $b = \bar{\lambda}^2$ .

Consider the perturbed equation  $x_{t+1} + ax_t + \tilde{b}x_{t-1} = 0$   
where  $a = -2\bar{\lambda}$  still and  $\tilde{b} = \bar{\lambda}^2 - \epsilon^2$  with  $\epsilon$  a small positive  
number.

We consider the behaviour of its general solution as  $\epsilon \rightarrow 0$ .

The auxiliary equation  $\lambda^2 + a\lambda + \tilde{b} = 0$   
can be written as  $\lambda^2 - 2\bar{\lambda}\lambda + \bar{\lambda}^2 - \epsilon^2 = 0$ .

Note that  $\lambda^2 - 2\bar{\lambda}\lambda + \bar{\lambda}^2 - \epsilon^2 = (\lambda - \bar{\lambda} + \epsilon)(\lambda - \bar{\lambda} - \epsilon)$ .

So the perturbed auxiliary equation  
has the two real roots  $\lambda = \bar{\lambda} \pm \epsilon$ .

## The Solution with Fixed Initial Conditions

Fix  $\bar{x}_0$  and  $\bar{x}_1$ .

The general solution satisfying  $x_0 = \bar{x}_0$  and  $x_1 = \bar{x}_1$  is  $x_t = A(\bar{\lambda} + \epsilon)^t + B(\bar{\lambda} - \epsilon)^t$  where  $\bar{x}_0 = A + B$  and  $\bar{x}_1 = A(\bar{\lambda} + \epsilon) + B(\bar{\lambda} - \epsilon) = (A + B)\bar{\lambda} + (A - B)\epsilon$ .

Hence  $A + B = \bar{x}_0$  and  $A - B = (1/\epsilon)(\bar{x}_1 - \bar{x}_0\bar{\lambda})$ , implying that  $A = \frac{1}{2} [\bar{x}_0 + (1/\epsilon)(\bar{x}_1 - \bar{x}_0\bar{\lambda})]$  and  $B = \frac{1}{2} [\bar{x}_0 - (1/\epsilon)(\bar{x}_1 - \bar{x}_0\bar{\lambda})]$ .

The solution for fixed  $\epsilon$  is therefore

$$x_t^\epsilon = \frac{1}{2} [\bar{x}_0 + (1/\epsilon)(\bar{x}_1 - \bar{x}_0\bar{\lambda})] (\bar{\lambda} + \epsilon)^t + \frac{1}{2} [\bar{x}_0 - (1/\epsilon)(\bar{x}_1 - \bar{x}_0\bar{\lambda})] (\bar{\lambda} - \epsilon)^t$$

which can be rewritten as

$$x_t^\epsilon = \frac{1}{2}\bar{x}_0 [(\bar{\lambda} + \epsilon)^t + (\bar{\lambda} - \epsilon)^t] + \frac{1}{2}(\bar{x}_1 - \bar{x}_0\bar{\lambda})(1/\epsilon) [(\bar{\lambda} + \epsilon)^t - (\bar{\lambda} - \epsilon)^t]$$

## The Limiting Solution as $\epsilon \rightarrow 0$

The limit of  $x_t^\epsilon$  as  $\epsilon \rightarrow 0$  takes the form

$$\bar{x}_0 \bar{\lambda}^t + \frac{1}{2}(\bar{x}_1 - \bar{x}_0 \bar{\lambda}) \lim_{\epsilon \rightarrow 0} (1/\epsilon) [(\bar{\lambda} + \epsilon)^t - (\bar{\lambda} - \epsilon)^t]$$

To evaluate the last limit, apply l'Hôpital's rule to obtain

$$\begin{aligned} & \lim_{\epsilon \rightarrow 0} [(\bar{\lambda} + \epsilon)^t - (\bar{\lambda} - \epsilon)^t] / \epsilon \\ &= \lim_{\epsilon \rightarrow 0} [t(\bar{\lambda} + \epsilon)^{t-1} + t(\bar{\lambda} - \epsilon)^{t-1}] / 1 \\ &= 2t\bar{\lambda}^{t-1} = (2t/\bar{\lambda})\bar{\lambda}^t \end{aligned}$$

Two linearly independent possible solutions

of the difference equation  $x_{t+1} + ax_t + bx_{t-1} = 0$

with  $x_0 \neq 0$  are  $x_t^{(1)} = x_0 \lambda^t$  and  $x_t^{(2)} = x_0 t \lambda^t$ .

There are two degrees of freedom in the difference equation.

Its general solution is  $x_t = (C + Dt)\lambda^t$   
for arbitrary real constants  $C$  and  $D$ .



## A Simpler Approach, I

We are trying to solve the homogeneous second-order difference equation with a repeated root  $\lambda$ , taking the form

$$x_{t+1} - 2\lambda x_t + \lambda^2 x_{t-1} = 0$$

We know that one solution is  $x_t = x_0 \lambda^t$  for arbitrary  $x_0$ .

To find a second linearly independent solution that we know must exist, try putting  $x_t = \lambda^t y_t$ .

Substituting into the original equation gives

$$\lambda^{t+1} y_{t+1} - 2\lambda^{t+1} y_t + \lambda^{t+1} y_{t-1} = 0$$

Disregarding the trivial case when  $\lambda = 0$ , one has  $y_{t+1} - 2y_t + y_{t-1} = 0$ .

## A Simpler Approach, II

To solve  $y_{t+1} - 2y_t + y_{t-1} = 0$ ,

try introducing yet another new variable  $z_t = y_{t+1} - y_t$ .

This leads to the new difference equation  $z_t - z_{t-1} = 0$  whose solution is obviously  $z_t = z_0$  for all  $t = 1, 2, \dots$

Then  $y_{t+1} - y_t = z_0$  for all  $t$ , implying that  $y_t = y_0 + z_0 t$ .

It follows that  $x_t = \lambda^t y_t = (y_0 + z_0 t)\lambda^t$ .

To conclude, two solutions are  $x_t^{(1)} = \lambda^t$  and  $x_t^{(2)} = t\lambda^t$ .

These are linearly independent because

$$\begin{vmatrix} x_0^{(1)} & x_0^{(2)} \\ x_1^{(1)} & x_1^{(2)} \end{vmatrix} = \begin{vmatrix} 1 & 0 \\ \lambda & \lambda \end{vmatrix} = \lambda \neq 0$$

The general solution is therefore  $x_t = (A + Bt)\lambda^t$  for arbitrary real constants  $A$  and  $B$ , where  $A = x_0$ .

# Lecture Outline

Solving Second-Order Equations

**Inhomogeneous Equations**

Particular Solutions in Two Special Cases

Method of Undetermined Coefficients

Higher-Order Linear Equations with Constant Coefficients

Stationary States and Stability for Second-Order Equations

## From Particular to General Solutions

The **homogeneous equation** with constant coefficients takes the form

$$x_{t+1} + ax_t + bx_{t-1} = 0$$

The associated **inhomogeneous equation** takes the form

$$x_{t+1} + ax_t + bx_{t-1} = f_t$$

for a general **forcing term**  $f_t$  on the RHS.

Let  $x_t^P$  denote a **particular solution**,  
and  $x_t^G$  any alternative **general solution**,  
of the inhomogeneous equation.

## Characterizing the General Solution

Our assumptions imply that, for each  $t = 1, 2, \dots$ , one has

$$x_{t+1}^P + ax_t^P + bx_{t-1}^P = f_t$$

$$x_{t+1}^G + ax_t^G + bx_{t-1}^G = f_t$$

Subtracting the first equation from the second implies that

$$x_{t+1}^G - x_{t+1}^P + a(x_t^G - x_t^P) + b(x_{t-1}^G - x_{t-1}^P) = 0$$

This shows that  $x_t^H := x_t^G - x_t^P$

solves the homogeneous equation  $x_{t+1} + ax_t + bx_{t-1} = 0$ .

So the general solution  $x_t^G$

of the inhomogeneous equation  $x_{t+1} + ax_t + bx_{t-1} = f_t$

with forcing term  $f_t$  is the sum  $x_t^P + x_t^H$  of

- ▶ any particular solution  $x_t^P$  of the inhomogeneous equation;
- ▶ the general solution  $x_t^H$  of the homogeneous equation.

## Linearity in the Forcing Term

### Theorem

Suppose that  $x_t^P$  and  $y_t^P$  are particular solutions of the two respective difference equations

$$x_{t+1} + ax_t + bx_{t-1} = d_t \quad \text{and} \quad y_{t+1} + ay_t + by_{t-1} = e_t$$

Then, for any scalars  $\alpha$  and  $\beta$ , the linear combination  $z_t^P := \alpha x_t^P + \beta y_t^P$  is a particular solution of the equation  $z_{t+1} + az_t + bz_{t-1} = \alpha d_t + \beta e_t$ .

### Proof.

Routine algebra. □

Consider any equation of the form  $x_{t+1} + ax_t + bx_{t-1} = f_t$  where  $f_t$  is a linear combination  $\sum_{k=1}^n \alpha_k f_t^k$  of  $n$  forcing terms.

The theorem implies that a particular solution is the corresponding linear combination  $\sum_{k=1}^n \alpha_k x_t^{Pk}$  of particular solutions to the equations  $x_{t+1} + ax_t + bx_{t-1} = f_t^k$ .

## Deriving an Explicit Particular Solution, I

In part A we were able to derive an explicit solution to the general first-order linear equation  $x_t - a_t x_{t-1} = f_t$ .

Here, for the special case of **constant coefficients**, we derive an explicit particular solution satisfying  $x_0 = x_1 = 0$  to the general second-order linear equation  $x_{t+1} + ax_t + bx_{t-1} = f_t$ .

Indeed, suppose that  $\lambda^2 + a\lambda + b = (\lambda - \lambda_1)(\lambda - \lambda_2)$  because  $\lambda_1$  and  $\lambda_2$  are the roots (possibly coincident, or possibly complex conjugates) of the auxiliary equation  $\lambda^2 + a\lambda + b = 0$ .

Introduce the new variable  $y_t = x_t - \lambda_1 x_{t-1}$ , implying that

$$\begin{aligned}y_{t+1} - \lambda_2 y_t &= x_{t+1} - \lambda_1 x_t - \lambda_2 x_t + \lambda_1 \lambda_2 x_{t-1} \\ &= x_{t+1} - (\lambda_1 + \lambda_2)x_t + \lambda_1 \lambda_2 x_{t-1} \\ &= x_{t+1} + ax_t + bx_{t-1} = f_t\end{aligned}$$

## Deriving an Explicit Particular Solution, II

Instead of the second-order equation  $x_{t+1} + ax_t + bx_{t-1} = f_t$ , we have the recursive pair of first-order equations

$$x_t - \lambda_1 x_{t-1} = y_t \quad \text{and} \quad y_{t+1} - \lambda_2 y_t = f_t \quad (\text{for } t = 1, 2, \dots)$$

where  $\lambda_1$  and  $\lambda_2$  are the roots of  $\lambda^2 + a\lambda + b = 0$ .

Given the initial conditions  $x_0 = x_1 = 0$  and so  $y_1 = 0$ , the explicit solutions like those derived in Part A are the sums

$$y_t = \sum_{k=1}^{t-1} \lambda_2^{t-k-1} f_k \quad \text{and} \quad x_t = \sum_{s=2}^t \lambda_1^{t-s} y_s \quad \text{for } t = 1, 2, \dots$$

Substituting the first equation in the second yields the double sum

$$x_t = \sum_{s=2}^t \lambda_1^{t-s} \sum_{k=1}^{s-1} \lambda_2^{s-k-1} f_k$$

which we would like to reduce to  $x_t = \sum_{k=1}^{t-1} \xi_{t-k-1} f_k$   
— i.e., a linear combination of the forcing terms  $(f_1, f_2, \dots, f_{t-1})$ .



## Deriving an Explicit Particular Solution, III

We begin by introducing

the mapping  $\mathbb{N} \times \mathbb{N} \ni (k, s) \mapsto 1_{ks}\{k < s\} \in \{0, 1\}$  defined by

$$1_{ks}\{k < s\} := \begin{cases} 1 & \text{if } k < s \\ 0 & \text{if } k \geq s \end{cases}$$

Then we can rewrite  $x_t = \sum_{s=2}^t \lambda_1^{t-s} \sum_{k=1}^{s-1} \lambda_2^{s-k-1} f_k$

as the double sum  $x_t = \sum_{s=2}^t \sum_{k=1}^{t-1} 1_{ks}\{k < s\} \lambda_1^{t-s} \lambda_2^{s-k-1} f_k$ .

Interchanging the order of summation gives

$$\begin{aligned} x_t &= \sum_{k=1}^{t-1} \sum_{s=2}^t 1_{ks}\{k < s\} \lambda_1^{t-s} \lambda_2^{s-k-1} f_k \\ &= \sum_{k=1}^{t-1} \left( \sum_{s=k+1}^t \lambda_1^{t-s} \lambda_2^{s-k-1} \right) f_k \\ &= \sum_{k=1}^{t-1} \left( \lambda_1^{t-k-1} + \lambda_1^{t-k-2} \lambda_2 + \dots + \lambda_1 \lambda_2^{t-k-2} + \lambda_2^{t-k-1} \right) f_k \end{aligned}$$

This reduces to  $x_t = \sum_{k=1}^{t-1} \xi_{t-k-1} f_k$  where  $\xi_m := \sum_{j=0}^m \lambda_1^{m-j} \lambda_2^j$ .

## Deriving an Explicit Particular Solution: IV

The value of the sum  $\xi_m = \sum_{j=0}^m \lambda_1^{m-j} \lambda_2^j$  depends on whether:

- ▶ we are in the **general case** when  $\lambda_1 \neq \lambda_2$ ;
- ▶ we are in the **degenerate case** when  $\lambda_1 = \lambda_2 = \lambda$ .

In the general case one has

$$(\lambda_1 - \lambda_2)\xi_m = \sum_{j=0}^m \left( \lambda_1^{m+1-j} \lambda_2^j - \lambda_1^{m-j} \lambda_2^{j+1} \right) = \lambda_1^{m+1} - \lambda_2^{m+1}$$

implying the particular solution

$$x_t^P = \frac{1}{\lambda_1 - \lambda_2} \sum_{k=1}^{t-1} \left( \lambda_1^{t-k} - \lambda_2^{t-k} \right) f_k$$

In the degenerate case one has  $\xi_m = (m+1)\lambda^m$ ,

implying the particular solution

$$x_t^P = \sum_{k=1}^{t-1} (t-k)\lambda^{t-k} f_k$$

# Lecture Outline

Solving Second-Order Equations

Inhomogeneous Equations

Particular Solutions in Two Special Cases

Method of Undetermined Coefficients

Higher-Order Linear Equations with Constant Coefficients

Stationary States and Stability for Second-Order Equations

## First Special Case with Distinct Real Roots, I

Consider the equation  $x_{t+1} + ax_t + bx_{t-1} = f_t$   
in the first special case when  $f_t = \mu^t$  with  $\mu \neq 0$ .

In the general case when the two roots  $\lambda_1$  and  $\lambda_2$   
of the auxiliary equation  $\lambda^2 + a\lambda + b = 0$  are distinct,  
the particular solution with  $x_0^P = x_1^P = 0$  is

$$x_t^P = \frac{1}{\lambda_1 - \lambda_2} \sum_{k=1}^{t-1} \left( \lambda_1^{t-k} - \lambda_2^{t-k} \right) \mu^k$$

But  $(\lambda - \mu) \sum_{k=1}^{t-1} \lambda^{t-k} \mu^k = \sum_{k=1}^{t-1} (\lambda^{t-k+1} \mu^k - \lambda^{t-k} \mu^{k+1})$ , so

$$x_t^P = \frac{1}{\lambda_1 - \lambda_2} \left( \frac{\lambda_1^t \mu - \lambda_1 \mu^t}{\lambda_1 - \mu} - \frac{\lambda_2^t \mu - \lambda_2 \mu^t}{\lambda_2 - \mu} \right)$$

in case  $\mu \notin \{\lambda_1, \lambda_2\}$ .

Disregarding the terms in  $\lambda_1^t$  and  $\lambda_2^t$   
that solve the corresponding homogeneous equation,  
the solution reduces to  $x_t^P = \alpha \mu^t$  for a suitable constant  $\alpha$ .

## First Special Case with Distinct Real Roots, II

The degenerate case when  $\mu \in \{\lambda_1, \lambda_2\}$  is more complicated.

In case  $\lambda_1 \neq \lambda_2 = \mu$ , the particular solution with  $x_0^P = x_1^P = 0$  is still

$$x_t^P = \frac{1}{\lambda_1 - \lambda_2} \sum_{k=1}^{t-1} \left( \lambda_1^{t-k} - \lambda_2^{t-k} \right) \mu^k$$

Because  $\lambda_2 = \mu$ , this reduces to

$$\begin{aligned} x_t^P &= \frac{1}{\lambda_1 - \mu} \sum_{k=1}^{t-1} \left( \lambda_1^{t-k} \mu^k - \mu^t \right) \\ &= \frac{1}{\lambda_1 - \mu} \left[ \frac{\lambda_1^t \mu - \lambda_1 \mu^t}{\lambda_1 - \mu} - (t-1) \mu^t \right] \end{aligned}$$

Disregarding the terms in  $\lambda_1^t$  and in  $\lambda_2^t = \mu^t$  that solve the corresponding homogeneous equation, the solution reduces to  $x_t^P = \alpha t \mu^t$  for a suitable constant  $\alpha$ .

## First Special Case with Coincident Real Roots

Consider now the degenerate case  
with coincident real roots  $\lambda_1 = \lambda_2 = \lambda$ .

So the inhomogeneous equation is  $x_{t+1} - 2\lambda x_t + \lambda^2 x_{t-1} = \mu^t$ .

As before, put  $y_t = x_t - \lambda x_{t-1}$  so that

$$y_{t+1} - \lambda y_t = x_{t+1} - \lambda x_t - \lambda x_t + \lambda^2 x_{t-1} = \mu^t$$

We consider again the particular solution  
with  $x_0 = x_1 = 0$  and so  $y_1 = 0$ .

## First Special Case with Coincident Real Roots: $\lambda \neq \mu$

Provided that  $\lambda \neq \mu$ , for  $t = 2, 3, \dots$  one has

$$y_t^P = \sum_{k=2}^t \lambda^{t-k} \mu^{k-1} = \frac{\mu(\lambda^{t-1} - \mu^{t-1})}{\lambda - \mu}$$

and then

$$\begin{aligned} x_t^P &= \sum_{k=2}^t \lambda^{t-k} y_k^P = \sum_{k=2}^t \lambda^{t-k} \mu \frac{\lambda^{k-1} - \mu^{k-1}}{\lambda - \mu} \\ &= \sum_{k=2}^t \frac{\mu \lambda^{t-1} - \lambda^{t-k} \mu^k}{\lambda - \mu} \\ &= \frac{\mu(t-1)\lambda^{t-1}}{\lambda - \mu} - \frac{\lambda^{t-1}\mu^2 - \mu^{t+1}}{(\lambda - \mu)^2} \end{aligned}$$

Hence  $x_t^P = (\alpha + \beta t)\lambda^t + \gamma\mu^t$  for suitable constants  $\alpha$ ,  $\beta$  and  $\gamma$  that depend on  $\lambda$  and  $\mu$ , but not on  $t$ .

Because  $(\alpha + \beta t)\lambda^t$  is a complementary solution of the homogeneous equation,

the particular solution can be reduced to  $x_t^P = \gamma\mu^t$ .

## First Special Case with Coincident Real Roots: $\lambda = \mu$

In case  $\lambda = \mu$ , however, for  $t = 2, 3, \dots$  one has

$$y_t^P = \sum_{k=2}^t \lambda^{t-k} \mu^{k-1} = (t-1)\lambda^{t-1}$$

and then

$$\begin{aligned} x_t^P &= \sum_{k=2}^t \lambda^{t-k} y_k^P = \sum_{k=2}^t \lambda^{t-k} (k-1)\lambda^{k-1} \\ &= \sum_{k=2}^t (k-1)\lambda^{t-1} = \frac{1}{2}t(t-1)\lambda^{t-1} \end{aligned}$$

Hence  $x_t^P = (\alpha t + \beta t^2)\lambda^t$  for suitable constants  $\alpha$  and  $\beta$  that depend on  $\lambda = \mu$ , but not on  $t$ .

Because  $\alpha t \lambda^t$  is a complementary solution of the homogeneous equation, the particular solution can be reduced to  $x_t^P = \beta t^2 \mu^t$ .



## Second Special Case: General Theorem

Consider next the equation  $x_{t+1} + ax_t + bx_{t-1} = f_t$   
in the second special case when  $f_t = t^r \mu^t$  with  $\mu \neq 0$  and  $r \in \mathbb{N}$ .

As before, let  $\lambda_1$  and  $\lambda_2$  denote the roots  
of the auxiliary equation  $\lambda^2 + a\lambda + b = 0$ .

### Theorem

*The difference equation  $x_{t+1} + ax_t + bx_{t-1} = t^r \mu^t$   
has a particular solution of the form  $x_t^P = \xi^P(t) \mu^t$   
where  $\xi^P(t) = \sum_{j=0}^d \xi_{rj} t^j$  is a polynomial in  $t$  which has degree:*

- ▶  $d = r$  in case  $\mu \notin \{\lambda_1, \lambda_2\}$ ;
- ▶  $d = r + 2$  in case  $\mu = \lambda_1 = \lambda_2$ ;
- ▶  $d = r + 1$  otherwise.

We begin the proof by introducing, as before,  
the new variable  $y_t := x_t - \lambda_1 x_{t-1}$ , implying that

$$\begin{aligned} y_{t+1} - \lambda_2 y_t &= x_{t+1} - \lambda_1 x_t - \lambda_2 x_t + \lambda_1 \lambda_2 x_{t-1} \\ &= x_{t+1} + ax_t + bx_{t-1} = t^r \mu^t \end{aligned}$$

## Continuing the Proof of the General Theorem

By the result in part A, the first-order equation  $y_{t+1} - \lambda_2 y_t = t^r \mu^t$  has a particular solution of the form  $y_t = Q(t)\mu^t$ , where  $Q(t) = \sum_{j=0}^d q_{rj} t^j$  is a polynomial in  $t$  which has degree:

- (i)  $d = r$  in case  $\mu \neq \lambda_2$ ;    (ii)  $d = r + 1$  in case  $\mu = \lambda_2$ .

By the linearity property of particular solutions, the equation

$$x_t - \lambda_1 x_{t-1} = y_t = Q(t)\mu^t = \sum_{j=0}^d q_{rj} t^j \mu^t$$

has a particular solution  $x_t^P = \xi^P(t)\mu^t$  where

$$x_t^P = \xi^P(t)\mu^t = \sum_{j=0}^d q_{rj} P_j(t)\mu^t$$

is the appropriate linear combination

of the particular solutions  $x_t = P_j(t)\mu^t$  ( $j = 0, 1, 2, \dots, d$ )

of the  $d + 1$  first-order equations  $x_t - \lambda_1 x_{t-1} = t^j \mu^t$ .

## Ending the Proof of the General Theorem

Again, using the result in part A,

for each  $j = 0, 1, 2, \dots, r$ , the solution  $x_t = P_j(t)\mu^t$  of the first-order difference equation  $x_t - \lambda_1 x_{t-1} = t^j \mu^t$  involves a polynomial  $P_j(t)$  in  $t$  which has degree:

- (i)  $j$  in case  $\mu \neq \lambda_1$ ;    (ii)  $j + 1$  in case  $\mu = \lambda_1$ .

So the degree of the highest order polynomial  $P_d(t)$  is

- (i)  $d$  in case  $\mu \neq \lambda_1$ ;    (ii)  $d + 1$  in case  $\mu = \lambda_1$ .

Combined with our previous result on whether  $d = r$  or  $d = r + 1$ , the degree  $d$  of  $\xi^P(t)$  is now easily seen to be

- ▶  $d = r$  in case  $\mu \notin \{\lambda_1, \lambda_2\}$ ;
- ▶  $d = r + 2$  in case  $\mu = \lambda_1 = \lambda_2$ ;
- ▶  $d = r + 1$  otherwise.



Using the notation  $\#S$  for the number of elements in a set  $S$ , these three cases can be summarized as  $d = r + 3 - \#\{\lambda_1, \lambda_2, \mu\}$ .

# Lecture Outline

Solving Second-Order Equations

Inhomogeneous Equations

Particular Solutions in Two Special Cases

**Method of Undetermined Coefficients**

Higher-Order Linear Equations with Constant Coefficients

Stationary States and Stability for Second-Order Equations

## First Special Case: A Simpler Approach

We have proved that the second-order difference equation

$$x_{t+1} + ax_t + bx_{t-1} = \mu^t$$

has a particular solution of the form  $x_t^P = \alpha\mu^t$ .

But there is a much easier way to find  $x_t^P$ , treating the parameter  $\alpha$  as an **undetermined coefficient**.

Indeed, for  $x_t = \alpha\mu^t$  to be a solution, one needs  $\alpha\mu^{t+1} + a\alpha\mu^t + b\alpha\mu^{t-1} = \mu^t$ .

Dividing each side by  $\mu^{t-1}$  yields the equation  $\alpha(\mu^2 + a\mu + b) = \mu$ .

In the **non-degenerate** case when  $\mu^2 + a\mu + b \neq 0$

because  $\mu$  is not a root

of the characteristic equation  $\lambda^2 + a\lambda + b = 0$ ,

one has  $\alpha = \mu(\mu^2 + a\mu + b)^{-1}$ .

Hence, a particular solution is  $x_t^P = (\mu^2 + a\mu + b)^{-1}\mu^{t+1}$ .

## Degenerate Case When $\mu$ is a Characteristic Root

The **simple degenerate case** occurs when  $\mu^2 + a\mu + b = 0$  because  $\mu$  equals one of the two **distinct** roots  $\lambda_1$  and  $\lambda_2$  of the characteristic equation  $\lambda^2 + a\lambda + b = 0$ .

Then we have proved that the second-order difference equation

$$x_{t+1} + ax_t + bx_{t-1} = \mu^t$$

has a particular solution of the form  $x_t^P = \alpha t \mu^t$ .

To determine the undetermined coefficient  $\alpha$ , we must solve

$$\alpha(t+1)\mu^{t+1} + a\alpha t\mu^t + b\alpha(t-1)\mu^{t-1} = \mu^t$$

Dividing each side by  $\mu^{t-1}$  and gathering terms yields the equation  $\alpha t(\mu^2 + a\mu + b) + \alpha(\mu^2 - b) = \mu$ .

Provided that  $\mu^2 \neq b$ , this reduces to  $\alpha = (\mu^2 - b)^{-1}\mu$ .

## Doubly Degenerate Case

When  $\mu^2 = b$ , however, the degenerate case is more complicated.

Indeed, the equation  $\mu^2 + a\mu + b = 0$  implies that  $a\mu + 2b = 0$ .

Hence  $\mu = -2b/a$ , so  $\mu^2 = b = 4b^2/a^2$  implying that  $a^2 = 4b$ .

Then the characteristic equation  $\lambda^2 + a\lambda + b = 0$  reduces to  $(\lambda - \mu)^2 = 0$ , with  $\mu$  as its repeated root.

Inspired by the earlier theorem,

we look for a particular solution of the form  $x_t^P = \alpha t^2 \mu^t$ .

To determine the undetermined coefficient  $\alpha$ , we must solve

$$\alpha(t+1)^2 \mu^{t+1} + a\alpha t^2 \mu^t + b\alpha(t-1)^2 \mu^{t-1} = \mu^t$$

Dividing each side by  $\mu^{t-1}$  and gathering terms yields

$$\alpha t^2 (\mu^2 + a\mu + b) + \alpha(2t+1)\mu^2 + \alpha b(-2t+1) = \mu$$

Because  $\mu^2 + a\mu + b = 0$  and  $0 \neq b = \mu^2$ ,

this equation reduces to  $2\alpha\mu^2 = \mu$ , implying that  $\alpha = 1/2\mu$ .

## Second Special Case

Again, inspired by earlier theorems, we can apply the method of undetermined coefficients to the equation

$$x_{t+1} + ax_t + bx_{t-1} = \sum_{k=1}^m \sum_{j=1}^{r_k} \alpha_{kj} t^j \mu_k^t$$

where we naturally assume that the constants  $\mu_k$  ( $k = 1, 2, \dots, m$ ) are all different.

A particular solution takes the form

$$x_t^P = \sum_{k=1}^m \sum_{j=1}^{d_k} \beta_{kj} t^j \mu_k^t$$

where the degree  $d_k$  of each polynomial  $\sum_{j=1}^{d_k} \beta_{kj} t^j$  with undetermined coefficients  $\langle \langle \beta_{kj} \rangle_{j=1}^{d_k} \rangle_{k=1}^m$  is

- ▶  $r_k$  in case  $\mu_k \notin \{\lambda_1, \lambda_2\}$ ;
- ▶  $r_k + 2$  in case  $\mu_k = \lambda_1 = \lambda_2$ ;
- ▶  $r_k + 1$  otherwise.



## Determining the Coefficients

The coefficients  $\langle\langle\beta_{kj}\rangle_{j=1}^{d_k}\rangle_{k=1}^m$  of the particular solution

$$x_t^P = \sum_{k=1}^m \sum_{j=1}^{d_k} \beta_{kj} t^j \mu_k^t$$

can be found (in principle!) by solving, for  $k = 1, 2, \dots, m$ , the  $m$  independent systems of linear equations that result from equating coefficients of powers of  $t$  in the expansions

$$\sum_{j=1}^{d_k} \beta_{kj} [(t+1)^j \mu_k^2 + at^j \mu_k^t + b(t-1)^j] = \sum_{j=1}^{r_k} \alpha_{kj} t^j \mu_k$$

# Lecture Outline

Solving Second-Order Equations

Inhomogeneous Equations

Particular Solutions in Two Special Cases

Method of Undetermined Coefficients

**Higher-Order Linear Equations with Constant Coefficients**

Stationary States and Stability for Second-Order Equations

# Higher-Order Linear Equations with Constant Coefficients

An  $n$ th order linear equation with constant coefficients takes the form

$$x_t + \sum_{r=1}^n a_r x_{t-r} = f_t$$

in the inhomogeneous case, and

$$x_t + \sum_{r=1}^n a_r x_{t-r} = 0$$

in the homogeneous case.

The corresponding auxiliary equation is  $\lambda^n + \sum_{r=1}^n a_r \lambda^{n-r} = 0$ .

## Roots of the Auxiliary Equation

The auxiliary equation can be written as  $p_n(\lambda) = 0$  whose LHS is the polynomial  $\lambda^n + \sum_{r=1}^n a_r \lambda^{n-r}$  of degree  $n$ .

By the **fundamental theorem of algebra**, this equation has at least one root  $\lambda_1$ , which may be complex.

Then  $p_n(\lambda)$  can be factored as  $p_n(\lambda) \equiv (\lambda - \lambda_1)p_{n-1}(\lambda)$ .

But now the equation  $p_{n-1}(\lambda) = 0$  also has at least one root  $\lambda_2$ , which may also be complex.

Repeating this argument  $n$  times, the auxiliary equation  $p_n(\lambda) = 0$  has  $n$  roots  $\lambda_1, \lambda_2, \dots, \lambda_n$ , some of which may be repeated.

In particular,  $p_n(\lambda) \equiv \prod_{i=1}^n (\lambda - \lambda_i)$ .

# Solving the Homogeneous Equation

## Theorem

Consider the homogeneous equation  $x_t + \sum_{r=1}^n a_r x_{t-r} = 0$ , and suppose that the auxiliary equation can be written as

$$0 = \lambda^n + \sum_{r=1}^n a_r \lambda^{t-r} = \prod_{j=1}^k (\lambda - \rho_j)^{m_j}$$

with  $k$  distinct roots  $\rho_j$  ( $j = 1, 2, \dots, k$ )

whose respective **multiplicities**  $m_j$  satisfy  $\sum_{j=1}^k m_j = n$ .

Then the general solution of the homogeneous equation takes the form

$$x_t = \sum_{j=1}^k \sum_{h=1}^{m_j} \alpha_{jh} t^{h-1} \rho_j^t$$

for  $n$  arbitrary constants  $\alpha_{jh}$  ( $h = 1, 2, \dots, m_j$  and  $j = 1, 2, \dots, k$ ).

That is, the general solution is an arbitrary linear combination of the functions  $t^{h-1} \rho_j^t$  ( $h = 1, 2, \dots, m_j$  and  $j = 1, 2, \dots, k$ ).

# Solving the Inhomogeneous Equation

## Theorem

The general solution of the inhomogeneous equation

$$x_t + \sum_{r=1}^n a_r x_{t-r} = \sum_{h=1}^i \sum_{j=1}^{q_h} \alpha_{hj} t^j \mu_h^t$$

is the sum of: (i) the general **complementary** solution to the corresponding homogeneous equation  $x_t + \sum_{r=1}^n a_r x_{t-r} = 0$ ; and (ii) any particular solution.

One particular solution takes the form  $x_t^P = \sum_{h=1}^i \sum_{j=1}^{d_h} \beta_{hj} t^j \mu_h^t$  where the degree  $d_h$  of each polynomial  $\sum_{j=1}^{d_h} \beta_{hj} t^j$  with undetermined coefficients  $\langle \langle \beta_{hj} \rangle_{j=1}^{d_h} \rangle_{h=1}^i$  is

- ▶  $q_h$  in case  $\mu_h \notin \{\rho_1, \rho_2, \dots, \rho_k\}$ ;
- ▶  $q_h + m_j$  in case  $\mu_h = \rho_j$ , a root of multiplicity  $m_j$ .

# Lecture Outline

Solving Second-Order Equations

Inhomogeneous Equations

Particular Solutions in Two Special Cases

Method of Undetermined Coefficients

Higher-Order Linear Equations with Constant Coefficients

Stationary States and Stability for Second-Order Equations

## Stationary States of a Linear Equation

Consider the second-order equation  $x_{t+1} + ax_t + bx_{t-1} = f$  for a constant **forcing term**  $f \in \mathbb{R}$ .

Here a stationary state  $x^* \in \mathbb{R}$  has the defining property that  $x_{t-1} = x_t = x^* \implies x_{t+1} = x^*$ .

This is satisfied if and only if  $x^* + ax^* + bx^* = f$ , or equivalently, if and only if  $(1 + a + b)x^* = f$ .

In case  $a + b = -1$ , there is:

- ▶ no stationary state unless  $f = 0$ ;
- ▶ a whole real line  $\mathbb{R}$  of stationary states if  $f = 0$ .

Otherwise, if  $a + b \neq -1$ , the only stationary state is  $x^* = (1 + a + b)^{-1}f$ .



## Stability of a Linear Equation

If  $a + b \neq -1$ , let  $y_t := x_t - x^*$  denote the **deviation** of state  $x_t$  from the stationary state  $x^* = (1 + a + b)^{-1}f$ . Then

$$\begin{aligned}y_{t+1} &= x_{t+1} - x^* = -ax_t - bx_{t-1}f - x^* \\ &= -a(y_t + x^*) - b(y_{t-1} + x^*) + f - x^* = -ay_t - by_{t-1}\end{aligned}$$

Thus  $y_t$  solves the homogenous equation  $x_{t+1} + ax_t + bx_{t-1} = 0$ .

As already seen, the solution to this homogeneous equation depends on the two roots  $\lambda_{1,2} = -\frac{1}{2}a \pm \frac{1}{2}\sqrt{a^2 - 4b}$  of the quadratic characteristic equation

$$f(\lambda) \equiv \lambda^2 + a\lambda + b \equiv (\lambda - \lambda_1)(\lambda - \lambda_2) = 0$$

There are three cases to consider:

1. two distinct real roots because  $a^2 - 4b > 0$ ;
2. two complex conjugate roots because  $a^2 - 4b < 0$ ;
3. two coincident real roots because  $a^2 - 4b = 0$ .

## Stability Condition

With two distinct roots  $\lambda_1$  and  $\lambda_2$ , real or complex, the general solution of the homogeneous equation is  $x_t = A\lambda_1^t + B\lambda_2^t$ .

Stability is satisfied if and only if for all  $A, B \in \mathbb{R}$  one has  $x_t \rightarrow 0$  as  $t \rightarrow \infty$ .

This is true if and only if the absolute values of both roots satisfy  $|\lambda_1| < 1$  and  $|\lambda_2| < 1$ .

With two coincident roots  $\lambda_1 = \lambda_2 = -\frac{1}{2}a = \sqrt{b}$ , the general solution of the homogeneous equation is  $x_t = (A + Bt)\lambda^t$ .

Again, stability is satisfied if and only if for all  $A, B \in \mathbb{R}$  one has  $x_t \rightarrow 0$  as  $t \rightarrow \infty$ .

This is true if and only if the absolute value of the double root satisfies  $|\lambda| < 1$ .

## Two Distinct Real Roots

Here  $\lambda^2 + a\lambda + b = (\lambda - \lambda_1)(\lambda - \lambda_2)$

where  $\lambda_1$  and  $\lambda_2$  are both real.

Note that the quadratic function  $f(\lambda) \equiv \lambda^2 + a\lambda + b$  is convex and satisfies  $f(\lambda) \rightarrow +\infty$  as  $\lambda \rightarrow \pm\infty$ .

So the real roots of  $f(\lambda) = 0$  satisfy  $|\lambda_1| < 1$  and  $|\lambda_2| < 1$  iff

$$f(-1) > 0 \text{ and } f(1) > 0 \text{ with } f'(-1) < 0 \text{ and } f'(1) > 0$$

These conditions are equivalent to

$$1 - a + b > 0 \text{ and } 1 + a + b > 0 \text{ with } -2 + a < 0 \text{ and } 2 + a > 0$$

or to  $|a| < 2$  and  $|a| < 1 + b$ .

Together with the condition  $a^2 > 4b$

for the equation  $f(\lambda) = 0$  to have two distinct real roots, these inequalities are equivalent to  $|a| - 1 < b < 1$ .

## Two Complex Conjugate Roots

The characteristic equation  $\lambda^2 + a\lambda + b = 0$  has two complex conjugate roots when  $a^2 - 4b < 0$ .

In this case, these characteristic roots are

$$\lambda_{1,2} = -\frac{1}{2}a \pm \frac{1}{2}i\sqrt{4b - a^2} = r e^{\pm i\theta} = r(\cos \theta \pm i \sin \theta)$$

where  $r = \sqrt{b}$  and  $\theta = \arccos(a/2\sqrt{b})$

Then the general solution of the homogeneous equation can be written as  $x_t = r^t(A \cos \theta t + B \sin \theta t)$ .

Stability is satisfied if and only if for all  $A, B \in \mathbb{R}$  one has  $x_t \rightarrow 0$  as  $t \rightarrow \infty$ .

This is true if and only if  $b < 1$ , as well as  $a^2 - 4b < 0$  which implies that  $b > 0$ .

## A Repeated Real Root

The characteristic equation  $\lambda^2 + a\lambda + b = 0$  has two coincident real roots when  $a^2 = 4b$ .

In this case,  $\lambda^2 + a\lambda + b = (\lambda + \frac{1}{2}a)^2$ .

The coincident real roots both equal  $-\frac{1}{2}a$ .

Then the general solution of the homogeneous equation is  $x_t = (A + Bt)(-\frac{1}{2}a)^t$ .

Stability is satisfied if and only if for all  $A, B \in \mathbb{R}$  one has  $x_t \rightarrow 0$  as  $t \rightarrow \infty$ .

This is true if and only if the modulus of the repeated root  $\lambda = -\frac{1}{2}a$  satisfies  $|\lambda| < 1$ , and so if and only if  $|a| < 2$ .

# A Simpler Stability Condition

## Theorem

*The linear autonomous equation  $x_{t+1} + ax_t + bx_{t-1} = f$  is stable, both locally and globally, if and only if  $|a| < 1 + b < 2$ .*


## Proof.

Stability requires one of the following three to hold:

1. distinct real roots because  $a^2 > 4b$ , with  $|a| - 1 < b < 1$ ;
2. complex conjugate roots because  $a^2 < 4b$ , with  $b < 1$ ;
3. a repeated real root because  $a^2 = 4b$ , with  $|a| < 2$ .

A diagram in the  $(a, b)$ -plane shows that one of these three holds if and only if  $|a| < 1 + b < 2$ . □

# Diagram of Stable Region



The stable region occurs where  $|a| - 1 < b < 1$ , in the interior of an isosceles right-angled triangle with corners at  $(a, b) = (0, -1)$  and  $(a, b) = (\pm 2, 1)$ .

## Stability with a Variable Forcing Term

Consider now the second-order equation  $x_{t+1} + ax_t + bx_{t-1} = f_t$  for a **variable** forcing term  $f_t$ .

The general solution takes the form  $x_t^G = x_t^H + x_t^P$  where:

- ▶  $x_t^P$  is one particular solution of  $x_{t+1} + ax_t + bx_{t-1} = f_t$ ;
- ▶  $x_t^H$  is any one of a two-dimension continuum of solutions to the homogeneous equation  $x_{t+1} + ax_t + bx_{t-1} = 0$ .

The stability condition  $|a| < 1 + b < 2$  is necessary and sufficient for any solution of the homogeneous equation to satisfy  $x_t^H \rightarrow 0$  as  $t \rightarrow \infty$ .

It is therefore also necessary and sufficient for the difference between any two solutions  $x_t^{(1)}$  and  $x_t^{(2)}$  of the inhomogeneous equation  $x_{t+1} + ax_t + bx_{t-1} = f_t$  to satisfy  $x_t^{(1)} - x_t^{(2)} \rightarrow 0$  as  $t \rightarrow \infty$ .

In the long run, this means that there is an **asymptotically unique** solution to  $x_{t+1} + ax_t + bx_{t-1} = f_t$ .