

# Lecture Notes 7: Dynamic Equations

## Part D: Differential Equations

Peter J. Hammond

Latest revision 2024 September 16th;  
typeset from `dynEqLects24D.tex`

# Lecture Outline

## First-Order Differential Equations in One Variable

### Introduction

Picard's Method

General First-Order Affine Equation

Constant and Undetermined Coefficients

Stability in the Autonomous Case

## Second-Order Differential Equations in One Variable

### Introduction

The Inhomogeneous Equation

The Method of Undetermined Coefficients

Stability

## First-Order Multivariable Differential Equations

### Introduction

Prominent Examples and Stability Conditions

Autonomous Nonlinear Equations in Many Variables

Existence and Uniqueness Theorem

# First-Order Differential Equations

The typical first-order differential equation in one variable  $x$  is

$$\dot{x} = \frac{dx}{dt} = f(x, t)$$

The equation is **autonomous** just in case  $f$  is independent of  $t$ , so it can be written as  $\dot{x} = f(x)$ .

Typically one imposes an **initial condition** requiring  $x(s) = \bar{x}_s$  at time  $s$  (not necessarily the earliest time).

Then any solution is a **fixed function**  $t \mapsto x(t)$  that satisfies the corresponding **integral equation**  $x(t) = \bar{x}_s + \int_s^t f(x(u), u) du$ .

**Picard's method of successive approximations** starts with an arbitrary function  $t \mapsto x^{(0)}(t)$  satisfying  $x^{(0)}(s) = \bar{x}_s$ .

Then it computes  $x^{(n)}(t) = \bar{x}_s + \int_s^t f(x^{(n-1)}(u), u) du$  for  $n \in \mathbb{N}$ .

**If** convergence occurs, the limit as  $n \rightarrow \infty$  will be a solution.

## Right-Hand Side Independent of $x$

A special case occurs when the right-hand side  $f(x, t)$  is independent of  $x$ .

Then the differential equation can be written as

$$\frac{dx}{dt} = g(t)$$

Its solution can be written as the **indefinite integral**

$$x(t) = \int g(t)dt$$

Introducing an **initial condition**  $x(s) = \bar{x}_s$

at a particular **start time**  $s$

allows the solution to be written as the **definite integral**

$$x(t) = \bar{x}_s + \int_s^t g(\tau)d\tau$$

CHECK that this alleged solution satisfies  $x(s) = \bar{x}_s$  and  $\dot{x}(t) = g(t)$  for all  $t \geq s$ .

## Leibniz's Rule for Differentiating an Integral

Consider the function  $F : \mathbb{R}^3 \rightarrow \mathbb{R}$  defined by

$$F(a, b, u) := \int_a^b f(t, u) dt$$

Its three first-order partial derivatives are:

$$(i) F'_a = -f(a, u); \quad (ii) F'_b = f(b, u); \quad (iii) F'_u = \int_a^b \frac{\partial}{\partial u} f(t, u) dt$$

Applying the chain rule, the total derivative of the integral function  $y \mapsto I(y) := \int_{a(y)}^{b(y)} f(t, y) dt$  satisfies

$$\begin{aligned} I'(y) &= \frac{d}{dy} F(a(y), b(y), y) = a'(y)F'_a + b'(y)F'_b + F'_u \\ &= b'(y)f(b(y), y) - a'(y)f(a(y), y) + \int_{a(y)}^{b(y)} \frac{\partial}{\partial y} f(t, y) dt \end{aligned}$$

# Lecture Outline

## First-Order Differential Equations in One Variable

Introduction

**Picard's Method**

General First-Order Affine Equation

Constant and Undetermined Coefficients

Stability in the Autonomous Case

## Second-Order Differential Equations in One Variable

Introduction

The Inhomogeneous Equation

The Method of Undetermined Coefficients

Stability

## First-Order Multivariable Differential Equations

Introduction

Prominent Examples and Stability Conditions

Autonomous Nonlinear Equations in Many Variables

Existence and Uniqueness Theorem

## Picard's Method of Successive Approximations

The simplest first-order equation with constant coefficients takes the form

$$\dot{x}(t) = ax(t), \text{ with } x(0) \text{ given}$$

It corresponds to the integral equation

$$x(t) - x(0) = \int_0^t ax(u) du \text{ for all } t \geq 0$$

Starting with even a very crude approximation such as the constant function  $x^{(0)}(t) \equiv x(0)$  for all  $t \geq 0$ , we can calculate a sequence  $t \mapsto x^{(n)}(t)$  ( $n \in \mathbb{N}$ ) of successive approximations to a solution  $[0, \infty) \ni t \mapsto x(t) \in \mathbb{R}$  using, for all  $t \geq 0$ , the iterative rule

$$x^{(n)}(t) - x(0) = \int_0^t f(x^{(n-1)}(u), u) du = \int_0^t ax^{(n-1)}(u) du$$

## Initial Three Iterations

Starting from  $x^{(0)}(t) \equiv x(0)$ , iterating once gives

$$x^{(1)}(t) - x(0) = \int_0^t a x^{(0)}(u) du = a x(0) t$$

Iterating a second time gives

$$x^{(2)}(t) - x(0) = \int_0^t a x(0)(1 + au) du = a x(0) t + \frac{1}{2} a^2 x(0) t^2$$

Iterating a third time gives

$$\begin{aligned} x^{(3)}(t) - x(0) &= \int_0^t [a x(0) + a^2 x(0) u + \frac{1}{2} a^3 x(0) u^2] du \\ &= a x(0) t + \frac{1}{2} a^2 x(0) t^2 + \frac{1}{6} a^3 x(0) t^3 \end{aligned}$$

## Terms of the Sum

Each time we are adding one term to a sum.

So, starting with  $y^{(0)}(t) \equiv x(0)$ ,

define the new incremental variable  $y^{(n)}(t) := x^{(n)}(t) - x^{(n-1)}(t)$ .

This implies that  $x^{(n)}(t) = x(0) + \sum_{k=1}^n y^{(k)}(t)$ .

Subtract  $x^{(n)}(t) - x(0) = \int_0^t a x^{(n-1)}(u) du$

from  $x^{(n+1)}(t) - x(0) = \int_0^t a x^{(n)}(u) du$

to obtain  $y^{(n+1)}(t) = \int_0^t a y^{(n)}(u) du$ .

Now we obtain successively

$$y^{(1)}(t) = \int_0^t a x(0) du = a x(0) t$$

$$y^{(2)}(t) = \int_0^t a^2 x(0) u du = \frac{1}{2} a^2 x(0) t^2$$

$$y^{(3)}(t) = \int_0^t \frac{1}{2} a^3 x(0) u^2 du = \frac{1}{6} a^3 x(0) t^3$$

This suggests the induction hypothesis  $y^{(n)}(t) = \frac{1}{n!} a^n x(0) t^n$ .

## Constructing the Sum

The induction hypothesis  $y^{(n)}(t) = \frac{1}{n!} a^n x(0) t^n$   
and the relation  $y^{(n+1)}(t) = \int_0^t a y^{(n)}(u) du$  together imply that

$$\begin{aligned} y^{(n+1)}(t) &= \int_0^t a \frac{1}{n!} a^n x(0) u^n du = \frac{1}{n!} a^{n+1} x(0) \int_0^t u^n du \\ &= \frac{1}{n!} a^{n+1} x(0) \frac{1}{n+1} t^{n+1} = \frac{1}{(n+1)!} a^{n+1} x(0) t^{n+1} \end{aligned}$$

This confirms the induction hypothesis with  $n$  replaced by  $n + 1$ .

It follows that  $y^{(n)}(t) = \frac{1}{n!} a^n x(0) t^n$  for all  $n \in \mathbb{N}$

and then that  $x^{(n)}(t) = x(0) + \sum_{k=1}^n \frac{1}{k!} a^k x(0) t^k$ .

# Euler's Number and the Exponential Function

**Euler's number** was invented by Jacob Bernoulli in 1683.

Euler chose to denote it by  $e$ .

Recall that it is given by

$$e = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n \approx 2.718281828$$

My late co-author Knut Sydsæter, as a cultured Norwegian, recognized 1828 as the year when their great playwright Henrik Ibsen was born.

So Knut remembered this 10 digit approximation as “2.7 Ibsen Ibsen”.



# The Exponential Function and Exponential Solution

The **exponential function**, which satisfies  $\exp x = e^x$ , satisfies

$$\exp x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots = 1 + \sum_{n=1}^{\infty} \frac{1}{n!} x^n = e^x$$

As  $n \rightarrow \infty$ , the Picard approximate solution  $x^{(n)}(t)$  to the differential equation that we found earlier converges to the infinite series

$$x(0) + \sum_{k=1}^{\infty} \frac{1}{k!} a^k x(0) t^k = x(0) \exp(at) = x(0) e^{at}$$

# Lecture Outline

## First-Order Differential Equations in One Variable

Introduction

Picard's Method

**General First-Order Affine Equation**

Constant and Undetermined Coefficients

Stability in the Autonomous Case

## Second-Order Differential Equations in One Variable

Introduction

The Inhomogeneous Equation

The Method of Undetermined Coefficients

Stability

## First-Order Multivariable Differential Equations

Introduction

Prominent Examples and Stability Conditions

Autonomous Nonlinear Equations in Many Variables

Existence and Uniqueness Theorem

## General First-Order Affine Equation

The general first-order affine equation takes the form

$$\dot{x}(t) = a(t)x(t) + b(t)$$

for arbitrary integrable functions  $t \mapsto a(t)$  and  $t \mapsto b(t)$ .

In the **homogeneous** case one has  $b(t) \equiv 0$ , and the equation takes the linear form  $\dot{x}(t) = a(t)x(t)$ .

Assuming that  $x > 0$  for all  $t$ , we can take logs and write the equation as

$$\frac{d}{dt} \ln x = \frac{\dot{x}}{x} = a(t)$$

After introducing the new variable  $y(t) := \ln x(t)$ , the equation becomes  $\dot{y} = a(t)$  whose solution is obviously

$$y(t) = y(s) + \int_s^t a(\tau) d\tau$$

## Solution in the Homogeneous Case

Because  $x(t) = \exp y(t)$ , the solution for  $x$  is

$$x(t) = \exp[y(t)] = \exp[y(s)] \exp \left[ \int_s^t a(\tau) d\tau \right] = x(s) \alpha_s(t)$$

where  $\alpha_s(t)$  denotes the **integrating factor**  $\exp \left[ \int_s^t a(\tau) d\tau \right]$ .

In the special case of an autonomous equation where  $a(\tau) = a$  constant, one has  $\int_s^t a(\tau) d\tau = a(t - s)$  and so  $\alpha_s(t) = e^{a(t-s)}$ .

## The Non-Homogeneous Case

The solution  $x(t) = x(s)\alpha_s(t)$

to the homogeneous equation  $\dot{x}(t) - a(t)x(t) = 0$

can be used to help solve the corresponding

non-homogeneous equation  $\dot{x}(t) - a(t)x(t) = f(t)$ .

Indeed, consider the result of dividing

each side of this non-homogeneous equation

by the **integrating factor**  $\alpha_s(t) := \exp\left[\int_s^t a(\tau)d\tau\right]$

whose reciprocal is  $1/\alpha_s(t) := \exp\left[-\int_s^t a(\tau)d\tau\right]$ .

Note that  $\frac{d}{dt}\left[-\int_s^t a(\tau)d\tau\right] = -a(t)$ ,

implying that  $\frac{d}{dt}[1/\alpha_s(t)] = -a(t)/\alpha_s(t)$  so, by the product rule

$$\frac{d}{dt}[x(t)/\alpha_s(t)] = [1/\alpha_s(t)]\dot{x}(t) - [a(t)/\alpha_s(t)]x(t) = f(t)/\alpha_s(t)$$

for any solution of the equation  $\dot{x}(t) - a(t)x(t) = f(t)$ .

## Solving the Non-Homogeneous Equation

Integrating each side of the equation  $\frac{d}{dt}[x(t)/\alpha_s(t)] = f(t)/\alpha_s(t)$  over the interval from  $s$  to  $t$  gives us

$$\int_s^t [x(u)/\alpha_s(u)]' du = \frac{x(t)}{\alpha_s(t)} - \frac{x(s)}{\alpha_s(s)} = \int_s^t \frac{f(u)}{\alpha_s(u)} du$$

The definition  $\alpha_s(t) = \exp\left[\int_s^t a(\tau)d\tau\right]$  implies that  $\alpha_s(s) = 1$  and also  $\alpha_s(t)/\alpha_s(u) = \alpha_u(t)$ .

Hence, multiplying each side by  $\alpha_s(t)$  gives the solution

$$\begin{aligned} x(t) &= \alpha_s(t) \left[ x(s) + \int_s^t [1/\alpha_s(u)] f(u) du \right] \\ &= \alpha_s(t)x(s) + \int_s^t \alpha_u(t) f(u) du \\ &= \exp\left[\int_s^t a(\tau)d\tau\right] x(s) + \int_s^t \exp\left[\int_u^t a(\tau)d\tau\right] f(u) du \end{aligned}$$

## Linearity in the Forcing Term

### Theorem

Suppose that  $x^P(t)$  and  $y^P(t)$  are particular solutions of the two respective differential equations

$$\dot{x}(t) - a(t)x(t) = d(t) \quad \text{and} \quad \dot{y}(t) - a(t)y(t) = e(t)$$

Then, for any scalars  $\alpha$  and  $\beta$ ,

the equation  $\dot{z}(t) - a(t)z(t) = f(t) = \alpha d(t) + \beta e(t)$

has as a particular solution

the corresponding linear combination  $z^P(t) := \alpha x^P(t) + \beta y^P(t)$ .

Consider any equation of the form  $\dot{x}(t) - a(t)x(t) = f(t)$

where  $f(t)$  is a linear combination  $\sum_{k=1}^n \alpha_k f^k(t)$

of  $n$  forcing terms.

The theorem implies that a particular solution

is the corresponding linear combination  $\sum_{k=1}^n \alpha_k x^{Pk}(t)$

of particular solutions to the  $n$  equations  $\dot{x}(t) - a(t)x(t) = f^k(t)$ .

# Lecture Outline

## First-Order Differential Equations in One Variable

Introduction

Picard's Method

General First-Order Affine Equation

**Constant and Undetermined Coefficients**

Stability in the Autonomous Case

## Second-Order Differential Equations in One Variable

Introduction

The Inhomogeneous Equation

The Method of Undetermined Coefficients

Stability

## First-Order Multivariable Differential Equations

Introduction

Prominent Examples and Stability Conditions

Autonomous Nonlinear Equations in Many Variables

Existence and Uniqueness Theorem

## First-Order Linear Equation with a Constant Coefficient

Next, consider the equation  $\dot{x}(t) - ax(t) = f(t)$

where the coefficient  $a$  of  $x(t)$  has become the **constant**  $a \neq 0$ .

The solution we found for the general case was

$$x(t) = \exp \left[ \int_s^t a(\tau) d\tau \right] x(s) + \int_s^t \exp \left[ \int_u^t a(\tau) d\tau \right] f(u) du$$

When  $a(t) = a$ , independent of  $t$ , this reduces to

$$x(t) = e^{a(t-s)} x(s) + \int_s^t e^{a(t-u)} f(u) du$$

We simplify further by choosing the initial time  $s = 0$ .

Then

$$x(t) = e^{at} x(0) + \int_0^t e^{a(t-u)} f(u) du$$

## First Special Case

An interesting special case occurs when the forcing term  $f(t)$  is the exponential function  $t \mapsto e^{\mu t}$ .

Then the solution is

$$x(t) = e^{at}x(0) + \int_0^t e^{a(t-u)+\mu u} du = e^{at} \left[ x(0) + \int_0^t e^{(\mu-a)u} du \right]$$

In the **degenerate case** when  $\mu = a$ , one has  $\int_0^t e^{(\mu-a)u} du = \int_0^t 1 du = t$ , so the solution collapses to

$$x(t) = e^{at} [x(0) + t]$$

This solution can be written as  $x(t) = x^H(t) + x^P(t)$  where:

1.  $x^H(t) = \xi^H e^{at}$  with  $\xi^H := x(0)$  is a complementary solution of the **homogeneous** equation  $\dot{x}(t) - ax(t) = 0$ ;
2.  $x^P(t) = \xi^P e^{at} t$  with  $\xi^P := 1$  is a **particular** solution of the **inhomogeneous** equation  $\dot{x}(t) - ax(t) = e^{at}$ .

## Non-Degenerate Case When $\mu \neq a$

In the **non-degenerate case** when  $\mu \neq a$ , one has

$$(\mu - a) \int_0^t e^{(\mu-a)u} du = \left|_0^t e^{(\mu-a)u} = e^{(\mu-a)t} - 1$$

So the solution is

$$x(t) = e^{at} \left[ x(0) + \frac{e^{(\mu-a)t} - 1}{\mu - a} \right] = e^{at} x(0) + \frac{e^{\mu t} - e^{at}}{\mu - a}$$

Again, this solution can be written as  $x(t) = x^H(t) + x^P(t)$  where:

1.  $x^H(t) = \xi^H e^{at}$  with  $\xi^H := x(0) - 1/(\mu - a)$  is a solution of the **homogeneous** equation  $\dot{x}(t) - ax(t) = 0$ ;
2.  $x^P(t) = \xi^P e^{\mu t}$  with  $\xi^P := 1/(\mu - a)$  is a **particular** solution of the **inhomogeneous** equation  $\dot{x}(t) - ax(t) = e^{\mu t}$ .

## Second Special Case

Another interesting special case occurs when  $f(t) = t^r e^{\mu t}$  for some  $r \in \mathbb{N}$ .

Then the solution  $x(t) = e^{at}x(0) + \int_0^t e^{a(t-u)}f(u)du$  becomes

$$x(t) = e^{at}x(0) + \int_0^t e^{a(t-u)}u^r e^{\mu u}du = e^{at} \left[ x(0) + \int_0^t u^r e^{(\mu-a)u}du \right]$$

In the **degenerate case** when  $\mu = a$ , the solution collapses to

$$x(t) = e^{at} [x(0) + \int_0^t (r+1)^{-1} u^{r+1}] = e^{at} [x(0) + (r+1)^{-1} t^{r+1}]$$

This solution can be written as  $x(t) = x^H(t) + x^P(t)$  where:

1.  $x^H(t) = \xi^H e^{at}$  with  $\xi^H := x(0)$  is a solution of the **homogeneous** equation  $\dot{x}(t) - ax(t) = 0$ ;
2.  $x^P(t) = \xi^P e^{at} t^{r+1}$  with  $\xi^P := (r+1)^{-1}$  is a **particular** solution of the **inhomogeneous** equation  $\dot{x}(t) - ax(t) = t^r e^{at}$ .

## Non-Degenerate Case When $\mu \neq a$

In the **non-degenerate case** when  $\mu \neq a$ , the solution is

$$x(t) = e^{at} \left[ x(0) + \int_0^t u^r e^{(\mu-a)u} du \right] = e^{at} [x(0) + I_r(t)]$$

where  $I_r(t) := \int_0^t u^r e^{(\mu-a)u} du$ .

In particular,  $I_0(t) = \int_0^t e^{(\mu-a)u} du = (\mu - a)^{-1} [e^{(\mu-a)t} - 1]$ .

Integrating by parts gives the first-order linear difference equation

$$\begin{aligned} I_r(t) &= \int_0^t u^r e^{(\mu-a)u} du \\ &= (\mu - a)^{-1} \Big|_0^t u^r e^{(\mu-a)u} - r(\mu - a)^{-1} \int_0^t u^{r-1} e^{(\mu-a)u} du \\ &= (a - \mu)^{-1} [r I_{r-1}(t) - t^r e^{(\mu-a)t}] \end{aligned}$$

## Solving the First-Order Linear Difference Equation

Let us divide each side of the difference equation

$$l_r(t) = (a - \mu)^{-1} \left[ r l_{r-1}(t) - t^r e^{(\mu-a)t} \right]$$

by the “summing factor”  $\prod_{k=1}^r (a - \mu)^{-1} = r!(a - \mu)^{-r}$  to get

$$\begin{aligned} J_r(t) &:= \frac{1}{r!} (a - \mu)^r l_r(t) \\ &= \frac{1}{r!} \left[ r(a - \mu)^{r-1} l_{r-1}(t) - (a - \mu)^{-1} t^r e^{(\mu-a)t} \right] \\ &= \frac{1}{(r-1)!} (a - \mu)^{r-1} l_{r-1}(t) - \frac{1}{r!} (a - \mu)^{r-1} t^r e^{(\mu-a)t} \\ &= J_{r-1}(t) - \frac{1}{r!} (a - \mu)^{r-1} t^r e^{(\mu-a)t} \end{aligned}$$

This obviously implies that

$$J_r(t) = J_0(t) - \sum_{k=1}^r \frac{1}{k!} (a - \mu)^{k-1} t^k e^{(\mu-a)t}$$

## Solving the Differential Equation

Because  $J_0(t) = I_0(t) = (\mu - a)^{-1}[e^{(\mu-a)t} - 1]$ , this implies that

$$J_r(t) = (\mu - a)^{-1}[e^{(\mu-a)t} - 1] - \sum_{k=1}^r \frac{1}{k!} (a - \mu)^{k-1} t^k e^{(\mu-a)t}$$

But  $J_r(t) = \frac{1}{r!} (a - \mu)^r I_r(t)$ , so

$$\begin{aligned} I_r(t) &:= r!(a - \mu)^{-r} J_r(t) \\ &= -r!(a - \mu)^{-r-1} [e^{(\mu-a)t} - 1] \\ &\quad - \sum_{k=1}^r \frac{r!}{k!} (a - \mu)^{k-r-1} t^k e^{(\mu-a)t} \end{aligned}$$

Then

$$\begin{aligned} x(t) &= e^{at} [x(0) + I_r(t)] \\ &= e^{at} \left[ x(0) + r!(a - \mu)^{-r-1} \right. \\ &\quad \left. - r!(a - \mu)^{-r-1} e^{\mu t} \left[ 1 + \sum_{k=1}^r \frac{1}{k!} (a - \mu)^k t^k \right] \right] \end{aligned}$$

## Particular and General Solution

For the equation  $\dot{x}(t) - ax(t) = t^r e^{\mu t}$  with  $\mu \neq a$ , the solution

$$x(t) = e^{at} \left[ x(0) + r!(a - \mu)^{-r-1} \right. \\ \left. - r!(a - \mu)^{-r-1} e^{\mu t} \left[ 1 + \sum_{k=1}^r \frac{1}{k!} (a - \mu)^k t^k \right] \right]$$

can be written as  $x(t) = x^H(t) + x^P(t)$  where:

1.  $x^H(t) = \xi^H e^{at}$  with  $\xi^H := x(0) + r!(a - \mu)^{-r-1}$   
is a solution of the **homogeneous** equation  $\dot{x}(t) - ax(t) = 0$ ;
2.  $x^P(t) = \xi^P(t) e^{\mu t}$ , where the polynomial

$$t \mapsto \xi^P(t) := -r!(a - \mu)^{-r-1} \left[ 1 + \sum_{k=1}^r \frac{1}{k!} (a - \mu)^k t^k \right]$$

of **degree**  $r$  in  $t$  is a **particular** solution  
of the **inhomogeneous** equation  $\dot{x}(t) - ax(t) = t^r e^{\mu t}$ .

# Method of Undetermined Coefficients

A practical issue is finding what polynomial

$$t \mapsto \xi^P(t) = \sum_{k=0}^r \xi_k t^k$$

of degree  $r$  (the power of  $t$  on the right-hand side)

makes  $\xi^P(t)e^{\mu t}$  a particular solution

of the inhomogeneous differential equation  $\dot{x}(t) - ax(t) = t^r e^{\mu t}$ .

The coefficients  $(\xi_0, \xi_1, \dots, \xi_r)$  of the polynomial  $t \mapsto \xi^P(t)$  are **undetermined** till we choose them

so that they make  $\xi^P(t)e^{\mu t}$  satisfy the differential equation.

## Remark

*Choosing  $r$  too low may exclude the solution you want.*

*On the other hand, choosing  $r$  too high just results in finding that the extra terms are zero and so could have been omitted.*

## Determining the Undetermined Coefficients

For  $x^P(t) = e^{\mu t} \sum_{k=0}^r \xi_k t^k$  to solve  $\dot{x}(t) - ax(t) = t^r e^{\mu t}$ , we need

$$\begin{aligned} t^r e^{\mu t} &= \mu e^{\mu t} \sum_{k=0}^r \xi_k t^k + e^{\mu t} \sum_{k=1}^r \xi_k k t^{k-1} - a e^{\mu t} \sum_{k=0}^r \xi_k t^k \\ &= (\mu - a) e^{\mu t} \xi_r t^r + e^{\mu t} \sum_{k=0}^{r-1} [(\mu - a)\xi_k + \xi_{k+1}(k+1)] t^k \end{aligned}$$

First consider the **non-degenerate case**  $\mu \neq a$ .

For  $k = r$ , this implies that  $(\mu - a)\xi_r = 1$ , so  $\xi_r = (\mu - a)^{-1}$ .

For  $k = 0, 1, \dots, r-1$ , it implies that  $(\mu - a)\xi_k + \xi_{k+1}(k+1) = 0$  or that  $\xi_k = (a - \mu)^{-1}(k+1)\xi_{k+1}$ , and so

$$\begin{aligned} \xi_k &= \left[ \prod_{j=k}^{r-1} (a - \mu)^{-1}(j+1) \right] \xi_r \\ &= \frac{r!}{k!} (a - \mu)^{k-r} \xi_r = -\frac{r!}{k!} (a - \mu)^{k-r+1} \end{aligned}$$

This matches our previous answer.

## Degenerate Case

In the **degenerate case** when  $\mu = a$ , the method of undetermined coefficients explained on the previous slides does not work.

Instead, to solve  $\dot{x}(t) - ax(t) = t^r e^{at}$ , we introduce the new variable  $y(t) = e^{-at}x(t)$ .

Then  $\dot{y}(t) = e^{-at}[\dot{x}(t) - ax(t)] = e^{-at}t^r e^{at} = t^r$ .

The solution to this differential equation is  $y(t) = y(0) + \int_0^t u^r du = y(0) + (r+1)^{-1}t^{r+1}$ .

The solution to the original differential equation is therefore  $x(t) = e^{at}y(t) = e^{at} [x(0) + (r+1)^{-1}t^{r+1}]$ .

The polynomial in  $t$  that occurs in this solution is now of degree  $r+1$  rather than  $r$ .

# Main Theorem

## Theorem

Consider the *inhomogeneous* first-order linear differential equation

$$\dot{x}(t) - ax(t) = t^r e^{\mu t}, \text{ where } a \neq 0 \text{ and } r \in \mathbb{Z}_+.$$

There exists a *particular solution* of the form  $x^P(t) = Q(t) e^{\mu t}$  where the function  $t \mapsto Q(t)$  is a polynomial in  $t$  of degree:

- ▶  $r$  in the regular case when  $\mu \neq a$ ;
- ▶  $r + 1$  in the degenerate case when  $\mu = a$ .

The *general solution* takes the form  $x(t) = x^P(t) + x^C(t)$  where:

- ▶  $x^P(t)$  is any particular solution;
- ▶  $x^C(t)$  is any member of the one-dimensional linear space of *complementary solutions* to the corresponding *homogeneous* equation  $\dot{x}(t) - ax(t) = 0$ .

# Lecture Outline

## First-Order Differential Equations in One Variable

Introduction

Picard's Method

General First-Order Affine Equation

Constant and Undetermined Coefficients

**Stability in the Autonomous Case**

## Second-Order Differential Equations in One Variable

Introduction

The Inhomogeneous Equation

The Method of Undetermined Coefficients

Stability

## First-Order Multivariable Differential Equations

Introduction

Prominent Examples and Stability Conditions

Autonomous Nonlinear Equations in Many Variables

Existence and Uniqueness Theorem

# The Autonomous Case

The **autonomous case** occurs when the first-order affine equation takes the form

$$\dot{x} = ax + b$$

with the right-hand side independent of  $t$ .

The **steady state** at which  $\dot{x}(t) = 0$  occurs when  $ax + b = 0$ , and so at  $x^* := -b/a$ .

Then the **deviation**  $y(t) := x(t) - x^*$  of  $x(t)$  from the steady state  $x^*$  satisfies the homogeneous equation

$$\dot{y}(t) = \dot{x}(t) = ax(t) + b = a[y(t) + x^*] + b = ay(t)$$

Hence  $y(t) = e^{at}y(0)$ , implying that  $x(t) = x^* + e^{at}[x(0) - x^*]$ .

# Stability

The steady state  $x^* := -b/a$  is **stable** just in case, for all  $x(0)$ , the solution  $x(t) = x^* + e^{at}[x(0) - x^*]$  satisfies  $x(t) \rightarrow x^*$  as  $t \rightarrow \infty$ .

A necessary and sufficient condition for stability is obviously that  $a < 0$ .

# Lecture Outline

## First-Order Differential Equations in One Variable

Introduction

Picard's Method

General First-Order Affine Equation

Constant and Undetermined Coefficients

Stability in the Autonomous Case

## Second-Order Differential Equations in One Variable

**Introduction**

The Inhomogeneous Equation

The Method of Undetermined Coefficients

Stability

## First-Order Multivariable Differential Equations

Introduction

Prominent Examples and Stability Conditions

Autonomous Nonlinear Equations in Many Variables

Existence and Uniqueness Theorem

## Second-Order Equations with Constant Coefficients

A general **second-order** differential equation takes the form

$$\ddot{x}(t) = F(\dot{x}(t), x(t), t)$$

To obtain a unique solution (if any solution exists), one typically needs two **initial conditions** such as  $x(s) = x_s$  and  $\dot{x}(s) = \dot{x}_s$  at an **initial time**  $s$ .

The equation is **autonomous** just in case it takes the form  $\ddot{x}(t) = F(\dot{x}(t), x(t))$ , with  $F$  independent of  $t$ .

The equation is **linear** just in case it takes the form  $\ddot{x}(t) + a(t)\dot{x}(t) + b(t)x(t) = 0$ , with  $F$  linear in  $(\dot{x}(t), x(t))$ .

The equation is linear with **constant coefficients** just in case it takes the form  $\ddot{x}(t) + a\dot{x}(t) + bx(t) = 0$ .

## Characteristic Equation

We know that the first-order equation  $\dot{x}(t) + ax(t) = 0$  has a solution of the form  $x(t) = x(0)e^{\lambda t}$  where  $\lambda$  solves the characteristic equation  $\lambda + a = 0$ .

So we look for solutions of the form  $x(t) = \xi e^{\lambda t}$  to the second-order equation  $\ddot{x}(t) + a\dot{x}(t) + bx(t) = 0$ .

Note that when  $x(t) = \xi e^{\lambda t}$ , then  $\dot{x}(t) = \lambda \xi e^{\lambda t}$  and  $\ddot{x}(t) = \lambda^2 \xi e^{\lambda t}$ .

So  $x(t) = \xi e^{\lambda t}$  is a **non-trivial** solution (with  $\xi \neq 0$ ) if and only if

$$0 = \lambda^2 \xi e^{\lambda t} + a \lambda \xi e^{\lambda t} + b \xi e^{\lambda t} = (\lambda^2 + a\lambda + b) \xi e^{\lambda t}$$

and so, given that  $\xi e^{\lambda t} \neq 0$ , if and only if  $\lambda$  is a root of the characteristic equation  $\lambda^2 + a\lambda + b = 0$ .

# Characteristic Equation for an Equation of Order $n$

## Definition

A **homogeneous linear** differential equation of order  $n$  with constant coefficients takes the form

$$\sum_{k=0}^n a_k \frac{d^k}{dt^k} x(t) = 0$$

Choose  $n$  so that the coefficient of the  $n$  derivative satisfies  $a_n \neq 0$ , and so can be normalized to take the value  $a_n = 1$ .

## Remark

*A similar technique based on roots of the characteristic equation applies to this  $n$ th order equation.*

*It implies that  $x(t) = \xi e^{\lambda t}$  is a non-trivial solution if and only if  $\lambda$  is a root of the characteristic equation*

$$\sum_{k=0}^n a_k \lambda^k = 0$$

## Characteristic Roots of a Second-Order Equation

Consider the second-order equation  $\ddot{x} + a\dot{x} + b = 0$ .

One can factorize the quadratic function  $q(\lambda) := \lambda^2 + a\lambda + b$  as  $q(\lambda) := (\lambda - \lambda_1)(\lambda - \lambda_2)$

where  $\lambda_1$  and  $\lambda_2$  are the two roots of the equation  $q(\lambda) = 0$ .

As with the corresponding discussion

of second-order difference equations, there are three cases:

1. in case  $a^2 > 4b$ , there are two distinct real roots  $\lambda_1$  and  $\lambda_2$  given by  $\lambda_{1,2} = -\frac{1}{2}a \pm \frac{1}{2}\sqrt{a^2 - 4b}$ .
2. in case  $a^2 < 4b$ , there are two complex conjugate roots given by  $\lambda_{1,2} = -\frac{1}{2}a \pm \frac{1}{2}i\sqrt{4b - a^2}$ .
3. in case  $a^2 = 4b$ , there are two coincident real roots given by  $\lambda = -\frac{1}{2}a = \sqrt{b}$ .

## Case 1: Two Distinct Real Roots

In this case  $a^2 > 4b$ , when the two characteristic roots are  $\lambda_{1,2} = -\frac{1}{2}a \pm \frac{1}{2}\sqrt{a^2 - 4b}$ .

Because  $\lambda_1 \neq \lambda_2$ , one has

$$\begin{vmatrix} e^{\lambda_1 0} & e^{\lambda_2 0} \\ e^{\lambda_1 1} & e^{\lambda_2 1} \end{vmatrix} = \begin{vmatrix} 1 & 1 \\ e^{\lambda_1} & e^{\lambda_2} \end{vmatrix} = e^{\lambda_2} - e^{\lambda_1} \neq 0$$

and so  $e^{\lambda_1 t}$  and  $e^{\lambda_2 t}$  are two linearly independent solutions.

So in this case the homogeneous equation  $\ddot{x} + a\dot{x} + b = 0$  has the general solution

$$x(t) = Ae^{\lambda_1 t} + Be^{\lambda_2 t}$$

## Case 2: Two Complex Conjugate Roots, I

In case  $a^2 < 4b$  the two characteristic roots are the complex conjugates  $\lambda_{1,2} = -\frac{1}{2}a \pm i\theta$ , with  $\theta := \frac{1}{2}\sqrt{4b - a^2}$ .

Then  $x(t) = e^{\lambda_1 t} = e^{-\frac{1}{2}at} e^{i\theta t} = e^{-\frac{1}{2}at} (\cos \theta t + i \sin \theta t)$   
and  $x(t) = e^{\lambda_2 t} = e^{-\frac{1}{2}at} e^{-i\theta t} = e^{-\frac{1}{2}at} (\cos \theta t - i \sin \theta t)$   
are two different solutions, where  $\theta \neq 0$ .

For any  $t$  such that  $\sin \theta t \neq 0$ , one has

$$\begin{vmatrix} e^{\lambda_1 0} & e^{\lambda_2 0} \\ e^{\lambda_1 t} & e^{\lambda_2 t} \end{vmatrix} = \begin{vmatrix} 1 & 1 \\ e^{\lambda_1 t} & e^{\lambda_2 t} \end{vmatrix} = e^{\lambda_2 t} - e^{\lambda_1 t} = -2e^{-\frac{1}{2}at} i \sin \theta t \neq 0$$

It follows that  $e^{\lambda_1 t}$  and  $e^{\lambda_2 t}$  are two linearly independent solutions in the complex plane  $\mathbb{C}$ .

## Case 2: Two Complex Conjugate Roots, II

Focusing on solutions in the real line  $\mathbb{R}$ ,  
we can consider  $e^{-\frac{1}{2}at} \cos \theta t$  and  $e^{-\frac{1}{2}at} \sin \theta t$ .

Again, for any  $t$  such that  $\sin \theta t \neq 0$ , one has

$$\begin{aligned} \begin{vmatrix} e^{-\frac{1}{2}a0} \cos \theta 0 & e^{-\frac{1}{2}a0} \sin \theta 0 \\ e^{-\frac{1}{2}at} \cos \theta t & e^{-\frac{1}{2}at} \sin \theta t \end{vmatrix} &= \begin{vmatrix} 1 & 0 \\ -\frac{1}{2}at \cos \theta t & e^{-\frac{1}{2}at} \sin \theta t \end{vmatrix} \\ &= e^{-\frac{1}{2}at} \sin \theta t \neq 0 \end{aligned}$$

It follows that  $e^{-\frac{1}{2}at} \cos \theta t$  and  $e^{-\frac{1}{2}at} \sin \theta t$  are two linearly independent real-valued solutions in the complex plane  $\mathbb{C}$ .

The general solution of the homogeneous equation  
is  $x = e^{-\frac{1}{2}at}(A \cos \theta t + B \sin \theta t)$ .

## Case 3: Two Coincident Real Roots

In this case  $a^2 = 4b$ , and so

$$q(\lambda) = \lambda^2 + a\lambda + b = (\lambda + \frac{1}{2}a)^2 = (\lambda - \sqrt{b})^2$$

The homogeneous equation  $\ddot{x}(t) + a\dot{x}(t) + bx(t) = 0$  has one solution given by  $x = e^{\lambda t}$  where  $\lambda = -\frac{1}{2}a = \sqrt{b}$ .

To find a second linearly independent solution, introduce the new variable  $y(t) := e^{-\lambda t}x(t)$ .

Then  $\dot{y}(t) = e^{-\lambda t}\dot{x}(t) - \lambda e^{-\lambda t}x(t)$  and so, when  $x = e^{\lambda t}$ , one has

$$\begin{aligned}\ddot{y}(t) &= e^{-\lambda t}\ddot{x}(t) - 2\lambda e^{-\lambda t}\dot{x}(t) + \lambda^2 e^{-\lambda t}x(t) \\ &= e^{-\lambda t}[\ddot{x}(t) - 2\lambda\dot{x}(t) + \lambda^2x(t)] \\ &= e^{-\lambda t}[\lambda^2 e^{\lambda t} - 2\lambda \cdot \lambda e^{\lambda t} + \lambda^2 e^{\lambda t}] = 0\end{aligned}$$

The obvious general solution to  $\ddot{y}(t) = 0$  satisfies  $\dot{y}(t) = \text{constant}$  and so  $y(t) = A + Bt = e^{-\lambda t}x(t)$ .

Hence  $x(t) = (A + Bt)e^{\lambda t}$  is the general solution.

# Lecture Outline

## First-Order Differential Equations in One Variable

Introduction

Picard's Method

General First-Order Affine Equation

Constant and Undetermined Coefficients

Stability in the Autonomous Case

## Second-Order Differential Equations in One Variable

Introduction

The Inhomogeneous Equation

The Method of Undetermined Coefficients

Stability

## First-Order Multivariable Differential Equations

Introduction

Prominent Examples and Stability Conditions

Autonomous Nonlinear Equations in Many Variables

Existence and Uniqueness Theorem

# The Inhomogeneous Equation

Consider next the inhomogeneous equation

$$\ddot{x}(t) + a(t)\dot{x}(t) + b(t)x(t) = f(t)$$

with a non-zero forcing term on the right-hand side.

Suppose that  $y(t)$  and  $z(t)$  are both solutions, implying that

$$\begin{aligned} \ddot{y}(t) + a(t)\dot{y}(t) + b(t)y(t) &= f(t) \\ \text{and } \ddot{z}(t) + a(t)\dot{z}(t) + b(t)z(t) &= f(t) \end{aligned}$$

Subtracting the second equation from the first tells us that the function  $x_H(t) := y(t) - z(t)$  is a solution of the corresponding homogeneous equation  $\ddot{x}(t) + a(t)\dot{x}(t) + b(t)x(t) = 0$ .

So the **general** solution of  $\ddot{x}(t) + a(t)\dot{x}(t) + b(t)x(t) = f(t)$  is the sum  $x_G(t) = x_P(t) + x_H(t)$  of:

- ▶ any **particular** solution  $x_P(t)$  of the inhomogeneous equation;
- ▶ any function  $x_H(t)$  in the two dimensional linear space of solutions to the homogeneous equation.

## Linearity in the Forcing Term, I

### Theorem

Suppose that  $x^P(t)$  and  $y^P(t)$  are particular solutions of the two respective differential equations

$$\begin{aligned}\ddot{x}(t) + a(t)\dot{x}(t) + b(t)x(t) &= d(t) \\ \text{and } \ddot{y}(t) + a(t)\dot{y}(t) + b(t)y(t) &= e(t)\end{aligned}$$

Then, for any scalars  $\alpha$  and  $\beta$ , a particular solution of the equation

$$\ddot{z}(t) + a(t)\dot{z}(t) + b(t)z(t) = f(t) = \alpha d(t) + \beta e(t) \quad (*)$$

is the linear combination  $z^P(t) := \alpha x^P(t) + \beta y^P(t)$ .

### Proof.

Verify the claimed solution

by inserting the specified linear combination  $z^P(t)$ , together with its first two derivatives  $\dot{z}^P(t)$  and  $\ddot{z}^P(t)$ , into the differential equation (\*). □

## Linearity in the Forcing Term, II

Consider the equation  $\ddot{x}(t) + a(t)\dot{x}(t) + b(t)x(t) = f(t)$  whose forcing term  $f(t)$  is a linear combination  $\sum_{k=1}^n \alpha_k f^k(t)$  of  $n$  forcing terms.

The theorem implies that a particular solution is the corresponding linear combination  $\sum_{k=1}^n \alpha_k x^{Pk}(t)$  of particular solutions  $x^{Pk}(t)$  to the respective  $n$  equations

$$\ddot{x}(t) + a(t)\dot{x}(t) + b(t)x(t) = f^k(t) \quad (k = 1, 2, \dots, n)$$

# Lecture Outline

## First-Order Differential Equations in One Variable

Introduction

Picard's Method

General First-Order Affine Equation

Constant and Undetermined Coefficients

Stability in the Autonomous Case

## Second-Order Differential Equations in One Variable

Introduction

The Inhomogeneous Equation

The Method of Undetermined Coefficients

Stability

## First-Order Multivariable Differential Equations

Introduction

Prominent Examples and Stability Conditions

Autonomous Nonlinear Equations in Many Variables

Existence and Uniqueness Theorem

## A Newtonian Example, I

Newton's law: force = mass  $\times$  acceleration.

A force of 1 Newton, by definition, accelerates a mass of 1 kilogram at the rate of 1 metre per second per second.

So we consider the equation  $\ddot{x}(t) = f(t)$  whose solution  $t \mapsto x(t)$  is the position (in one dimension) of a 1 kilogram weight that has been subjected to a force function  $t \mapsto f(t)$ .

Integrating once gives us the equation  $\dot{x}(t) = \dot{x}(0) + \int_0^t f(u)du$ .

Integrating a second time gives us the solution

$$\begin{aligned}x(t) &= x(0) + \int_0^t \dot{x}(v)dv = x(0) + \int_0^t [\dot{x}(0) + \int_0^v f(u)du] dv \\ &= x(0) + \dot{x}(0)t + \int_0^t [\int_0^v f(u)du] dv\end{aligned}$$

Note that  $x(0) + \dot{x}(0)t$  solves the homogeneous equation  $\ddot{x}(t) = 0$ , whereas the iterated double integral  $\int_0^t [\int_0^v f(u)du] dv$  is a particular solution.

# An Important Theorem on Iterated Double Integrals, I

## Theorem

For any integrable function  $(x, y) \mapsto \phi(x, y) \in \mathbb{R}$  defined on the square domain  $[a, b] \times [a, b] \subset \mathbb{R}^2$ , one has

$$\int_a^b \left[ \int_a^y \phi(x, y) dx \right] dy = \int_a^b \left[ \int_x^b \phi(x, y) dy \right] dx$$

## Proof.

Define the **indicator function**  $1_{x \leq y}(x, y) := \begin{cases} 1 & \text{if } x \leq y \\ 0 & \text{if } x > y \end{cases}$ . Then

$$\begin{aligned} \int_a^b \left[ \int_a^y \phi(x, y) dx \right] dy &= \int_a^b \left[ \int_a^b 1_{x \leq y}(x, y) \phi(x, y) dx \right] dy \\ \int_a^b \left[ \int_x^b \phi(x, y) dy \right] dx &= \int_a^b \left[ \int_a^b 1_{x \leq y}(x, y) \phi(x, y) dy \right] dx \end{aligned}$$

But both right-hand sides equal  $\int_a^b \int_a^b 1_{x \leq y}(x, y) \phi(x, y) dx dy$ .  $\square$

## An Important Theorem on Iterated Double Integrals, II

An alternative simple proof involves noticing that the two integrals

$$\int_a^b \left[ \int_a^y \phi(x, y) dx \right] dy \quad \text{and} \quad \int_a^b \left[ \int_x^b \phi(x, y) dy \right] dx$$

are simply two different ways of writing the integral  $\iint_T \phi(x, y) dx dy$  of the function  $\phi$  of two variables over the isosceles right-angled triangle

$$T := \{(x, y) \in [a, b] \times [a, b] \subset \mathbb{R}^2 \mid x \leq y\}$$

Note that  $T$  consists of points above and to the left of the diagonal that joins the two corner points  $(a, a)$  and  $(b, b)$  of the square  $[a, b] \times [a, b]$ .

The set  $T$  is also the convex hull of the three points  $(a, a)$ ,  $(a, b)$  and  $(b, b)$ .

## A Newtonian Example, II

Reversing the order of integration allows the particular solution in the form of the iterated double integral  $\int_0^t \left[ \int_0^v f(u) du \right] dv$  to be rewritten as

$$\int_0^t \left[ \int_u^t f(u) dv \right] du = \int_0^t \left[ \int_u^t 1 dv \right] f(u) du = \int_0^t (t - u) f(u) du$$

Ultimately, then, one has

$$x(t) = x(0) + \dot{x}(0)t + \int_0^t (t - u) f(u) du$$

## Linear Equation with Constant Coefficients, I

Next, consider the equation  $\ddot{x}(t) + a\dot{x}(t) + bx(t) = f(t)$  where the coefficients  $a$  of  $\dot{x}(t)$  and  $b$  of  $x(t)$  have both become **constants**, with  $b \neq 0$ .

Consider the quadratic function  $q(\lambda) := \lambda^2 + a\lambda + b$  that appears in the characteristic equation  $\lambda^2 + a\lambda + b = 0$ .

One can factorize it as

$$q(\lambda) = \lambda^2 + a\lambda + b = (\lambda - \lambda_1)(\lambda - \lambda_2)$$

where  $\lambda_1$  and  $\lambda_2$  are the two roots of the equation  $q(\lambda) = 0$ .

Recall that  $\lambda_1 + \lambda_2 = -a$  and  $\lambda_1\lambda_2 = b$ .

Define the new variable  $y(t) := \dot{x}(t) - \lambda_1 x(t)$ .

Note that, if we could find the function  $t \mapsto y(t)$ , then we would have

$$x(t) = e^{\lambda_1 t} x(0) + \int_0^t e^{\lambda_1(t-u)} y(u) du$$

## Linear Equation with Constant Coefficients, II

We are considering the equation  $\ddot{x}(t) + a\dot{x}(t) + bx(t) = f(t)$ , with  $b \neq 0$ .

We have introduced the new variable  $y(t) := \dot{x}(t) - \lambda_1 x(t)$ , implying that  $x(t) = e^{\lambda_1 t} x(0) + \int_0^t e^{\lambda_1(t-u)} y(u) du$ .

But the characteristic roots satisfy  $\lambda^2 + a\lambda + b = (\lambda - \lambda_1)(\lambda - \lambda_2)$ , implying that  $\lambda_1 + \lambda_2 = -a$  and  $\lambda_1 \lambda_2 = b$ , and so

$$\begin{aligned} \dot{y}(t) - \lambda_2 y(t) &= \ddot{x}(t) - \lambda_1 \dot{x}(t) - \lambda_2 \dot{x}(t) + \lambda_1 \lambda_2 x(t) \\ &= \ddot{x}(t) + a\dot{x}(t) + bx(t) \end{aligned}$$

Hence  $y(t)$  satisfies the first-order equation  $\dot{y}(t) - \lambda_2 y(t) = f(t)$  whose solution is

$$y(t) = e^{\lambda_2 t} y(0) + \int_0^t e^{\lambda_2(t-v)} f(v) dv$$

## Linear Equation with Constant Coefficients, III

Substituting  $y(t) = e^{\lambda_2 t} y(0) + \int_0^t e^{\lambda_2(t-v)} f(v) dv$   
in the expression  $x(t) = e^{\lambda_1 t} x(0) + \int_0^t e^{\lambda_1(t-u)} y(u) du$  gives

$$\begin{aligned} x(t) &= e^{\lambda_1 t} x(0) + \int_0^t e^{\lambda_1(t-u)} y(u) du \\ &= e^{\lambda_1 t} x(0) + \int_0^t e^{\lambda_1(t-u)} \left[ e^{\lambda_2 u} y(0) + \int_0^u e^{\lambda_2(u-v)} f(v) dv \right] du \end{aligned}$$

## Linear Equation with Constant Coefficients, IV

This form of the solution is the sum of the following two parts:

1. the **complementary** solution

$$\begin{aligned}t \mapsto x^C(t) &:= e^{\lambda_1 t} x(0) + y(0) \int_0^t e^{\lambda_1(t-u)} e^{\lambda_2 u} du \\ &= e^{\lambda_1 t} \left[ x(0) + y(0) \int_0^t e^{(\lambda_2 - \lambda_1)u} du \right]\end{aligned}$$

to the homogeneous equation  $\ddot{x}(t) + a\dot{x}(t) + bx(t) = 0$ ;

2. a **particular** solution in the form of the iterated double integral

$$t \mapsto x^P(t) := \int_0^t e^{\lambda_1(t-u)} \left[ \int_0^u e^{\lambda_2(u-v)} f(v) dv \right] du$$

to the inhomogeneous equation  $\ddot{x}(t) + a\dot{x}(t) + bx(t) = f(t)$ .

## Degenerate Case

In the degenerate case when  $\lambda_1 = \lambda_2 = \lambda$ ,

1. the **complementary** solution takes the form:

$$\begin{aligned}x^C(t) &= e^{\lambda t}x(0) + y(0) \int_0^t e^{\lambda u} du \\ &= e^{\lambda t} [x(0) + y(0)t]\end{aligned}$$

2. the **particular** solution takes the form:

$$\begin{aligned}x^P(t) &= \int_0^t e^{\lambda(t-u)} \left[ \int_0^u e^{\lambda(u-v)} f(v) dv \right] du \\ &= e^{\lambda t} \int_0^t \left[ \int_0^u e^{-\lambda v} f(v) dv \right] du \\ &= e^{\lambda t} \int_0^t \left[ \int_v^t 1 du \right] e^{-\lambda v} f(v) dv \\ &= \int_0^t (t-v) e^{\lambda(t-v)} f(v) dv\end{aligned}$$

The overall solution is therefore

$$x(t) = e^{\lambda t} \left[ x(0) + y(0)t + \int_0^t (t-v) e^{-\lambda v} f(v) dv \right]$$

## Non-Degenerate Case: Complementary Solution

In the non-degenerate case when  $\lambda_1 \neq \lambda_2$ , the **complementary** solution takes the form

$$\begin{aligned}x^C(t) &= e^{\lambda_1 t} \left[ x(0) + y(0) \int_0^t e^{(\lambda_2 - \lambda_1)u} du \right] \\&= e^{\lambda_1 t} x(0) + \frac{1}{\lambda_2 - \lambda_1} e^{\lambda_1 t} y(0) [e^{(\lambda_2 - \lambda_1)t} - 1] \\&= x(0)e^{\lambda_1 t} + y(0) \frac{e^{\lambda_2 t} - e^{\lambda_1 t}}{\lambda_2 - \lambda_1}\end{aligned}$$

After substituting  $\dot{x}(0) - \lambda_1 x(0)$  for  $y(0)$ , the right-hand side becomes

$$\frac{1}{\lambda_2 - \lambda_1} \left\{ (\lambda_2 - \lambda_1)x(0)e^{\lambda_1 t} + [\dot{x}(0) - \lambda_1 x(0)](e^{\lambda_2 t} - e^{\lambda_1 t}) \right\}$$

and so

$$x^C(t) = \frac{1}{\lambda_2 - \lambda_1} \left[ x(0)(\lambda_2 e^{\lambda_1 t} - \lambda_1 e^{\lambda_2 t}) + \dot{x}(0)(e^{\lambda_2 t} - e^{\lambda_1 t}) \right]$$

## Non-Degenerate Case: Particular Solution

Using our rule for reversing the order of recursive integration, the **particular** solution takes the form

$$\begin{aligned}x^P(t) &= \int_0^t e^{\lambda_1(t-u)} \left[ \int_0^u e^{\lambda_2(u-v)} f(v) dv \right] du \\&= \int_0^t \left[ \int_v^t e^{\lambda_1(t-u)} e^{\lambda_2(u-v)} du \right] f(v) dv \\&= \int_0^t e^{\lambda_1 t - \lambda_2 v} \left[ \int_v^t e^{(\lambda_2 - \lambda_1)u} du \right] f(v) dv \\&= \frac{1}{\lambda_2 - \lambda_1} \int_0^t e^{\lambda_1 t - \lambda_2 v} \left[ e^{(\lambda_2 - \lambda_1)t} - e^{(\lambda_2 - \lambda_1)v} \right] f(v) dv \\&= \frac{1}{\lambda_2 - \lambda_1} \int_0^t \left[ e^{\lambda_2(t-v)} - e^{\lambda_1(t-v)} \right] f(v) dv\end{aligned}$$

## First Special Case

An interesting first special case of the particular solution

$$x^P(t) = \frac{1}{\lambda_2 - \lambda_1} \int_0^t \left[ e^{\lambda_2(t-v)} - e^{\lambda_1(t-v)} \right] f(v) dv$$

occurs when  $f(t)$  is the exponential function  $e^{\mu t}$ , and so

$$x^P(t) = \frac{1}{\lambda_2 - \lambda_1} \int_0^t \left[ e^{\lambda_2(t-v)} - e^{\lambda_1(t-v)} \right] e^{\mu v} dv$$

In the degenerate case when  $\lambda_2 = \mu \neq \lambda_1$ , this reduces to

$$\begin{aligned} x^P(t) &= \frac{1}{\lambda_2 - \lambda_1} \left[ e^{\lambda_2 t} t - e^{\lambda_1 t} \int_0^t e^{(\mu - \lambda_1)v} dv \right] \\ &= \frac{e^{\lambda_2 t} t}{\lambda_2 - \lambda_1} - \frac{e^{\lambda_1 t} (e^{(\lambda_2 - \lambda_1)t} - 1)}{(\lambda_2 - \lambda_1)^2} \\ &= \frac{e^{\lambda_2 t} t}{\lambda_2 - \lambda_1} - \frac{e^{\lambda_2 t} - e^{\lambda_1 t}}{(\lambda_2 - \lambda_1)^2} \end{aligned}$$

## Non-Degenerate Case

In the **non-degenerate case** when  $\lambda_1$ ,  $\lambda_2$  and  $\mu$  are all different, one has the particular solution

$$\begin{aligned}x^P(t) &= \frac{e^{\lambda_2 t}}{\lambda_2 - \lambda_1} \int_0^t e^{(\mu - \lambda_2)v} dv - \frac{e^{\lambda_1 t}}{\lambda_2 - \lambda_1} \int_0^t e^{(\mu - \lambda_1)v} dv \\&= \frac{e^{\lambda_2 t} [e^{(\mu - \lambda_2)t} - 1]}{(\lambda_2 - \lambda_1)(\mu - \lambda_2)} - \frac{e^{\lambda_1 t} [e^{(\mu - \lambda_1)t} - 1]}{(\lambda_2 - \lambda_1)(\mu - \lambda_1)} \\&= \frac{1}{\lambda_2 - \lambda_1} \left( \frac{e^{\mu t} - e^{\lambda_2 t}}{\mu - \lambda_2} - \frac{e^{\mu t} - e^{\lambda_1 t}}{\mu - \lambda_1} \right)\end{aligned}$$

But the multiples of  $e^{\lambda_1 t}$  and  $e^{\lambda_2 t}$  can be incorporated in the complementary solution to the homogeneous equation, so this particular solution can be reduced to

$$\tilde{x}^P(t) = \frac{e^{\mu t}}{\lambda_2 - \lambda_1} \left( \frac{1}{\mu - \lambda_2} - \frac{1}{\mu - \lambda_1} \right) = \frac{e^{\mu t}}{(\mu - \lambda_1)(\mu - \lambda_2)}$$

## Second Special Case

An interesting second special case of the particular solution

$$x^P(t) = \frac{1}{\lambda_2 - \lambda_1} \int_0^t \left[ e^{\lambda_2(t-v)} - e^{\lambda_1(t-v)} \right] f(v) dv$$

occurs when  $f(t)$  is the exponential function  $t^r e^{\mu t}$ , and so

$$x^P(t) = \frac{1}{\lambda_2 - \lambda_1} \int_0^t \left[ e^{\lambda_2(t-v)} - e^{\lambda_1(t-v)} \right] v^r e^{\mu v} dv$$

In the **non-degenerate case** when  $\lambda_1$ ,  $\lambda_2$  and  $\mu$  are all different, this particular solution  $x^P(t)$  becomes

$$\frac{1}{\lambda_2 - \lambda_1} \left[ e^{\lambda_2 t} \int_0^t v^r e^{(\mu - \lambda_2)v} dv - e^{\lambda_1 t} \int_0^t v^r e^{(\mu - \lambda_1)v} dv \right]$$

It follows that  $x^P(t) = P_2(t)e^{\lambda_2 t} - P_1(t)e^{\lambda_1 t}$   
for polynomials  $t \mapsto P_1(t)$  and  $t \mapsto P_2(t)$  of degree  $r$   
whose coefficients are functions of the parameter triple  $(\lambda_1, \lambda_2, \mu)$ .

# Lecture Outline

## First-Order Differential Equations in One Variable

Introduction

Picard's Method

General First-Order Affine Equation

Constant and Undetermined Coefficients

Stability in the Autonomous Case

## Second-Order Differential Equations in One Variable

Introduction

The Inhomogeneous Equation

The Method of Undetermined Coefficients

**Stability**

## First-Order Multivariable Differential Equations

Introduction

Prominent Examples and Stability Conditions

Autonomous Nonlinear Equations in Many Variables

Existence and Uniqueness Theorem

# The Autonomous Equation

Now consider the autonomous equation

$$\ddot{x}(t) + a\dot{x}(t) + bx(t) = c$$

with a constant right-hand side.

There is a constant solution  $x(t) = \bar{x}$   
where  $\bar{x} = c/b$  is the unique steady state.

The new variable  $y(t) := x(t) - \bar{x}$  satisfies  
the homogeneous equation  $\ddot{y}(t) + a\dot{y}(t) + by(t) = 0$ .

The associated characteristic equation is

$$\lambda^2 + a\lambda + b = (\lambda - \lambda_1)(\lambda - \lambda_2) = 0$$

## A Stability Condition

1. In case there are two real characteristic roots

$$\lambda_{1,2} = -\frac{1}{2}a \pm \frac{1}{2}\sqrt{a^2 - 4b}$$

the general solution  $Ae^{\lambda_1 t} + Be^{\lambda_2 t} \rightarrow 0$  as  $t \rightarrow \infty$   
if and only if both  $\lambda_1$  and  $\lambda_2$  are negative.

2. In case there are two complex conjugate characteristic roots

$$\lambda_{1,2} = -\frac{1}{2}a \pm \frac{1}{2}i\sqrt{4b - a^2}$$

one has  $e^{\lambda t} = e^{-\frac{1}{2}at} e^{\pm \frac{1}{2}it\sqrt{4b - a^2}}$ .

The general solution  $Ae^{\lambda_1 t} + Be^{\lambda_2 t} \rightarrow 0$  as  $t \rightarrow \infty$   
iff  $a > 0$ , or iff both  $\lambda_1$  and  $\lambda_2$  have negative real parts.

3. In case there are two coincident real characteristic roots,  
the general solution  $(A + Bt)e^{\lambda t} \rightarrow 0$  as  $t \rightarrow \infty$  iff  $\lambda < 0$ .

All these conditions can be subsumed into one: stability holds  
if and only if each characteristic root has a negative real part.

# Lecture Outline

## First-Order Differential Equations in One Variable

Introduction

Picard's Method

General First-Order Affine Equation

Constant and Undetermined Coefficients

Stability in the Autonomous Case

## Second-Order Differential Equations in One Variable

Introduction

The Inhomogeneous Equation

The Method of Undetermined Coefficients

Stability

## First-Order Multivariable Differential Equations

Introduction

Prominent Examples and Stability Conditions

Autonomous Nonlinear Equations in Many Variables

Existence and Uniqueness Theorem

# Linear Differential Equation in $n$ Variables

A **linear differential equation in  $n$  variables** specifies the time derivative  $\dot{\mathbf{x}}(t)$  of the  $n$ -vector  $\mathbf{x}(t)$  as an affine function  $\mathbf{A}(t)\mathbf{x}(t) + \mathbf{b}(t)$  of  $\mathbf{x}(t)$ .

That is

$$\dot{\mathbf{x}}(t) = \mathbf{A}(t)\mathbf{x}(t) + \mathbf{b}(t)$$

where

- ▶  $t \mapsto \mathbf{A}(t) \in \mathbb{R}^{n \times n}$  is a matrix-valued function of time;
- ▶  $t \mapsto \mathbf{b}(t) \in \mathbb{R}^n$  is a vector-valued function of time.

# Matrix Differentiation

Consider the  $m \times n$  matrix function  $t \mapsto \mathbf{A}(t)$  whose elements  $(a_{ij}(t))_{m \times n}$  are differentiable functions of  $t$ .

For all  $h \neq 0$ , the **Newton quotient** matrix  $\frac{1}{h}[\mathbf{A}(t+h) - \mathbf{A}(t)]$  has elements equal to

the Newton quotients  $\frac{1}{h}(a_{ij}(t+h) - a_{ij}(t))_{m \times n}$  of the matrix  $(a_{ij}(t))_{m \times n}$ .

As  $h \rightarrow 0$ , these converge to the derivatives  $(\frac{d}{dt}a_{ij}(t))_{m \times n}$ .

For this reason, the matrix  $\mathbf{A}(t)$  is said to be **differentiable** with **derivative**  $\dot{\mathbf{A}}(t) = \frac{d}{dt}\mathbf{A}(t)$  whose elements are  $(\frac{d}{dt}a_{ij}(t))_{m \times n}$ .

# Differentiating the Product of Matrices

Suppose that  $t \mapsto \mathbf{A}(t)$  and  $t \mapsto \mathbf{B}(t)$  are differentiable, where each  $\mathbf{A}(t)$  is  $\ell \times m$  and each  $\mathbf{B}(t)$  is  $m \times n$ .

Then  $t \mapsto \mathbf{C}(t) = \mathbf{A}(t)\mathbf{B}(t)$  is well defined as a matrix product with elements given by  $c_{ik}(t) = \sum_{j=1}^m a_{ij}(t)b_{jk}(t)$  whose time derivatives are

$$\dot{c}_{ik}(t) = \sum_{j=1}^m [\dot{a}_{ij}(t)b_{jk}(t) + a_{ij}(t)\dot{b}_{jk}(t)]$$

Hence  $t \mapsto \mathbf{C}(t)$  is differentiable, with  $\dot{\mathbf{C}}(t) = \dot{\mathbf{A}}(t)\mathbf{B}(t) + \mathbf{A}(t)\dot{\mathbf{B}}(t)$ .

# Differentiating the Square of a Square Matrix

Suppose that  $\mathbf{A}(t)$  is an  $n \times n$  matrix for all  $t$ ,  
and that each element is a differentiable function of  $t$ .

Then the square matrix  $\mathbf{A}^2(t)$  is well defined and differentiable,  
with derivative  $\frac{d}{dt}\mathbf{A}^2(t) = \dot{\mathbf{A}}(t)\mathbf{A}(t) + \mathbf{A}(t)\dot{\mathbf{A}}(t)$ .

Unless the matrices  $\dot{\mathbf{A}}(t)$  and  $\mathbf{A}(t)$  happen to commute,  
in the sense that  $\dot{\mathbf{A}}(t)\mathbf{A}(t) = \mathbf{A}(t)\dot{\mathbf{A}}(t)$ ,  
this will **not** be equal to  $2\dot{\mathbf{A}}(t)\mathbf{A}(t)$  or to  $2\mathbf{A}(t)\dot{\mathbf{A}}(t)$ .

## Example

Note that, even if each  $\mathbf{A}(t)$  is square, it may not commute with  $\dot{\mathbf{A}}(t)$ .

For example, when  $\mathbf{A}(t) = \begin{pmatrix} 0 & 1 \\ 1 & t \end{pmatrix}$ , then  $\dot{\mathbf{A}}(t) = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ ,  
implying that  $\mathbf{A}(t)\dot{\mathbf{A}}(t) = \begin{pmatrix} 0 & 1 \\ 0 & t \end{pmatrix} \neq \dot{\mathbf{A}}(t)\mathbf{A}(t) = \begin{pmatrix} 0 & 0 \\ 1 & t \end{pmatrix}$ .

Note that in this example  $\mathbf{A}$  is symmetric; so therefore is  $\dot{\mathbf{A}}$ .  
Hence  $\mathbf{A}(t)\dot{\mathbf{A}}(t) = \mathbf{A}(t)^\top \dot{\mathbf{A}}^\top(t) = [\dot{\mathbf{A}}(t)\mathbf{A}(t)]^\top$ .

Also  $\mathbf{A}^2(t) = \begin{pmatrix} 1 & t \\ t & 1+t^2 \end{pmatrix}$  whose derivative satisfies

$$\frac{d}{dt}\mathbf{A}^2(t) = \dot{\mathbf{A}}(t)\mathbf{A}(t) + \mathbf{A}(t)\dot{\mathbf{A}}(t) = \begin{pmatrix} 0 & 1 \\ 1 & 2t \end{pmatrix}$$

This differs from both  $2\mathbf{A}(t)\dot{\mathbf{A}}(t)$  and  $2\dot{\mathbf{A}}(t)\mathbf{A}(t)$ .

## The Exponential of a Square Matrix

Recall that the exponential function of a scalar is **defined** so that the solution of the differential equation  $\dot{x} = ax$  is  $x(t) = e^{at}x(0)$ .

Similarly, we define the **exponential function of a square matrix** so that the solution of the differential equation system  $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x}$  is  $\mathbf{x}(t) = \exp(\mathbf{A}t)\mathbf{x}(0)$ .

The function  $t \mapsto \exp(\mathbf{A}t)$  is often called the **resolvent**.

Recall that, for a scalar, there is the convergent power series

$$e^{at} = 1 + \frac{1}{1!}at + \frac{1}{2!}(at)^2 + \frac{1}{3!}(at)^3 \dots = \sum_{r=0}^{\infty} \frac{1}{r!}(at)^r$$

with the convention that  $0! = 1$ .

Similarly, for a square matrix, with the convention that  $(\mathbf{A}t)^0 = \mathbf{I}$  one can use a convergent power series to give,

$$\exp(\mathbf{A}t) = \mathbf{I} + \frac{1}{1!}\mathbf{A}t + \frac{1}{2!}(\mathbf{A}t)^2 + \frac{1}{3!}(\mathbf{A}t)^3 \dots = \sum_{r=0}^{\infty} \frac{1}{r!}(\mathbf{A}t)^r$$

## The Exponential of a Diagonal Matrix

Dropping the time argument, it follows that we define

$$\exp(\mathbf{C}) := \mathbf{I} + \frac{1}{1!}\mathbf{C} + \frac{1}{2!}(\mathbf{C})^2 + \frac{1}{3!}(\mathbf{C})^3 \dots = \sum_{r=0}^{\infty} \frac{1}{r!}(\mathbf{C})^r$$

Suppose that  $\mathbf{C}$  is the diagonal matrix  $\mathbf{diag}(c_1, c_2, \dots, c_n) = \mathbf{diag} \mathbf{c}$  where  $\mathbf{c}$  is the vector  $(c_1, c_2, \dots, c_n)$ .

Now, each matrix power  $(\mathbf{diag} \mathbf{c})^r = \mathbf{diag}(c_1^r, c_2^r, \dots, c_n^r)$  as is readily proved by induction on  $r$ .

So, with this notation for the exponential of a matrix, we have

$$\begin{aligned} \exp(\mathbf{C}) &= \sum_{r=0}^{\infty} \frac{1}{r!}\mathbf{C}^r = \sum_{r=0}^{\infty} \frac{1}{r!}\mathbf{diag}(c_1^r, c_2^r, \dots, c_n^r) \\ &= \mathbf{diag}(e^{c_1}, e^{c_2}, \dots, e^{c_n}) \end{aligned}$$

Also, suppose matrix  $\mathbf{C}$  has  $\mathbf{C} = \mathbf{V}\mathbf{\Lambda}\mathbf{V}^{-1}$  as a diagonalization.

Then each matrix power  $\mathbf{C}^r = \mathbf{V}\mathbf{\Lambda}^r\mathbf{V}^{-1}$  implying that  $\exp(\mathbf{C}) = \mathbf{V}\exp(\mathbf{\Lambda})\mathbf{V}^{-1}$ .

## Integrating and Differentiating an Exponential Matrix

From the definition  $\exp(\mathbf{A}s) = \sum_{r=0}^{\infty} \frac{1}{r!} (\mathbf{A}s)^r$ ,  
either post- or premultiplying by  $\mathbf{A}$  and then integrating gives

$$\int_0^t \exp(\mathbf{A}s) \mathbf{A} ds = \int_0^t \mathbf{A} \exp(\mathbf{A}s) ds = \int_0^t \sum_{r=0}^{\infty} \frac{1}{r!} \mathbf{A}^{r+1} s^r ds$$

Next, integrating term by term, the last expression becomes

$$\sum_{r=0}^{\infty} \frac{1}{r!} \mathbf{A}^{r+1} \int_0^t s^r ds = \sum_{r=0}^{\infty} \frac{1}{r!} \mathbf{A}^{r+1} \cdot \left| \frac{1}{r+1} s^{r+1} \right|_0^t$$

Simplifying converts this to

$$\sum_{r=0}^{\infty} \frac{1}{(r+1)!} \mathbf{A}^{r+1} t^{r+1} = \sum_{r=1}^{\infty} \frac{1}{r!} \mathbf{A}^r t^r = \exp(\mathbf{A}t) - \mathbf{I}$$

So  $\int_0^t \exp(\mathbf{A}s) \mathbf{A} ds = \int_0^t \mathbf{A} \exp(\mathbf{A}s) ds = \exp(\mathbf{A}t) - \mathbf{I}$ ,  
implying that

$$\frac{d}{dt} \exp(\mathbf{A}t) = \mathbf{A} \exp(\mathbf{A}t) = \exp(\mathbf{A}t) \mathbf{A}$$

## Affine Equation in $n$ Variables

Consider what happens when we multiply each side of the non-homogeneous **affine** equation  $\dot{\mathbf{x}}(t) - \mathbf{A}\mathbf{x}(t) = \mathbf{b}(t)$  by the **matrix integrating factor**  $\exp(-\mathbf{A}t)$ .

Because the product rule of differentiation applies to matrices,

$$\begin{aligned}\frac{d}{dt} [\exp(-\mathbf{A}t) \mathbf{x}(t)] &= \exp(-\mathbf{A}t) \dot{\mathbf{x}}(t) + \frac{d}{dt} [\exp(-\mathbf{A}t)] \mathbf{x}(t) \\ &= \exp(-\mathbf{A}t) \dot{\mathbf{x}}(t) - \exp(-\mathbf{A}t) \mathbf{A} \mathbf{x}(t) \\ &= \exp(-\mathbf{A}t) \mathbf{b}(t)\end{aligned}$$

if and only if  $\mathbf{x}(t)$  solves the equation  $\dot{\mathbf{x}}(t) - \mathbf{A}\mathbf{x}(t) = \mathbf{b}(t)$ .

Hence  $\exp(-\mathbf{A}t) \mathbf{x}(t) - \exp(-\mathbf{A}s) \mathbf{x}(s) = \int_s^t \exp(-\mathbf{A}\tau) \mathbf{b}(\tau) d\tau$ .

Multiplying each side by  $\exp(\mathbf{A}t)$  gives the unique solution

$$\mathbf{x}(t) = \exp[\mathbf{A}(t - s)] \mathbf{x}(s) + \int_s^t \exp[\mathbf{A}(t - \tau)] \mathbf{b}(\tau) d\tau$$

# The Diagonal Case

The **diagonal case** occurs when  $\mathbf{A} = \mathbf{\Lambda} = \mathbf{diag}(\lambda_1, \dots, \lambda_n)$ .

Then the system  $\dot{\mathbf{x}}(t) - \mathbf{Ax}(t) = \mathbf{b}(t)$  of  $n$  coupled equations reduces to the system of  $n$  uncoupled equations

$$\dot{x}_i(t) = a_{ii}x_i(t) + b_i(t) = \lambda_i x_i(t) + b_i(t) \quad (i = 1, \dots, n)$$

one in each variable  $x_i$ , with respective solutions

$$x_i(t) = e^{\lambda_i t} x_i(s) + \int_s^t e^{\lambda_i(t-\tau)} b_i(\tau) d\tau$$

## The Diagonalizable Case

Suppose that  $\mathbf{A}$  has  $n$  distinct eigenvalues — or if not, then  $n$  linearly independent eigenvectors that make up the columns of the matrix  $\mathbf{V}$ .

Then  $\mathbf{A} = \mathbf{V}\mathbf{\Lambda}\mathbf{V}^{-1}$  and  $\mathbf{A}t = \mathbf{V}(\mathbf{\Lambda}t)\mathbf{V}^{-1}$  implying that  $\exp(\mathbf{A}t) = \mathbf{V}\exp(\mathbf{\Lambda}t)\mathbf{V}^{-1}$ .

Hence the solution

$$\mathbf{x}(t) = \exp[\mathbf{A}(t-s)] \mathbf{x}(s) + \int_s^t \exp[\mathbf{A}(t-\tau)] \mathbf{b}(\tau) d\tau$$

simplifies to

$$\mathbf{x}(t) = \mathbf{V} \exp[\mathbf{\Lambda}(t-s)] \mathbf{V}^{-1} \mathbf{x}(s) + \int_s^t \mathbf{V} \exp[\mathbf{\Lambda}(t-\tau)] \mathbf{V}^{-1} \mathbf{b}(\tau) d\tau$$

Of course, the transformation  $\mathbf{y}(t) := \mathbf{V}^{-1} \mathbf{x}(t)$  takes us back to the diagonal case, with  $\dot{\mathbf{y}}(t) := \mathbf{\Lambda} \mathbf{y}(t) + \mathbf{V}^{-1} \mathbf{b}(t)$ .

## A Stability Condition

When  $\mathbf{\Lambda} = \mathbf{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$ ,  
one has  $\exp(\mathbf{\Lambda}) = \mathbf{diag}(e^{\lambda_1}, e^{\lambda_2}, \dots, e^{\lambda_n})$ .

Furthermore  $\exp(\mathbf{\Lambda}t) = \mathbf{diag}(e^{\lambda_1 t}, e^{\lambda_2 t}, \dots, e^{\lambda_n t})$ .

This converges to the zero matrix as  $t \rightarrow \infty$   
if and only if each  $e^{\lambda_i t} \rightarrow 0$ ,  
which is true iff each eigenvalue  $\lambda_i$  has a negative real part.

Similarly, if  $\mathbf{A}$  is diagonalizable, with  $\mathbf{A} = \mathbf{V}\mathbf{\Lambda}\mathbf{V}^{-1}$ ,  
then consider the new variables  $\mathbf{y} = \mathbf{V}^{-1}\mathbf{x}$ .

The differential equation  $\dot{\mathbf{x}}(t) = \mathbf{A}\mathbf{x}(t)$  becomes transformed to

$$\dot{\mathbf{y}}(t) = \mathbf{V}^{-1}\dot{\mathbf{x}}(t) = \mathbf{V}^{-1}\mathbf{A}\mathbf{x}(t) = \mathbf{\Lambda}\mathbf{V}^{-1}\mathbf{x}(t) = \mathbf{\Lambda}\mathbf{y}(t)$$

Because  $\mathbf{V}$  is invertible, one has  $\mathbf{x}(t) \rightarrow \mathbf{0} \iff \mathbf{y}(t) \rightarrow \mathbf{0}$ .

Once again, stability holds  
iff each eigenvalue of the matrix  $\mathbf{A}$  has a negative real part.

This is true even when  $\mathbf{A}$  is not diagonalizable.

## The Schrödinger Equation in $\mathbb{C}^n$

A **wave function** is a mapping  $\mathbb{R} \ni t \mapsto \psi(t) \in \mathbb{C}^n$ .

**Schrödinger's wave equation** is a linear equation that, in a simple case, can be written in the form  $\dot{\psi}(t) = -i \mathbf{H} \psi(t)$  where  $\mathbf{H}$  is a **Hamiltonian** “energy” matrix with complex elements that is self-adjoint.

Because  $\mathbf{H}$  is self-adjoint, it can be diagonalized so that, after a change of variables,

one has  $\dot{\psi}(t) = -i \mathbf{diag}(h_1, \dots, h_n) \psi(t)$

and so  $\dot{\psi}_k(t) = -i h_k \psi_k(t)$  for each  $k \in \mathbb{N}_n$ .

For each possible initial value  $\psi(0) \in \mathbb{C}^n$ , and for each  $k \in \mathbb{N}_n$ , the unique solution is

$$\psi_k(t) = \psi_k(0) e^{-i h_k t} = \psi_k(0) [\cos(-h_k t) + i \sin(-h_k t)]$$

This is a wave or oscillatory solution with frequency  $h_k$ .

Generally, the eigenvalues in the spectrum of  $\mathbf{H}$ , which are all real, are possible frequencies of oscillatory solutions.

# Lecture Outline

## First-Order Differential Equations in One Variable

Introduction

Picard's Method

General First-Order Affine Equation

Constant and Undetermined Coefficients

Stability in the Autonomous Case

## Second-Order Differential Equations in One Variable

Introduction

The Inhomogeneous Equation

The Method of Undetermined Coefficients

Stability

## First-Order Multivariable Differential Equations

Introduction

Prominent Examples and Stability Conditions

**Autonomous Nonlinear Equations in Many Variables**

Existence and Uniqueness Theorem

# Autonomous First-Order Equations

Let  $\mathbf{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a general function that may be non-linear.

Consider the autonomous differential equation  $\dot{\mathbf{x}} = \mathbf{F}(\mathbf{x})$ .

A **solution** satisfying the initial condition  $\mathbf{x}(s) = \bar{\mathbf{x}}$  is a differentiable function  $[s, t) \ni t \mapsto \mathbf{x}(t)$  that satisfies  $\dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{x}(t))$  for almost all  $t \geq s$ .

Equivalently, for almost all  $t \geq s$ , one must have

$$\mathbf{x}(t) = \mathbf{x}(s) + \int_s^t \mathbf{F}(\mathbf{x}(\tau)) d\tau$$

# Stationary States and Rest Points

A **stationary state** is a point  $\mathbf{x}^* \in \mathbb{R}^n$  with the property that if  $\mathbf{x}(s) = \mathbf{x}^*$  at any time  $s$ , then  $\mathbf{x}(t) = \mathbf{x}^*$  at all times  $t \geq s$ .

A **rest point** is a state  $\bar{\mathbf{x}} \in \mathbb{R}^n$  with the property that  $\mathbf{F}(\bar{\mathbf{x}}) = \mathbf{0}$ .

## Theorem

*Any rest point is a stationary state, and conversely.*

## Proof.

If  $\mathbf{F}(\bar{\mathbf{x}}) = \mathbf{0}$ , then the solution of  $\mathbf{x}(t) = \mathbf{x}(s) + \int_s^t \mathbf{F}(\mathbf{x}(\tau)) d\tau$  with  $\mathbf{x}(s) = \bar{\mathbf{x}}$  satisfies  $\mathbf{x}(t) = \mathbf{x}(s) = \bar{\mathbf{x}}$  for all  $t \geq s$ .

Conversely, if that solution satisfies  $\mathbf{x}(t) = \mathbf{x}(s) = \mathbf{x}^*$  for all  $t \geq s$ , then  $\dot{\mathbf{x}}(t) = \mathbf{F}(\mathbf{x}(t)) = \mathbf{F}(\mathbf{x}^*) = \mathbf{0}$  for all  $t \geq s$ . □

# Local Stability of a Stationary State

Let  $\mathbf{F}'(\mathbf{x})$  denote the  $n \times n$  **Jacobian matrix** whose elements are the partial derivatives  $\mathbf{F}'_{ij}(\mathbf{x}) = \frac{\partial}{\partial x_j} F_i(\mathbf{x})$  of the different components  $(F_i(\mathbf{x}))_{i=1}^n$ .

Any particular steady state  $\mathbf{x}^*$  is locally asymptotically stable if and only if all the eigenvalues of  $\mathbf{F}'(\mathbf{x}^*)$  have negative real parts.

## A System with Two Variables

Consider the **coupled pair**  $\dot{x} = f(x, y)$ ,  $\dot{y} = g(x, y)$  of differential equations.

Let  $(a, b)$  be any stationary point satisfying both  $f(a, b) = 0$  and  $g(a, b) = 0$ .

The Jacobian matrix at the stationary point takes the form

$$\mathbf{J}(a, b) = \begin{pmatrix} \frac{\partial f}{\partial x}(a, b) & \frac{\partial f}{\partial y}(a, b) \\ \frac{\partial g}{\partial x}(a, b) & \frac{\partial g}{\partial y}(a, b) \end{pmatrix}$$

## Local Saddle Point with Two Variables

The product of the two eigenvalues  $\lambda_1, \lambda_2$  of  $\mathbf{J}(a, b)$  equals its determinant  $\det \mathbf{J}(a, b)$ .

The two eigenvalues are real and have opposite signs if and only if  $\det \mathbf{J}(a, b) < 0$ .

This is a sufficient condition for the steady state to be unstable.

But if  $(x(0) - a, y(0) - b)^\top$  is an eigenvector corresponding to the negative eigenvalue, then in the case when the equations are linear and so  $\mathbf{J}$  is constant, the solution will converge to the steady state.

This is **saddle point** stability.

# The Lotka–Volterra Predator–Prey Model

Foxes are predators; their prey includes rabbits.

Let  $x$  denote the expected population of rabbits, and  $y$  denote expected population of foxes.

Assume these populations are linked by the **coupled** differential equations

$$\begin{aligned}\dot{x} &= x(k - ay) \\ \dot{y} &= y(bx - h)\end{aligned}$$

where  $a, b, h, k$  are all positive parameters.

Thus:

1. the rabbit population growth rate  $\frac{d}{dt} \ln x = \dot{x}/x$  is a decreasing affine function of the fox population;
2. whereas the fox population growth rate  $\frac{d}{dt} \ln y = \dot{y}/y$  is an increasing affine function of the rabbit population.

## Lotka–Volterra: Phase Plane Analysis

Given the system  $\dot{x} = x(k - ay)$  and  $\dot{y} = y(bx - h)$ , the two **nullclines** where  $\dot{x} = 0$  and  $\dot{y} = 0$  are given by  $y = k/a$  and  $x = h/b$  respectively.

So the steady state is at  $(x, y) = (h/b, k/a)$ .

The Jacobian matrix  $\mathbf{J}(x, y) = \begin{pmatrix} \frac{\partial \dot{x}}{\partial x} & \frac{\partial \dot{x}}{\partial y} \\ \frac{\partial \dot{y}}{\partial x} & \frac{\partial \dot{y}}{\partial y} \end{pmatrix}$

satisfies  $\mathbf{J}(x, y) = \begin{pmatrix} k - ay & -ax \\ yb & bx - h \end{pmatrix}$ .

It reduces to  $\begin{pmatrix} 0 & -ah/b \\ bk/a & 0 \end{pmatrix}$  at the steady state  $(h/b, k/a)$ .

The characteristic equation is  $\det \begin{pmatrix} -\lambda & -ah/b \\ bk/a & -\lambda \end{pmatrix} = \lambda^2 + hk = 0$ ,

whose roots are the complex numbers  $\lambda = \pm i\sqrt{hk}$ .

As the following diagram suggests, there can be **limit cycles** with  $x(t) = \xi \cos \sqrt{hkt}$  and  $y(t) = \eta \sin \sqrt{hkt}$ .



## Saddle Point Example

Consider a macro model where: (i)  $K$  denotes capital;  
(ii)  $Y$  denotes output; and (iii)  $C$  denotes consumption.

Suppose that net investment  $\dot{K} = Y - C$ , that  $Y = aK - bK^2$ ,  
and  $\dot{C} = w(a - 2bk)C$ , where  $a, b, k, w$  are positive constants.

This gives the coupled system with

$$\dot{K} = aK - bK^2 - C \text{ and } \dot{C} = w(a - 2bK)C$$

The two nullclines are  $C = aK - bK^2$  and  $K = a/2b$ .


These intersect at the stationary point  $(K^*, C^*) = (a/2b, a^2/4b)$ .

The Jacobian matrix is


$$\mathbf{J}(K, C) = \begin{pmatrix} \frac{\partial \dot{K}}{\partial K} & \frac{\partial \dot{K}}{\partial C} \\ \frac{\partial \dot{C}}{\partial K} & \frac{\partial \dot{C}}{\partial C} \end{pmatrix} = \begin{pmatrix} a - 2bK & -1 \\ -2wbC & w(a - 2bK) \end{pmatrix}$$

This reduces to  $\begin{pmatrix} 0 & -1 \\ -\frac{1}{2}a^2w & 0 \end{pmatrix}$  at the steady state.

## Phase Diagram I



## Phase Diagram II



## Stability Analysis

The Jacobian matrix at the steady state is  $\begin{pmatrix} 0 & -1 \\ -\frac{1}{2}a^2w & 0 \end{pmatrix}$ .

This matrix has trace 0 and negative determinant  $-\frac{1}{2}a^2w$ .

So the two eigenvalues have sum 0 and product  $-\frac{1}{2}a^2w$ .

It follows that the eigenvalues are  $\pm\lambda$  where  $\lambda^2 = \frac{1}{2}a^2w$  and so  $\lambda = a\sqrt{w/2}$ .

The general solution near the steady state takes the form

$$\begin{pmatrix} K - K^* \\ C - C^* \end{pmatrix} = \begin{pmatrix} A_1 \\ A_2 \end{pmatrix} e^{\lambda t} + \begin{pmatrix} B_1 \\ B_2 \end{pmatrix} e^{-\lambda t}$$

for arbitrary constant vectors  $(A_1, A_2)^\top$  and  $(B_1, B_2)^\top$ .

This converges to the steady state at  $(K^*, C^*) = (a/2b, a^2/4b)$  if and only if  $A_1 = A_2 = 0$ .

It follows that the steady state is a saddle point.

# Lecture Outline

## First-Order Differential Equations in One Variable

Introduction

Picard's Method

General First-Order Affine Equation

Constant and Undetermined Coefficients

Stability in the Autonomous Case

## Second-Order Differential Equations in One Variable

Introduction

The Inhomogeneous Equation

The Method of Undetermined Coefficients

Stability

## First-Order Multivariable Differential Equations

Introduction

Prominent Examples and Stability Conditions

Autonomous Nonlinear Equations in Many Variables

**Existence and Uniqueness Theorem**

# Existence and Uniqueness Theorem, I

**Note:** In the following,  
we use ordinary Roman rather than bold letters  
for vectors in the finite-dimensional space  $\mathbb{R}^d$ .

Extract from pp. 355–356 in ch. 6 of David Applebaum (2009)  
*Lévy Processes and Stochastic Calculus, 2nd edn.* (Cambridge)

Let  $b : \mathbb{R}^d \rightarrow \mathbb{R}^d$ , so that  $b = (b^1, \dots, b^d)$   
where  $b^i : \mathbb{R}^d \rightarrow \mathbb{R}$  for  $1 \leq i \leq d$ .

We study the **initial value problem** posed  
by the vector-valued differential equation  $\frac{d}{dt}c(t) = b(c(t))$   
with fixed initial condition  $c(0) = c_0 \in \mathbb{R}^d$ ,  
whose solution, if it exists, is a curve  $(c(t), t \in \mathbb{R})$  in  $\mathbb{R}^d$ .

## Existence and Uniqueness Theorem, II

We say that  $b$  is (globally) **Lipschitz** if there exists  $K > 0$  such that, for all  $x, y \in \mathbb{R}^d$ ,  $\|b(x) - b(y)\| \leq K\|x - y\|$ .

**Exercise 6.1.1** Show that, if  $b$  is differentiable with bounded partial derivatives, then it is Lipschitz.

**Exercise 6.1.2** Deduce that, if  $b$  is Lipschitz, then it satisfies a linear growth condition  $\|b(x)\| \leq L(1 + \|x\|)$  for all  $x \in \mathbb{R}^d$ , where  $L = \max\{K, \|b(0)\|\}$ .

**Theorem 6.1.3** If  $b : \mathbb{R}^d \rightarrow \mathbb{R}^d$  is (globally) Lipschitz, then there exists a unique solution  $c : \mathbb{R} \rightarrow \mathbb{R}^d$  of the initial value problem  $\frac{d}{dt}c(t) = b(c(t))$  with fixed initial condition  $c(0) = c_0 \in \mathbb{R}^d$ .

The proof offered by Applebaum does not use a contraction mapping theorem.

Rather, it bounds possible solutions within error bands that are exponential functions that converge to zero.