

# EC9A0: Pre-sessional Advanced Mathematics Course

## Real Analysis

Pablo F. Beker  
Department of Economics  
University of Warwick

September 2024

# Slides Outline

- 1 Sets
  - Definition and Operations
  - Binary Relation
  - Correspondences and Functions
  - The Real Numbers
  - The Euclidean Space
  - Fields
  - Vector Spaces
  - Metric Spaces
  - Normed Vector Spaces
- 2 Sequences
  - Sequences in  $\mathbb{R}^K$
  - Limits of Sequences
- 3 Topology of  $\mathbb{R}^K$ 
  - Open Sets
  - Closed Sets
  - Compact Sets
  - Limits of Functions in  $\mathbb{R}$
  - Continuity of Functions
  - Differentiability
  - Mean Value Theorem and Taylor's Theorem

# Sets: Definition and Operations

- A set is a collection of (finitely or infinitely many) objects.
  - For any set  $A$ , we use the notation  $x \in A$  to indicate that “ $x$  is an element of  $A$ ” (“or belongs to  $A$ ” or “is a member of  $A$ ”).
  - The *empty set*,  $\emptyset$ , is the only set with no elements at all.
  - $\mathbb{N} := \{1, 2, \dots\}$  denotes the (countably infinite) set of *natural numbers*
  - $\mathbb{R}$  denotes the (uncountable) set of *real numbers*.
- Two sets  $A$  and  $B$  are equal ( $A = B$ ) if they have the same elements.
- If every member of  $A$  is also a member of  $B$ , we say that  $A$  is a **subset** of  $B$  and write  $A \subseteq B$ .
  - $A = B$  if and only if  $A \subseteq B$  and  $B \subseteq A$ .
  - If  $A \subseteq B$  but  $A \neq B$ , then  $A$  is said to be a proper subset of  $B$ , denoted  $A \subset B$ .
  - The set of all subsets of  $A$  is called **the power set** of  $A$  and denoted  $2^A$ .
- Given any sets  $A$  and  $B$ , **their union** is  $A \cup B \equiv \{x : x \in A \text{ or } x \in B\}$ .
- Given any sets  $A$  and  $B$ , **their intersection** is  $A \cap B \equiv \{x : x \in A \ \& \ x \in B\}$ .
- Given any sets  $A$  and  $B$ , **the cartesian product**  $A \times B$  is the set  $\{(a_1, b_1), (a_2, b_2), \dots\}$  where  $a_i \in A$  and  $b_i \in B$  for all  $i$ .

# Binary Relation

## Definition

Let  $X$  and  $Y$  be two nonempty subsets. A subset  $R$  of  $X \times Y$  is called a **binary relation** from  $X$  to  $Y$ . If  $(x, y) \in R$ , then we write  $x R y$

# Correspondences and Functions

## Definition

A *correspondence*  $f$  from a set  $X \neq \emptyset$  into a set  $Y \neq \emptyset$ , denoted  $f : X \rightarrow Y$ , is a relation  $f \subseteq X \times 2^Y$

- 1 for every  $x \in X$ , there exists a  $Y' \subseteq Y$  such that  $x f Y'$ ,
- 2 for every  $Y', Z' \subseteq Y$  with  $x f Y'$  and  $x f Z'$ , we have  $Y' = Z'$ .

(a rule that assigns to each  $x \in X$  a unique set  $f(x) \subseteq Y$ ).

## Definition

A *function*  $f$  from a set  $X \neq \emptyset$  into a set  $Y \neq \emptyset$ , denoted  $f : X \rightarrow Y$ , is a relation  $f \subseteq X \times Y$  such that

- 1 for every  $x \in X$ , there exists a  $y \in Y$  such that  $x f y$ ,
- 2 for every  $y, z \in Y$  with  $x f y$  and  $x f z$ , we have  $y = z$ .

(a rule that assigns to each  $x \in X$  a unique  $f(x) \in Y$ )

# Functions

Given function  $f : X \mapsto Y$ ,

- $X$  is said to be the *domain*  $Y$  its *target set* or *co-domain*.
- If  $f : X \rightarrow Y$  and  $A \subseteq X$ , the *image of  $A$  under  $f$* , denoted by  $f[A]$ , is the set

$$f[A] = \{y \in Y \mid \exists x \in A : f(x) = y\}.$$

- The image  $f[X]$  of the whole domain is called the *range* of  $f$ .
- If  $f : X \rightarrow Y$ , and  $B \subset Y$ , the *inverse image of  $B$  under  $f$* , denoted  $f^{-1}[B]$ , is the set

$$f^{-1}[B] = \{x \in X \mid f(x) \in B\} \quad (\text{or } f^{-1}[B] = \{x \in X \mid f(x) \in B\}).$$

# Properties of Functions

## Definition

Function  $f : X \rightarrow Y$  is said to be:

- *Onto*, or *surjective*, if  $f[X] = Y$ ;
- *One-to-one*, or *injective*, if  $f(x_1) = f(x_2) \Rightarrow x_1 = x_2$ ;
- *Bijjective*, if it is both onto and one-to-one.

## Examples

- $f : \mathbb{R}^2 \mapsto \mathbb{R}$  defined by  $f(x_1, x_2) = x_1^2 + x_2^2$  is neither one-to-one nor onto.
- $f : \mathbb{R} \setminus \{0\} \mapsto \mathbb{R}$  defined by  $f(x) = \frac{1}{x}$  is one-to-one but not onto.
- $f : \mathbb{R}^2 \mapsto \mathbb{R}$  defined by  $f(x_1, x_2) = x_1 + x_2$  is onto but not one-to-one.
- $f : \mathbb{R} \mapsto \mathbb{R}$  defined by  $f(x) = x$  is one-to-one and onto.

# Inverse Function

## Definition

If  $f : X \rightarrow Y$  is a one-to-one function, the *inverse function*  $f^{-1} : f[X] \rightarrow X$  is implicitly defined by  $f^{-1}(y) = f^{-1}[\{y\}]$ .

## Theorem

The function  $f : X \rightarrow Y$  is onto iff  $f^{-1}[B] \neq \emptyset$  for all non-empty  $B \subseteq Y$ .

**Proof:** ( $\Rightarrow$ ) Suppose  $f : X \mapsto Y$  is onto.

- ① Let  $B \subseteq Y$ . We need to show  $f^{-1}[B] \equiv \{x \in X \mid f(x) \in B\} \neq \emptyset$ .
- ② Let  $\tilde{y} \in B$ . Since  $f$  is onto,  $\{y \in Y \mid f(x) = y \text{ for some } x \in X\} = Y$ .
- ③ Then, there exists  $x \in X$  such that  $f(x) = \tilde{y}$ . Thus,  $f^{-1}[B] \neq \emptyset$ .

( $\Leftarrow$ ) Suppose  $f^{-1}[B] \neq \emptyset$  for all non-empty  $B \subseteq Y$ .

- ① We need to show that  $f[X] \equiv \{y \in Y \mid f(x) = y \text{ for some } x \in X\} = Y$ .
- ② Since  $f[X] \subseteq Y$ , it suffices to show that  $Y \subseteq f[X]$ . Let  $\tilde{y} \in Y$ .
- ③ By hypothesis,  $f^{-1}(\{\tilde{y}\}) \neq \emptyset$ . Hence, there is  $x \in X$  such that  $f(x) = \tilde{y}$ .
- ④ Then,  $\tilde{y} \in \{y \in Y \mid f(x) = y \text{ for some } x \in X\} \equiv f[X]$ . Thus,  $Y \subseteq f[X]$  ■.

# The Real Numbers: Infimum and Supremum

## Definition

Fix a set  $Y \subseteq \mathbb{R}$ . A number  $\alpha \in \mathbb{R}$  is *an upper bound of  $Y$*  if  $y \leq \alpha$  for all  $y \in Y$ , and is *a lower bound of  $Y$*  if the opposite inequality holds.

## Definition

$\alpha \in \mathbb{R}$  is *the least upper bound of  $Y$* , denoted  $\alpha = \sup Y$ , if:

- 1  $\alpha$  is an upper bound of  $Y$ ; and
- 2  $\gamma \geq \alpha$  for any other upper bound  $\gamma$  of  $Y$ .

## Definition

$\beta \in \mathbb{R}$  is *the greatest lower bound of  $Y$* , denoted  $\beta = \inf Y$ , if:

- 1  $\beta$  is a lower bound of  $Y$ ; and
- 2 if  $\gamma$  is a lower bound of  $Y$ , then  $\gamma \leq \beta$ .

**THE COMPLETENESS AXIOM:** Every nonempty subset  $S$  of  $\mathbb{R}$  that is bounded from above has a supremum in  $\mathbb{R}$ .

# The Euclidean Space

- For any  $K \in \mathbb{N}$ , the  $K$ -dimensional real (Euclidean) space is the  $K$ -fold Cartesian product of  $\mathbb{R}$ , denoted by  $\mathbb{R}^K$ .
  - $x \in \mathbb{R}^K \implies x = (x_1 \ x_2 \ \dots \ x_K)$ .
- The *origin* of  $\mathbb{R}^K$  is the vector zero given by  $(0, 0, \dots, 0)$ .
- Given any pair  $x, y \in \mathbb{R}^K$  where  $K \geq 2$ ,
  - 1  $x \gg y$  if  $x_i > y_i$  for all  $i = 1, \dots, K$ ,
  - 2  $x > y$  if  $x \neq y$  and  $x_i \geq y_i$  for all  $i = 1, \dots, K$ ,
  - 3  $x \geq y$  if  $x_i \geq y_i$  for all  $i = 1, \dots, K$ .
- The *non-negative orthant* of  $\mathbb{R}^K$  is  $\mathbb{R}_+^K := \{x \in \mathbb{R}^K \mid x \geq 0\}$ ;
- The *positive orthant* of  $\mathbb{R}^K$  is  $\mathbb{R}_{++}^K := \{x \in \mathbb{R}^K \mid x \gg 0\}$ ;
- No special notation for the set  $\mathbb{R}_+^K \setminus \{0\} = \{x \in \mathbb{R}^K \mid x > 0\}$ ;
- Define vector addition by  $x + y = (x_1 + y_1 \ x_2 + y_2 \ \dots \ x_K + y_K)$ ;
- Define scalar multiplication by  $\alpha x = (\alpha x_1 \ \alpha x_2 \ \dots \ \alpha x_K)$ .

# Fields

## Definition

A set  $\mathbb{F}$  is said to be a **field** if there are two binary operations  $(x, y) \mapsto x \oplus y$  from  $\mathbb{F} \times \mathbb{F}$  to  $\mathbb{F}$  and  $(x, y) \mapsto x \otimes y$  from  $\mathbb{F} \times \mathbb{F}$  to  $\mathbb{F}$  called addition and multiplication, respectively, such that for all  $x, y, z \in \mathbb{F}$ :

- 1  $x \oplus y = y \oplus x$  (addition commutes);
- 2  $(x \oplus y) \oplus z = x \oplus (y \oplus z)$  (addition is associative);
- 3 There exists an element  $0 \in \mathbb{F}$ , such that  $x \oplus 0 = x$  (additive identity);
- 4 For each  $x \in \mathbb{F}$ , there is a unique element in  $\mathbb{F}$ , denoted  $-x$ , such that  $x \oplus (-x) = 0$  (negative);
- 5  $x \otimes y = y \otimes x$  (multiplication is commutative);
- 6  $(x \otimes y) \otimes z = x \otimes (y \otimes z)$  (multiplication is associative);
- 7 There is an element  $1 \in \mathbb{F}$  s.t.  $1 \neq 0$  and  $1 \otimes x = x$ ; (multiplicative identity)
- 8 If  $x \in \mathbb{F}$  and  $x \neq 0$ , there is an element  $\frac{1}{x} \in \mathbb{F}$  such that  $x \otimes (\frac{1}{x}) = 1$
- 9  $x \otimes (y \oplus z) = x \otimes y \oplus x \otimes z$  (distributive law);

# Vector Spaces

## Definition

A set  $L$  is said to be a **vector (or linear) space over the scalar field  $\mathbb{F}$**  if there are two binary operations  $(x, y) \mapsto x \oplus y$  from  $L \times L$  to  $L$  and  $(\lambda, x) \mapsto \lambda \otimes x$  from  $\mathbb{F} \times L$  to  $L$  called addition and scalar multiplication, respectively, and a unique *null vector*  $\theta \in L$ , such that for all  $x, y, z \in L$  and  $\lambda, \mu \in \mathbb{F}$ :

- 1  $x \oplus y = y \oplus x$  (addition commutes);
- 2  $(x \oplus y) \oplus z = x \oplus (y \oplus z)$  (addition is associative);
- 3  $x \oplus \theta = x$  (additive identity);
- 4 for each  $x \in L$ , there is a unique *inverse*  $-x$  such that  $x \oplus (-x) = \theta$ ;
- 5  $\lambda \otimes (\mu \otimes x) = (\lambda \cdot \mu) \otimes x$  (scalar mult. is associative);
- 6  $1 \otimes x = x$  (multiplicative identity);
- 7  $0 \otimes x = \theta$ ;
- 8  $(\lambda + \mu) \otimes x = \lambda \otimes x \oplus \mu \otimes x$  (first distributive law);
- 9  $\lambda \otimes (x \oplus y) = \lambda \otimes x \oplus \lambda \otimes y$  (second distributive law).

# Vector Spaces: Examples

## Examples

- $\mathbb{R}^K$  is a vector space over the field  $\mathbb{R}$ .
- The set  $\mathbb{R}^\infty$  consisting of all infinite sequences  $\{x_0, x_1, x_2, \dots\}$  is a vector space.
- The unit circle in  $\mathbb{R}^2$  is not a vector space over the field  $\mathbb{R}$ .
- The set of all nonnegative functions on  $[a, b]$  is not a vector space over the field  $\mathbb{R}$ .
- The set  $\mathbb{R}$  with  $x \oplus y \equiv x + y + 7$  and  $r \otimes x \equiv rx + 7(r - 1)$ , is a vector space over the field  $\mathbb{R}$ .

# Metric Spaces: Distance Function

## Definition

Given any set  $X$ , the function  $d : X \times X \rightarrow \mathbb{R}$  is a *metric or distance function* on  $X$  if the following properties hold:

- *Positivity*:  $d(x, y) \geq 0$  for all  $x, y \in X$ , with  $d(x, y) = 0$  iff  $x = y$ .
- *Symmetry*:  $d(x, y) = d(y, x)$ .
- *Triangle Inequality*:  $d(x, z) \leq d(x, y) + d(y, z), \forall x, y, z \in X$ .

## Example

Euclidean distance:  $d(x, y) = (\sum_{i \in K} (x_i - y_i)^2)^{1/2}$ .

## Example

Let  $p \in \mathbb{R}_+$  and  $d_p : \mathbb{R}^K \times \mathbb{R}^K \rightarrow \mathbb{R}$  by  $d_p(x, y) = (\sum_{i \in K} |x_i - y_i|^p)^{\frac{1}{p}}$ .

- $d_p$  is a distance iff  $p \geq 1$ .

# Metric Spaces: Definition

## Definition

A **metric space** is  $(X, d)$  where  $X$  is a set and  $d : X \times X \rightarrow \mathbb{R}$  is a metric.

## Examples

- 1 the set of integers with  $d(x, y) = |x - y|$ .
- 2 the set of integers with
$$d(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y. \end{cases}$$
- 3  $\mathbb{R}$  with  $d(x, y) = f(|x - y|)$ , where  $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  is strictly increasing, and strictly concave, with  $f(0) = 0$ .

## Definition

A neighborhood with radius  $\epsilon$  around  $x \in X$  is the set  $B_\epsilon(x) \equiv \{y \in X : d(x, y) \leq \epsilon\}$

# Normed Vector Spaces: Norms

## Definition

Given any vector space  $X$ , a **norm on  $X$**  is a function  $\|\cdot\| : X \mapsto \mathbb{R}$  such that for all  $x, y \in X$  and  $\alpha \in \mathbb{R}$ :

- 1  $\|x\| \geq 0$ , with equality if and only if  $x = \theta$ ;
  - 2  $\|\alpha x\| = |\alpha| \|x\|$ ; and
  - 3  $\|x + y\| \leq \|x\| + \|y\|$  (the triangle inequality)
- In order to measure how far from 0 an element  $x$  of  $\mathbb{R}^K$  is, we use the *Euclidean norm* which is defined as

$$\|x\| = \left( \sum_{k=1}^K x_k^2 \right)^{1/2} .$$

# Normed Vector Spaces: Definition

## Definition

A **normed vector space** is a pair  $(X, \|\cdot\|)$  where  $X$  is a vector space and  $\|\cdot\| : X \mapsto \mathbb{R}$  is a norm.

- It is standard to view any normed vector space  $(X, \|\cdot\|)$  as a metric space where the metric  $d(x, y) = \|x - y\|$  for all  $x, y \in X$ .

## Examples

- $X = \mathbb{R}^K$ , with  $\|x\| = \left[ \sum_{k=1}^K x_k^2 \right]^{\frac{1}{2}}$  (Euclidean Space)
- $X = \mathbb{R}^K$ , with  $\|x\| = \max_j |x_j|$ .
- $X = \mathbb{R}^K$ , with  $\|x\| = \sum_{k=1}^K |x_k|$ .
- $X$  is the set of all bounded infinite sequences  $\{x_k\}_{k=1}^{\infty}$  with  $\|x\| = \sup_k |x_k|$ . (This space is called  $l_{\infty}$ )

# Sequences in $\mathbb{R}^K$

## Definition

A **sequence** in  $\mathbb{R}^K$  is a function  $f : \mathbb{N} \rightarrow \mathbb{R}^K$ .

- $(a_1, a_2, \dots)$  or  $(a_n)_{n=1}^{\infty}$ , where  $a_n = f(n)$ , for  $n \in \mathbb{N}$ .
- $(a_n)_{n=1}^{\infty}$  is
  - *nondecreasing (increasing)* if  $a_{n+1} \geq (>)a_n$  for all  $n \in \mathbb{N}$ ;
  - *nonincreasing (decreasing)* if  $a_{n+1} \leq (<)a_n$  for all  $n \in \mathbb{N}$ ;
  - *bounded above* if there exists  $\bar{a} \in \mathbb{R}^K$  such that  $a_n \leq \bar{a}$  for all  $n$ ;
  - *bounded below* if there exists  $\underline{a} \in \mathbb{R}^K$  such that  $a_n \geq \underline{a}$  for all  $n$ ;
  - *bounded* if it is bounded both above and below.

## Definition

Given a sequence  $(a_n)_{n=1}^{\infty}$ , a sequence  $(b_m)_{m=1}^{\infty}$  is a **subsequence** of  $(a_n)_{n=1}^{\infty}$  if there exists an increasing sequence  $(n_m)_{m=1}^{\infty}$  such that  $n_m \in \mathbb{N}$  and  $b_m = a_{n_m}$  for all  $m \in \mathbb{N}$ .

## Example

$(1/\sqrt{2m+5})_{m=1}^{\infty}$  is a subsequence of  $(1/\sqrt{n})_{n=1}^{\infty}$  for  $(n_m)_{m=1}^{\infty} = (2m+5)_{m=1}^{\infty}$ .

# Limits of Sequences

## Definition

A sequence  $(a_n)_{n=1}^{\infty}$  in  $\mathbb{R}^K$  **converges** to  $a \in \mathbb{R}^K$  (written  $a_n \rightarrow a$ ), if for each  $\varepsilon > 0$  there exists some  $N_\varepsilon \in \mathbb{N}$  such that

$$d(a_n, a) < \varepsilon \text{ for all } n \geq N_\varepsilon.$$

## Theorem

*Let  $d$  be the Euclidean distance. Then,  $(a_n)_{n=1}^{\infty}$  in  $\mathbb{R}^K$  converges to  $a$  if and only if  $(a_{k,n})_{n=1}^{\infty}$  in  $\mathbb{R}$  converges to  $a_k$  for all  $k = 1, \dots, K$ .*

## Theorem

*Sequence  $(a_n)_{n=1}^{\infty}$  in  $\mathbb{R}^K$  converges to  $a \in \mathbb{R}^K$  if and only if every subsequence of  $(a_n)_{n=1}^{\infty}$  converges to  $a$ .*

# Limits of Sequences

## Definition

For a sequence  $(a_n)_{n=1}^{\infty}$  in  $\mathbb{R}$ , we say that  $\lim_{n \rightarrow \infty} a_n = \infty$  if for all  $\Delta > 0$  there exists some  $n^* \in \mathbb{N}$  such that  $a_n > \Delta$  for all  $n \geq n^*$ . We say that  $\lim_{n \rightarrow \infty} a_n = -\infty$  when  $\lim_{n \rightarrow \infty} (-a_n) = \infty$ . We say that a sequence  $(a_n)_{n=1}^{\infty}$  in  $\mathbb{R}$  **diverges to  $\infty$  ( $-\infty$ )** if  $\lim_{n \rightarrow \infty} a_n = \infty$  ( $-\infty$ ).

## Examples

- 1 Does  $((-1)^n)_{n=1}^{\infty}$  converge? Does  $(-1/n)_{n=1}^{\infty}$ ?
- 2 Does the sequence  $(\frac{3n}{\sqrt{n}})_{n=1}^{\infty}$  have a limit? Does it converge?

# Limits of Sequences: Properties I

## Theorem

If  $a_n \rightarrow x$  and  $a_n \rightarrow y$ , then  $x = y$ .

## Theorem

For sequences  $(a_n)_{n=1}^{\infty}$  in  $\mathbb{R}$  such that  $a_n > 0$  for all  $n \in \mathbb{N}$ ,

$$\lim_{n \rightarrow \infty} a_n = \infty \Leftrightarrow \lim_{n \rightarrow \infty} \frac{1}{a_n} = 0.$$

## Limits of Sequences: Properties II

### Theorem (Arithmetic of Limits)

Let  $(a_n)_{n=1}^{\infty}$  and  $(b_n)_{n=1}^{\infty}$  be sequences in  $\mathbb{R}$ . Suppose that  $a, b \in \mathbb{R}$ , we have that  $\lim_{n \rightarrow \infty} a_n = a$  and  $\lim_{n \rightarrow \infty} b_n = b$ . Then,

- 1  $\lim_{n \rightarrow \infty} (a_n + b_n) = a + b$ ;
- 2  $\lim_{n \rightarrow \infty} (\alpha a_n) = \alpha a$ , for all  $\alpha \in \mathbb{R}$ ;
- 3  $\lim_{n \rightarrow \infty} (a_n b_n) = ab$ ;
- 4 if  $b \neq 0$  and  $b_n \neq 0$  for all  $n \in \mathbb{N}$ , then  $\lim_{n \rightarrow \infty} (a_n / b_n) = a / b$ .

### Theorem (Weak Inequalities are Preserved under Sequential Limits)

If  $a_n \leq \alpha$ , for all  $n \in \mathbb{N}$ , and  $\lim_{n \rightarrow \infty} a_n = a$ , then  $a \leq \alpha$ .

- Can we strengthen the last Theorem to strict inequalities?

## Limits of Sequences: Properties III

### Theorem

*Every sequence  $(a_n)_{n=1}^{\infty}$  has a monotone subsequence.*

### Theorem

*If sequence  $(a_n)_{n=1}^{\infty}$  in  $\mathbb{R}$  is convergent, then it is bounded.*

### Theorem

*If a sequence  $(a_n)_{n=1}^{\infty}$  is monotone and bounded, then it is convergent.*

### Theorem (Bolzano-Weierstrass)

*If sequence  $(a_n)_{n=1}^{\infty}$  is bounded, then it has a convergent subsequence.*

# Cauchy Sequences

## Definition

A sequence  $\{a_n\}_{n=1}^{\infty}$  is a **Cauchy sequence** (or satisfies the **Cauchy criterion**) if for each  $\varepsilon > 0$ , there exists  $N_\varepsilon$  such that

$$d(a_n, a_m) < \varepsilon, \text{ for all } n, m \geq N_\varepsilon.$$

## Example

Is the sequence  $(1/\sqrt{n})_{n=1}^{\infty}$  in  $\mathbb{R}$  Cauchy?

## Theorem

- 1 *If a sequence is convergent, then it is a Cauchy sequence.*
- 2 *If a sequence is Cauchy, then it is bounded.*

## Definition

A metric space  $(X, d)$  is **complete** if every Cauchy sequence in  $X$  converges to an element of  $X$ .

**Fact:**  $\mathbb{R}$  with  $d(x, y) = |x - y|$  is a complete metric space.

# Open Sets

## Definition

A set  $X \subset \mathbb{R}^k$  is *open* if for all  $x \in X$ , there is some  $\varepsilon > 0$  for which  $B_\varepsilon(x) \subseteq X$ .

## Theorem

*The empty set, the open intervals in  $\mathbb{R}$  and  $\mathbb{R}^k$  are open.*

## Theorem

*The union of any collection of open sets is an open set. The intersection of any finite collection of open sets is an open set.*

## Exercise

- 1 Do we really need finiteness in the second part of the last Theorem? Consider  $I_n = (-\frac{1}{n}, \frac{1}{n})$  for all  $n \in \mathbb{N}$ . Find the intersection of all those intervals, denoted  $\bigcap_{n=1}^{\infty} I_n$ . Is it an open set?
- 2 Whether or not a set is open depends on the metric space. So changing either the set or the metric can change the openness of a set.  
For example,  $\{1\}$  is open in  $\mathbb{N}$  under the Euclidean metric. However  $\{1\}$  is not open in  $\mathbb{R}$  under the Euclidean metric. But it is open in  $\mathbb{R}$  under the discrete metric.

# Closed Sets

## Definition

A set  $X \subset \mathbb{R}^K$  is *closed* if for every sequence  $(x_n)_{n=1}^{\infty} \in X$  that converges to  $\bar{x}$ , then  $\bar{x} \in X$ .

## Theorem

*The empty set, the closed intervals in  $\mathbb{R}$  and  $\mathbb{R}^K$  are closed.*

## Theorem

*A set  $X$  is closed if and only if  $X^c$  is open.*

## Theorem

*The intersection of any collection of closed sets is closed. The union of any finite collection of closed sets is closed.*

# Compact Sets

## Definition

A set  $X \subseteq \mathbb{R}^K$  is said to be *bounded above* if there exists  $\alpha \in \mathbb{R}^K$  such that  $x \leq \alpha$  for all  $x \in X$ ; it is said to be *bounded below* if for some  $\beta \in \mathbb{R}^K$  one has that  $x \geq \beta$  is true for all  $x \in X$ ; and it is said to be *bounded* if it is bounded above and below.

## Definition

A set  $X \subseteq \mathbb{R}^K$  is said to be *compact* if it is closed and bounded.

## Exercise

Prove the following statement: if  $(x_n)_{n=1}^{\infty}$  is a sequence defined on a compact set  $X$ , then it has a subsequence that converges to a point in  $X$ .

## Theorem

*A set  $X \subset \mathbb{R}^K$  is compact if and only if every sequence in  $X$  has a subsequence that converges to a point in  $X$ .*

# Limit Points

## Definition

Let  $x \in \mathbb{R}^K$  and  $\delta > 0$ . The **open ball of radius  $\delta$  around  $x$** , denoted  $B_\delta(x)$ , is the set

$$B_\delta(x) = \{y \in \mathbb{R} : d(y, x) < \delta\}.$$

## Definition

The **punctured open ball of radius  $\delta$  around  $x$** , denoted  $B'_\delta(x)$ , is the set  $B'_\delta(x) = B_\delta(x) \setminus \{x\}$ .

## Definition

A point  $\bar{x} \in \mathbb{R}^K$  is a **limit point of  $X \subseteq \mathbb{R}^K$**  if for all  $\varepsilon > 0$ ,  $B'_\varepsilon(\bar{x}) \cap X \neq \emptyset$

# Limits of Functions in $\mathbb{R}$

## Definition

Consider  $f : X \rightarrow \mathbb{R}$ , where  $X \subseteq \mathbb{R}^K$ . Suppose that  $\bar{x} \in \mathbb{R}^K$  is a limit point of  $X$  and that  $\bar{y} \in \mathbb{R}$ . We say that  $\lim_{x \rightarrow \bar{x}} f(x) = \bar{y}$  when for all  $\varepsilon > 0$  there exists  $\delta > 0$  such that  $d(f(x), \bar{y}) < \varepsilon$  for all  $x \in B'_\delta(\bar{x}) \cap X$ .

## Definition

Consider  $f : X \rightarrow \mathbb{R}$ , where  $X \subseteq \mathbb{R}^K$ . Suppose that  $\bar{x} \in \mathbb{R}^K$  is a limit point of  $X$ . We say that  $\lim_{x \rightarrow \bar{x}} f(x) = \infty$  when for all  $\Delta > 0$ , there exists  $\delta > 0$  such that  $f(x) \geq \Delta$  for all  $x \in B'_\delta(\bar{x}) \cap X$ . We say that  $\lim_{x \rightarrow \bar{x}} f(x) = -\infty$  when  $\lim_{x \rightarrow \bar{x}} (-f)(x) = \infty$ .

## Limits of Functions: Examples

### Example

Suppose that  $X = \mathbb{R}$  and  $f : X \rightarrow \mathbb{R}$  is defined by

$$f(x) = \begin{cases} 1/x, & \text{if } x \neq 0, \\ 0, & \text{otherwise.} \end{cases}$$

What is  $\lim_{x \rightarrow 5} f(x)$ ? What is  $\lim_{x \rightarrow 0} f(x)$ ?

### Example

Let  $X = \mathbb{R} \setminus \{0\}$  and  $f : X \rightarrow \mathbb{R}$  is defined by

$$f(x) = \begin{cases} 1, & \text{if } x > 0, \\ -1, & \text{otherwise.} \end{cases}$$

In this case, we claim that  $\lim_{x \rightarrow 0} f(x)$  does not exist.

# Limits of Functions and Sequences

## Theorem

Consider a function  $f : X \rightarrow \mathbb{R}$ , where  $X \subseteq \mathbb{R}^K$ . Suppose that  $\bar{x} \in \mathbb{R}^K$  is a limit point of  $X$  and that  $\bar{y} \in \mathbb{R}$ . Then,  $\lim_{x \rightarrow \bar{x}} f(x) = \bar{y}$  if and only if for every  $(x_n)_{n=1}^{\infty} \in X \setminus \{\bar{x}\}$  that converges to  $\bar{x}$ ,  $\lim_{n \rightarrow \infty} f(x_n) = \bar{y}$ .

**Proof:** ( $\Rightarrow$ ) Suppose  $\lim_{x \rightarrow \bar{x}} f(x) = \bar{y}$  and let  $(x_n)_{n=1}^{\infty}$  in  $X \setminus \{\bar{x}\}$  conv. to  $\bar{x}$ .

- ① Let  $\varepsilon > 0$ .  $\exists \delta > 0$  such that  $|f(x) - \bar{y}| < \varepsilon$  for all  $x \in B'_\delta(\bar{x}) \cap X$ .
- ② Since  $\lim_{n \rightarrow \infty} x_n = \bar{x}$ , there is  $N$  such that  $x_n \in B_\delta(\bar{x})$  for all  $n \geq N$ .
- ③ Since each  $x_n \in X \setminus \{\bar{x}\}$ , we have that  $x_n \in B'_\delta(\bar{x}) \cap X$ .
- ④ Therefore,  $|f(x_n) - \bar{y}| < \varepsilon$  for all  $n \geq N \Rightarrow \lim_{n \rightarrow \infty} f(x_n) = \bar{y}$ .

( $\Leftarrow$ ) Suppose  $\forall (x_n)_{n=1}^{\infty}$  in  $X \setminus \{\bar{x}\}$  &  $x_n \rightarrow \bar{x}$ ,  $\lim_{n \rightarrow \infty} f(x_n) = \bar{y}$  &  $\lim_{x \rightarrow \bar{x}} f(x) \neq \bar{y}$ .

- ①  $\exists \varepsilon > 0$  such that  $\forall \delta > 0$ ,  $\exists x \in B'_\delta(\bar{x}) \cap X$  &  $|f(x) - \bar{y}| \geq \varepsilon$ .
- ② For each  $n \in \mathbb{N}$  there is  $x_n \in B'_{1/n}(\bar{x}) \cap X$  for which  $|f(x_n) - \bar{y}| \geq \varepsilon$ .
- ③ We can construct a sequence  $(x_n)_{n=1}^{\infty}$  in  $X \setminus \{\bar{x}\}$  and  $\lim_{n \rightarrow \infty} x_n = \bar{x}$ .
- ④ Then,  $\lim_{n \rightarrow \infty} f(x_n) \neq \bar{y}$ , a contradiction ■

# Limits of Functions: Properties I

Define:

- $(f + g) : X \rightarrow \mathbb{R}$  by  $(f + g)(x) = f(x) + g(x)$ .
- $(\alpha f) : X \times \mathbb{R} \rightarrow \mathbb{R}$  by  $(\alpha f)(x) = \alpha f(x)$ .
- $(f \cdot g) : X \rightarrow \mathbb{R}$  by  $(f \cdot g)(x) = f(x)g(x)$
- $(\frac{f}{g}) : X_g^* \rightarrow \mathbb{R}$  by  $(\frac{f}{g})(x) = \frac{f(x)}{g(x)}$ , where  $X_g^* = \{x \in X \mid g(x) \neq 0\}$ .

## Theorem

Let  $f : X \rightarrow \mathbb{R}$  and  $g : X \rightarrow \mathbb{R}$ . Let  $\bar{x}$  be a limit point of  $X$ . Suppose that  $\bar{y}_1, \bar{y}_2 \in \mathbb{R}$  and that  $\lim_{x \rightarrow \bar{x}} f(x) = \bar{y}_1$  and  $\lim_{x \rightarrow \bar{x}} g(x) = \bar{y}_2$ .

- 1  $\lim_{x \rightarrow \bar{x}} (f + g)(x) = \bar{y}_1 + \bar{y}_2$ ;
- 2  $\lim_{x \rightarrow \bar{x}} (\alpha f)(x) = \alpha \bar{y}_1$ , for all  $\alpha \in \mathbb{R}$ ;
- 3  $\lim_{x \rightarrow \bar{x}} (f \cdot g)(x) = \bar{y}_1 \cdot \bar{y}_2$ ;
- 4 if  $\bar{y}_2 \neq 0$ , then  $\lim_{x \rightarrow \bar{x}} (f/g)(x) = \bar{y}_1 / \bar{y}_2$ .

# Limits of Functions: Properties II

## Theorem

Consider  $f : X \rightarrow \mathbb{R}$  and  $\bar{y} \in \mathbb{R}$ , and let  $\bar{x} \in \mathbb{R}^K$  be a limit point of  $X$ . If  $f(x) \leq \gamma$  for all  $x \in X$ , and  $\lim_{x \rightarrow \bar{x}} f(x) = \bar{y}$ , then  $\bar{y} \leq \gamma$ .

## Corollary

Consider  $f : X \rightarrow \mathbb{R}$  and  $g : X \rightarrow \mathbb{R}$ , let  $\bar{y}_1, \bar{y}_2 \in \mathbb{R}$ , and let  $\bar{x} \in \mathbb{R}^K$  be a limit point of  $X$ . If  $f(x) \geq g(x)$ , for all  $x \in X$ ,  $\lim_{x \rightarrow \bar{x}} f(x) = \bar{y}_1$  and  $\lim_{x \rightarrow \bar{x}} g(x) = \bar{y}_2$ , then  $\bar{y}_1 \geq \bar{y}_2$ .

# Continuity of Functions

## Definition

Function  $f : X \rightarrow \mathbb{R}$  is *continuous at  $\bar{x} \in X$*  if for every  $\varepsilon > 0$ , there exists  $\delta > 0$  such that  $|f(x) - f(\bar{x})| < \varepsilon$  for all  $x \in B_\delta(\bar{x}) \cap X$ . It is *continuous* if it is continuous at all  $\bar{x} \in X$ .

## Theorem

*Suppose that  $f : X \rightarrow \mathbb{R}$  and  $g : X \rightarrow \mathbb{R}$  are continuous at  $\bar{x} \in X$ , and let  $\alpha \in \mathbb{R}$ . Then, the functions  $f + g$ ,  $\alpha f$  and  $f \cdot g$  are continuous at  $\bar{x}$ .*

*Moreover, if  $g(\bar{x}) \neq 0$ , then  $\frac{f}{g}$  is continuous at  $\bar{x}$ .*

# Properties of Continuous Functions

## Theorem

*The image of a compact set under a continuous function is compact.*

## Theorem

*Function  $f : \mathbb{R}^k \rightarrow \mathbb{R}$  is continuous if and only if for every open set  $U \subseteq \mathbb{R}$  the set  $f^{-1}[U]$  is open.*

## Theorem (The Intermediate Value Theorem in $\mathbb{R}$ )

*If function  $f : [a, b] \rightarrow \mathbb{R}$  is continuous, then for every number  $\gamma$  between  $f(a)$  and  $f(b)$  there exists an  $x \in [a, b]$  for which  $f(x) = \gamma$ .*

# Left- and Right- Continuity

## Definition

One says that  $\lim_{x \searrow \bar{x}} f(x) = \ell$ , if for every  $\varepsilon > 0$  there is a number  $\delta > 0$  such that  $|f(x) - \ell| < \varepsilon$  whenever  $x \in X \cap B_\delta(\bar{x})$  and  $x > \bar{x}$ . In such case, function  $f$  is said to converge to  $\ell$  as  $x$  tends to  $\bar{x}$  from above.

Similarly,  $\lim_{x \nearrow \bar{x}} f(x) = \ell$ , when for every  $\varepsilon > 0$  there is  $\delta > 0$  such that  $|f(x) - \ell| < \varepsilon$  for all  $x \in X \cap B_\delta(\bar{x})$  satisfying that  $x < \bar{x}$ . In this case,  $f$  is said to converge to  $\ell$  as  $x$  tends to  $\bar{x}$  from below.

## Definition

Function  $f : X \rightarrow \mathbb{R}$  is *right-continuous at  $\bar{x} \in X$* , where  $\bar{x}$  is a limit point of  $X$ , if  $\lim_{x \searrow \bar{x}} f(x) = f(\bar{x})$ . It is *right-continuous* if it is right-continuous at every  $\bar{x} \in X$  that is a limit point of  $X$ . Similarly,  $f : X \rightarrow \mathbb{R}$  is *left-continuous at  $\bar{x}$*  if  $\lim_{x \nearrow \bar{x}} f(x) = f(\bar{x})$ , and one says that  $f$  is *left-continuous* if it is left-continuous at all limit point  $\bar{x} \in X$ .

# Differentiability

## Definition

Let  $f : \mathbb{R} \mapsto \mathbb{R}$  be a function defined in a neighbourhood of  $x_0$ . Then  $f$  is said to be **differentiable** at  $x_0$  with derivative equal to the real number  $f'(x_0)$  if for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that  $|x - x_0| < \delta$  implies

$$\left| \frac{f(x) - f(x_0)}{x - x_0} - f'(x_0) \right| \leq \varepsilon$$

- Since  $x - x_0 \neq 0$ , multiply the inequality above by  $|x - x_0|$  to obtain

$$|f(x) - f(x_0) - f'(x_0)(x - x_0)| \leq \varepsilon |x - x_0|$$

to see that  $|f(x) - f(x_0) - f'(x_0)(x - x_0)|$  goes to zero faster than  $|x - x_0|$ .

# Mean Value Theorem and Taylor's Theorem

## Theorem (Mean Value Theorem)

Let  $f$  be a continuous function on  $[a, b]$  that is differentiable in  $(a, b)$ . Then there exists  $x_0 \in (a, b)$  such that  $f'(x_0) = \frac{f(b) - f(a)}{b - a}$ .

## Theorem (Taylor's Theorem)

Let  $f$  be  $\mathbb{C}^n$  in a neighborhood of  $x_0$ , and let

$$T_n(x_0, x) = f(x_0) + f'(x_0)(x - x_0) + \frac{1}{2}f''(x_0)(x - x_0)^2 + \dots + \frac{1}{n!}f^n(x_0)(x - x_0)^n.$$

Then for any  $\varepsilon > 0$ , there exists  $\delta$  such that  $|x - x_0| \leq \delta$  implies

$$|f(x) - T_n(x_0, x)| \leq \varepsilon |x - x_0|^n.$$

## Theorem (Lagrange Remainder Theorem)

Suppose  $f$  is  $\mathbb{C}^{n+1}$  in a neighborhood of  $x_0$ . Then for every  $x$  in the neighbourhood there exists  $x_1$  between  $x_0$  and  $x$  such that

$$f(x) = T_n(x_0, x) + \frac{1}{(n+1)!}f^{n+1}(x_1)(x - x_0)^{n+1}$$