



WARWICK

ECONOMICS

CRETA

Centre for Research in Economic Theory and its Applications

Discussion Paper Series

**Roberts' Weak Welfarism Theorem: A Minor Correction**

Peter J. Hammond

[\(This paper also appears in  
Warwick Economics Research Papers series No: 1308\)](#)

October 2020

(This paper has been updated &  
accepted for publication in  
[Social Choice and Welfare](#))

No: 66

CRETA

Centre for Research in Economic Theory and its Applications

Department of Economics  
University of Warwick, Coventry,  
CV4 7AL, United Kingdom

[warwick.ac.uk/fac/soc/economics/research/centres/creta/papers](http://warwick.ac.uk/fac/soc/economics/research/centres/creta/papers)

## Roberts' Weak Welfarism Theorem: A Minor Correction

Peter J. Hammond: [p.j.hammond@warwick.ac.uk](mailto:p.j.hammond@warwick.ac.uk)

Dept. of Economics, University of Warwick, Coventry CV4 7AL, UK.

Latest version: 2021 March 20th, typeset from `RobertsThmSCWrev3.tex`

### Abstract:

Roberts' "weak neutrality" or "weak welfarism" theorem concerns Sen social welfare functionals which are defined on an unrestricted domain of utility function profiles and satisfy independence of irrelevant alternatives, the Pareto condition, and a form of weak continuity. Roberts (1980) claimed that the induced welfare ordering on social states has a one-way representation by a continuous, monotonic real-valued welfare function defined on the Euclidean space of interpersonal utility vectors — that is, an increase in this welfare function is sufficient, but may not be necessary, for social strict preference. A counter-example shows that weak continuity is insufficient; a minor strengthening to pairwise continuity is proposed instead and its sufficiency demonstrated.

*JEL Classification:* D71

*Keywords:* social welfare functionals, weak welfarism.

# 1 Introduction: Roberts' Claim

Consider a society with a non-empty finite set  $N$  of individuals  $i$ .<sup>1</sup> Let  $X$  be a domain of at least three social states, and  $\mathcal{R}(X)$  the set of all logically possible (complete and transitive) social weak preference orderings  $R$  on  $X$ . For each  $R \in \mathcal{R}(X)$ , let  $P$  and  $I$  denote the corresponding strict preference and indifference relations.

Let  $\mathbb{R}^N$  denote the Euclidean space that consists of the Cartesian product of  $\#N$  copies of the real line  $\mathbb{R}$ . A *utility function profile*  $\mathbf{u}^N$  is a mapping  $X \ni x \mapsto \mathbf{u}^N(x) \in \mathbb{R}^N$ . Let  $\mathcal{U}^N$  denote the set of all utility function profiles on  $X$ . Following Sen (1970, 1977), a *social welfare functional* (or SWFL)  $f$  on a domain  $\mathcal{D} \subseteq \mathcal{U}^N$  is a mapping  $\mathcal{D} \ni \mathbf{u}^N \mapsto f(\mathbf{u}^N) \in \mathcal{R}(X)$  that determines a (complete and transitive) social preference ordering  $R = f(\mathbf{u}^N)$  for each utility function profile in  $\mathcal{D}$ . With some slight abuse of notation, we let  $R(\mathbf{u}^N)$ ,  $P(\mathbf{u}^N)$  and  $I(\mathbf{u}^N)$  denote respectively the weak preference, strict preference, and indifference relations associated with  $f(\mathbf{u}^N)$ .

Given any non-empty subset  $A \subset X$ , say that:

1. two utility function profiles  $\mathbf{u}^N, \tilde{\mathbf{u}}^N \in \mathcal{U}^N$  are *equal on  $A$*  just in case one has  $\mathbf{u}^N(x) = \tilde{\mathbf{u}}^N(x)$  for all  $x \in A$ ;
2. two social preference orderings  $R$  and  $\tilde{R}$  on  $X$  are *equal on  $A$*  just in case, for all  $y, z \in A$ , one has  $y R z \iff y \tilde{R} z$ .

The main part of this paper considers social welfare functionals  $f$  which satisfy at least the first two of the following three axioms:

**Unrestricted domain (U)** The domain  $\mathcal{D}$  of  $f$  is the whole of  $\mathcal{U}^N$ .

**Independence (I)** Given any non-empty subset  $A \subset X$ , if the two utility function profiles  $\mathbf{u}^N, \tilde{\mathbf{u}}^N \in \mathcal{D}$  are equal on  $A$ , then the two associated social orderings  $f(\mathbf{u}^N), f(\tilde{\mathbf{u}}^N) \in \mathcal{R}(X)$  are also equal on  $A$ .

**Pareto indifference ( $\mathbf{P}^0$ )** In case  $y, z \in X$  and  $\mathbf{u}^N \in \mathcal{D}$  satisfy  $\mathbf{u}^N(y) = \mathbf{u}^N(z)$ , the associated social indifference relation satisfies  $y I(\mathbf{u}^N) z$ .

An important result in social choice theory with interpersonal comparisons is the “strong neutrality” or “welfarism” result due to d’Aspremont and Gevers (1977) and Sen (1977, p. 1553). This states that, when  $f$  satisfies all three conditions (U), (I), and ( $\mathbf{P}^0$ ), then there exists a (complete

---

<sup>1</sup>Most notation and definitions are based on Roberts (1980). Where appropriate, however, bold letters are used to indicate vectors.

and transitive) *social welfare ordering*  $R^*$  on  $\mathbb{R}^N$  with the property that  $y R z \iff \mathbf{u}^N(y) R^* \mathbf{u}^N(z)$ . This plays a prominent role among the results appearing in the surveys by Sen (1984), Blackorby, Donaldson and Weymark (1984), d'Aspremont (1985), Mongin and d'Aspremont (1998), and Bossert and Weymark (2004). Both Sen (1977) and d'Aspremont (1985, p. 34) provide complete proofs.<sup>2</sup>

While the Pareto indifference axiom ( $P^0$ ) is appealing, the impossibility theorem in Arrow (1963) is the most prominent of many results that replace it with the following alternative:

**Pareto (P)** In case the two social states  $y, z \in X$  and the utility function profile  $\mathbf{u}^N \in \mathcal{D}$  satisfy  $\mathbf{u}^N(y) \gg \mathbf{u}^N(z)$ , the associated strict preference relation  $P(\mathbf{u}^N)$  on  $X$  satisfies  $y P(\mathbf{u}^N) z$ .<sup>3</sup>

Specifically, under the assumption that individuals' utility functions are ordinally non-comparable, Arrow's impossibility theorem states that (U), (I) and (P) together imply a dictatorship. To develop a theory general enough to cover this important case, Roberts (1980, p. 427) specifies an additional condition that can be restated as follows:<sup>4</sup>

**Weak continuity (WC)** For all utility function profiles  $\mathbf{u}^N \in \mathcal{D}$  and all vectors  $\boldsymbol{\epsilon} \in \mathbb{R}^N$  with  $\boldsymbol{\epsilon} \gg \mathbf{0}$ , there exists a utility function profile  $\mathbf{u}_\epsilon^N \in \mathcal{D}$  satisfying  $\boldsymbol{\epsilon} \gg \mathbf{u}^N(x) - \mathbf{u}_\epsilon^N(x) \gg \mathbf{0}$  for all social states  $x \in X$  with the property that  $f(\mathbf{u}^N) = f(\mathbf{u}_\epsilon^N)$ .

Then Roberts (1980, p. 428) claims the following:

**Claim 1.** *Suppose that the SWFL  $\mathcal{D} \ni \mathbf{u}^N \mapsto f(\mathbf{u}^N) \in \mathcal{R}(X)$  satisfies (U), (I), (P), and (WC). Then there exists a continuous function  $\mathbb{R}^N \ni \mathbf{w} \mapsto W(\mathbf{w}) \in \mathbb{R}$ , strictly increasing with an increase in all its arguments, with the property that for all utility function profiles  $\mathbf{u}^N \in \mathcal{D}$  and all social states  $y, z \in X$  one has*

$$W(\mathbf{u}^N(y)) > W(\mathbf{u}^N(z)) \implies y P(\mathbf{u}^N) z$$

<sup>2</sup>Unfortunately, d'Aspremont's proof, which is otherwise the more elegant of the two, includes a crucial typographical error. The option  $e$  should be chosen so that  $b \neq e \neq d$ .

<sup>3</sup>Given any pair  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^N$  with  $\mathbf{a} = (a_i)_{i \in N}$  and  $\mathbf{b} = (b_i)_{i \in N}$ , we use the following notation for vector orderings: (i)  $\mathbf{a} \geq \mathbf{b}$  in case  $a_i \geq b_i$  for all  $i \in N$ ; (ii)  $\mathbf{a} \gg \mathbf{b}$  in case  $a_i > b_i$  for all  $i \in N$ ; (iii)  $\mathbf{a} > \mathbf{b}$  in case  $\mathbf{a} \geq \mathbf{b}$  but  $\mathbf{a} \neq \mathbf{b}$ . We also let  $\mathbb{R}_{++}^N$  denote the set  $\{\mathbf{a} \in \mathbb{R}^N \mid \mathbf{a} \gg \mathbf{0}\}$  of *strictly positive* vectors in  $\mathbb{R}^N$ .

<sup>4</sup>Roberts' original condition has been restated in a way that seems more appropriate if there is a restricted domain  $\mathcal{D} \subset \mathcal{U}^N$  of admissible utility profiles.

This claim has come to be known as Roberts’ “weak neutrality” or “weak welfarism” theorem.<sup>5</sup> In many of the surveys mentioned above, it was cited as an alternative to the strong neutrality result of d’Aspremont and Gevers (1977) and Sen (1977, p. 1553). The unpublished results by Le Breton (1987) and by Bordes and Le Breton (1987) investigating Roberts’ theorem for restricted economic domains have since been amalgamated with related results that appear in Bordes, Hammond and Le Breton (2005).

Condition (WC), however, is too weak for the claim to hold. To show this, Section 2 provides a counter example which even satisfies the following familiar condition:

**Strict Pareto (P\*)** In case the two social states  $y, z \in X$  and the utility function profile  $\mathbf{u}^N \in \mathcal{D}$  satisfy  $\mathbf{u}^N(y) \geq \mathbf{u}^N(z)$ , the associated social preferences satisfy  $y R(\mathbf{u}^N) z$ , with  $y P(\mathbf{u}^N) z$  unless  $\mathbf{u}^N(y) = \mathbf{u}^N(z)$ .

The same example shows the error in Roberts’ attempt to prove his intermediate Lemma 6. Then Section 3 uses a modified form of the alternative “shift invariance” condition due to Roberts (1983, p. 74) himself in order to prove the crucial Lemma 6 in Roberts (1980). This establishes that a slight alteration to Claim 1 makes it valid.

## 2 Weak Continuity: A Counter Example

### 2.1 Definition of a Discontinuous Social Welfare Ordering

The following is an example of a society with two individuals and a strictly increasing and symmetric utilitarian welfare function

$$\mathbb{R}^2 \ni (u_1, u_2) = \mathbf{u} \mapsto W(u_1, u_2) = W(\mathbf{u}) \rightarrow \mathbb{R} \quad (1)$$

such that the induced SWFL defined on  $X$  by

$$a R b \iff W(u_1(a), u_2(a)) \geq W(u_1(b), u_2(b)) \quad (2)$$

satisfies conditions (U), (I), (P\*) and (WC). Yet the function  $W$  that we will define has a discontinuity at the origin  $\mathbf{0} = (0, 0)$  which implies a discontinuity in the preference ordering  $R$  on  $\mathbb{R}^2$  defined by (2). This implies that no continuous function  $W$  can satisfy Claim 1 in this example.

---

<sup>5</sup>The Roberts’ theorem which is the topic of this paper concerns social choice in the sense of aggregating preferences. It differs from the Roberts’ theorem on revelation of preferences that appeared in Roberts (1979), and has subsequently been discussed by, amongst others, Lavi et al. (2009) and Mishra and Sen (2012).

Indeed, first define the function  $\mathbb{R}^2 \ni (v_1, v_2) = \mathbf{v} \mapsto w(\mathbf{v}) \in \mathbb{R}$  by

$$w(\mathbf{v}) := \min\{v_1 + 2v_2, 2v_1 + v_2\} \quad (3)$$

Next, partition  $\mathbb{R}^2$  into the three subdomains  $S_1, S_2, S_3$  and then define the symmetric function  $S_i \ni \mathbf{v} \mapsto W(\mathbf{v}) \in \mathbb{R}$  for  $i = 1, 2, 3$  so that

$$W(\mathbf{v}) := v_1 + v_2 \quad \text{on} \quad S_1 := \{\mathbf{v} \in \mathbb{R}^2 \mid v_1 + v_2 \leq 0\} \quad (4)$$

$$W(\mathbf{v}) := 1 + w(\mathbf{v}) \quad \text{on} \quad S_2 := \{\mathbf{v} \in \mathbb{R}^2 \mid w(\mathbf{v}) > 0\} \quad (5)$$

$$W(\mathbf{v}) := \exp \frac{(v_1 + 2v_2)(2v_1 + v_2)}{3(v_1 + v_2)} \quad \text{on} \quad S_3 := \mathbb{R}^2 \setminus (S_1 \cup S_2) \quad (6)$$

The indifference map corresponding to this function is illustrated in Figure 1, which shows how:

- (i) the closed set  $S_1$  is separated from  $S_3$  by the common boundary line  $\{\mathbf{v} \in \mathbb{R}^2 \mid v_1 + v_2 = 0\}$ , which constitutes the closed indifference curve  $W(\mathbf{v}) = 0$ ;
- (ii) the open set  $S_2$  is separated from  $S_3$  by the common boundary set  $\{\mathbf{v} \in \mathbb{R}^2 \mid w(\mathbf{v}) = 0\}$ ;
- (iii)  $(0, 0)$  belongs to  $S_1$ , but is a common boundary point of all three subdomains.

Consider first the intermediate subdomain  $S_3$ , which consists of two wedges where  $v_1 + v_2 > 0$  and  $w(\mathbf{v}) \leq 0$ . On  $S_3$ , it follows from (6) that

$$\ln W(\mathbf{v}) = \frac{2(v_1)^2 + 5v_1v_2 + 2(v_2)^2}{3(v_1 + v_2)} = \frac{2(v_1 + v_2)}{3} + \frac{v_1v_2}{3(v_1 + v_2)} \quad (7)$$

In  $S_3$ , because  $v_1 + v_2 > 0$ , this is a differentiable function of  $\mathbf{v}$ . Its first partial derivative is

$$\frac{\partial}{\partial v_1} \ln W = \frac{2}{3} + \frac{v_2(v_1 + v_2) - v_1v_2}{3(v_1 + v_2)^2} = \frac{2}{3} + \frac{(v_2)^2}{3(v_1 + v_2)^2} > 0$$

By symmetry, its second partial derivative is also positive. This shows that  $S_3 \ni \mathbf{v} \mapsto W(\mathbf{v}) \in \mathbb{R}$  is strictly increasing.

Furthermore, consider any sequence  $\mathbb{N} \ni n \mapsto \mathbf{v}^n = (v_1^n, v_2^n) \in S_3$  which converges to a point  $\bar{\mathbf{v}} = (\bar{v}_1, \bar{v}_2) \neq (0, 0)$  that lies on the lower boundary of  $S_3$  because  $\bar{v}_1 + \bar{v}_2 = 0$ . Evidently it follows from (7) that, as  $n \rightarrow \infty$ , so  $\ln W(v_1^n, v_2^n) \rightarrow -\infty$ , implying that  $W(v_1^n, v_2^n) \rightarrow 0$ . Also, it follows from




Figure 1: Eight level curves of the function  $(v_1, v_2) \mapsto W(v_1, v_2)$

(3) and (6) that  $W(\mathbf{v}) = 1$  when  $\mathbf{v}$  lies on the upper boundary of  $S_3$ , where  $w(\mathbf{v}) = 0$ . So the range set  $W(S_3)$  must be the whole interval  $(0, 1]$ .

Putting these results for the function  $\mathbb{R}^2 \ni \mathbf{v} \mapsto W(\mathbf{v})$  on the subdomain  $S_3$ , together with obvious properties on the subdomains  $S_1$  and  $S_2$  where  $W$  is defined by (4) and (5), we see that the respective range sets are the three pairwise disjoint line intervals

$$W(S_1) = (-\infty, 0], \quad W(S_2) = (1, \infty), \quad W(S_3) = (0, 1]$$

From this it is evident that the function  $\mathbf{v} \mapsto W(\mathbf{v})$  is strictly increasing throughout  $\mathbb{R}^2$ , and continuous except at  $(0, 0)$ , where  $W(0, 0) = 0$ .

The three-dimensional graph of  $W(v_1, v_2)$  has a boundary that includes the vertical ‘‘cliff’’  $\{(0, 0)\} \times [0, 1]$  in  $\mathbb{R}^3$  of height 1. The base of this cliff is at the origin  $(0, 0, 0)$ , which is on the graph of  $W$  because  $W(0, 0) = 0$ . Thus, the mapping  $\mathbb{R}^2 \ni \mathbf{v} \mapsto W(\mathbf{v})$  is discontinuous at  $\mathbf{v} = (0, 0)$ . Not only is the function  $W$  discontinuous; so is the preference relation it induces. Indeed, for each fixed  $\bar{\mathbf{v}} \in S_3$  where  $W(\bar{\mathbf{v}}) \in (0, 1]$ , the upper contour set  $\{(v_1, v_2) \in \mathbb{R}^2 \mid W(v_1, v_2) \geq W(\bar{\mathbf{v}})\}$  is not closed because it excludes the point  $(0, 0)$  which is in its closure.

## 2.2 Verifying Weak Continuity

Consider now the SWFL  $\mathcal{U}^2 \ni (u_1, u_2) \mapsto f(u_1, u_2)$  defined as in Section 2.1. Obviously, this induced SWFL satisfies conditions (U), (I) and (P\*). To verify condition (WC) it is enough to construct, for each fixed  $\epsilon = (\epsilon_1, \epsilon_2) \in \mathbb{R}_{++}^2$ , a transformation

$$\mathbb{R}^2 \ni \mathbf{v} \mapsto \phi^\epsilon(\mathbf{v}) = (\phi_1^\epsilon(\mathbf{v}), \phi_2^\epsilon(\mathbf{v})) \in \mathbb{R}^2$$

satisfying

$$(0, 0) \ll (v_1, v_2) - \phi^\epsilon(\mathbf{v}) \ll \epsilon$$

together with the requirement that  $\mathbf{v} \mapsto W(\phi^\epsilon(\mathbf{v}))$  and  $\mathbf{v} \mapsto W(\mathbf{v})$  are ordinally equivalent welfare functions in the sense that there exists a strictly increasing transformation  $\mathbb{R} \ni \mathbf{w} \mapsto \psi^\epsilon(\mathbf{w}) \in \mathbb{R}$  for which  $W(\phi^\epsilon(\mathbf{v})) \equiv \psi^\epsilon(W(\mathbf{v}))$ .

In the following constructions, given any fixed  $\epsilon = (\epsilon_1, \epsilon_2) \in \mathbb{R}_{++}^2$ , let

$$\epsilon_* := \min\{\epsilon_1, \epsilon_2\} \in \mathbb{R} \text{ and } \mathbf{e} := (1, 1) \in \mathbb{R}^2 \tag{8}$$

Then  $\epsilon_* > 0$ , of course. The transformation will take the form

$$\mathbb{R}^2 \ni \mathbf{v} \mapsto \phi^\epsilon(\mathbf{v}) := \mathbf{v} - \lambda^\epsilon(\mathbf{v}) \mathbf{e} \in \mathbb{R}^2 \tag{9}$$



for a suitably constructed scalar function  $\mathbb{R}^2 \ni \mathbf{v} \mapsto \lambda^\epsilon(\mathbf{v}) \in \mathbb{R}$  taking values in the open interval  $(0, \epsilon_*)$ .

**Case 1:** The simplest case is when

$$v_1 + v_2 \leq 0 \text{ and so } W(\mathbf{v}) = v_1 + v_2 \leq 0 \quad (10)$$

In this case, define  $\lambda^\epsilon(\mathbf{v}) := \frac{1}{2} \epsilon_*$  for  $\epsilon_*$  given by (8). Then (9) implies that

$$\phi^\epsilon(\mathbf{v}) = (v_1 - \frac{1}{2} \epsilon_*, v_2 - \frac{1}{2} \epsilon_*)$$

So (9) and (10) imply that  $\phi_1^\epsilon(\mathbf{v}) + \phi_2^\epsilon(\mathbf{v}) = v_1 + v_2 - \epsilon_* \leq -\epsilon_* < 0$ . Now, whenever  $v_1 + v_2 \leq 0$ , it follows that

$$W(\phi^\epsilon(\mathbf{v})) = \phi_1^\epsilon(\mathbf{v}) + \phi_2^\epsilon(\mathbf{v}) = \psi^\epsilon(W(\mathbf{v}))$$

provided we define

$$\psi^\epsilon(W) := W - \epsilon_* \text{ for all } W \leq 0 \quad (11)$$

**Case 2:** This case occurs when

$$w(\mathbf{v}) > 0 \text{ and so } W(\mathbf{v}) = 1 + w(\mathbf{v}) > 1 \quad (12)$$

In this case, define

$$\lambda^\epsilon(\mathbf{v}) := \frac{1}{6} \min\{\epsilon_*, w(\mathbf{v})\} \quad (13)$$

Clearly, this definition implies that  $\lambda^\epsilon(\mathbf{v}) \in (0, \epsilon_*)$ . Also

$$\phi_1^\epsilon(\mathbf{v}) + 2\phi_2^\epsilon(\mathbf{v}) = v_1 + 2v_2 - 3\lambda^\epsilon(\mathbf{v}) \quad (14)$$

$$\text{and } 2\phi_1^\epsilon(\mathbf{v}) + \phi_2^\epsilon(\mathbf{v}) = 2v_1 + v_2 - 3\lambda^\epsilon(\mathbf{v}) \quad (15)$$

But  $w(\mathbf{v}) := \min\{v_1 + 2v_2, 2v_1 + v_2\}$  and (13) implies that  $\lambda^\epsilon(\mathbf{v}) \leq \frac{1}{6} w(\mathbf{v})$ . So it follows from (14) and (15) that

$$\min\{\phi_1^\epsilon(\mathbf{v}) + 2\phi_2^\epsilon(\mathbf{v}), 2\phi_1^\epsilon(\mathbf{v}) + \phi_2^\epsilon(\mathbf{v})\} = w(\mathbf{v}) - 3\lambda^\epsilon(\mathbf{v}) \geq \frac{1}{2} w(\mathbf{v}) > 0 \quad (16)$$

So the definition of  $\mathbb{R}^2 \ni \mathbf{v} \mapsto W(\mathbf{v}) \in \mathbb{R}$  in (5) and of  $\mathbb{R}^2 \ni \mathbf{v} \mapsto \lambda^\epsilon(\mathbf{v}) \in \mathbb{R}$  in (13) imply, together with (16), that

$$\begin{aligned} 1 < W(\phi^\epsilon(\mathbf{v})) &= 1 + \min\{\phi_1^\epsilon(\mathbf{v}) + 2\phi_2^\epsilon(\mathbf{v}), 2\phi_1^\epsilon(\mathbf{v}) + \phi_2^\epsilon(\mathbf{v})\} \\ &= 1 + w(\mathbf{v}) - 3\lambda^\epsilon(\mathbf{v}) \\ &= 1 + w(\mathbf{v}) - \frac{1}{2} \min\{\epsilon_*, w(\mathbf{v})\} \\ &= \max\{1 + w(\mathbf{v}) - \frac{1}{2} \epsilon_*, 1 + \frac{1}{2} w(\mathbf{v})\} \\ &= \max\{W(\mathbf{v}) - \frac{1}{2} \epsilon_*, \frac{1}{2} [W(\mathbf{v}) + 1]\} \end{aligned}$$

It follows that  $W(\phi^\epsilon(\mathbf{v})) = \psi^\epsilon(W(\mathbf{v}))$  provided that we define

$$\psi^\epsilon(W) := \max\{W - \frac{1}{2}\epsilon_*, \frac{1}{2}(W + 1)\} \text{ for all } W > 1 \quad (17)$$

**Case 3:** This leaves the hardest third case, when

$$w(\mathbf{v}) \leq 0 \text{ and also } v_1 + v_2 > 0 \quad (18)$$

In this case, the definition in (6) implies that  $0 < W(\mathbf{v}) \leq 1$ .

Fix any  $\mathbf{v} = (v_1, v_2) \in \mathbb{R}^2$  satisfying (18). Then, given any  $\epsilon \in \mathbb{R}^2$  satisfying  $\epsilon \gg 0$ , consider the non-empty open interval of  $\mathbb{R}$  defined by

$$I^\epsilon(\mathbf{v}) := (0, \min\{\epsilon_*, \frac{1}{2}(v_1 + v_2)\}) = (0, \epsilon_*) \cap (0, \frac{1}{2}(v_1 + v_2)) \quad (19)$$

Now consider the function  $g$  defined on the open interval in (19) by

$$I^\epsilon(\mathbf{v}) \ni \lambda \mapsto g(\lambda) := W(\mathbf{v} - \lambda \mathbf{e}) \in \mathbb{R} \quad (20)$$

In Section 2.1 we saw that  $W$  is strictly increasing as a function of two variables, so the function  $g$  is strictly decreasing. Also, when  $\lambda > 0$ , it is evident that

$$w(\mathbf{v} - \lambda \mathbf{e}) < w(\mathbf{v}) \leq 0 \quad (21)$$

On the other hand, when  $\lambda < \frac{1}{2}(v_1 + v_2)$ , because  $e_1 = e_2 = 1$ , one has

$$(v_1 - \lambda e_1) + (v_2 - \lambda e_2) = v_1 + v_2 - 2\lambda > 0 \quad (22)$$

So for all  $\lambda \in I^\epsilon(\mathbf{v})$  the inequalities (21) and (22) imply that the 2-vector  $\mathbf{v} - \lambda \mathbf{e}$  satisfies (18). It follows that  $W(\mathbf{v} - \lambda \mathbf{e})$  is also defined by (6), implying that

$$g(\lambda) = W(\mathbf{v} - \lambda \mathbf{e}) = \exp \frac{(v_1 + 2v_2 - 3\lambda)(2v_1 + v_2 - 3\lambda)}{3(v_1 + v_2 - 2\lambda)} \quad (23)$$

Now let  $\mu$  denote a suitably chosen positive scalar constant which is independent of both  $\mathbf{v}$  and  $\epsilon$ , and whose possible range will be specified later. For each  $\mathbf{v} \in \mathbb{R}^2$  that satisfies the inequalities (18), because  $g$  is strictly decreasing and positive, we can define  $\lambda^\epsilon(\mathbf{v})$  implicitly as the unique value of  $\lambda$  that solves the equation

$$g(\lambda) = W(\mathbf{v} - \lambda \mathbf{e}) = W(\mathbf{v}) \exp(-\mu \epsilon_*) \quad (24)$$

Then  $\lambda^\epsilon(\mathbf{v})$  will be well defined and positive, with

$$W(\phi^\epsilon(\mathbf{v})) = W(\mathbf{v}) \exp(-\mu \epsilon_*) = \psi^\epsilon(W(\mathbf{v})) < 1$$

where  $\psi^\epsilon(W) := W \exp(-\mu \epsilon_*) \in (0, 1)$  whenever  $0 < W \leq 1$ .

It remains only to choose  $\mu > 0$  so that the corresponding solution to equation (24) exists in the open interval  $I^\epsilon(\mathbf{v})$  defined by (19) and so satisfies  $\lambda^\epsilon(\mathbf{v}) < \epsilon_*$ . Because we are assuming that the inequalities (18) hold, definition (6) implies that any  $\lambda^\epsilon(\mathbf{v})$  satisfying (24) and (19) must be a value of  $\lambda$  which solves the equation

$$\frac{(v_1 + 2v_2 - 3\lambda)(2v_1 + v_2 - 3\lambda)}{3(v_1 + v_2 - 2\lambda)} = \frac{(v_1 + 2v_2)(2v_1 + v_2)}{3(v_1 + v_2)} - \mu \epsilon_*$$

But  $v_1 + v_2 > 2\lambda > 0$  in the relevant interval of values of  $\lambda$ , so we can clear fractions to obtain the quadratic equation  $q(\lambda) = 0$ , where

$$\begin{aligned} q(\lambda) := & (v_1 + v_2)(v_1 + 2v_2 - 3\lambda)(2v_1 + v_2 - 3\lambda) \\ & - (v_1 + v_2 - 2\lambda)(v_1 + 2v_2)(2v_1 + v_2) \\ & + 3\mu \epsilon_* (v_1 + v_2)(v_1 + v_2 - 2\lambda) \end{aligned} \quad (25)$$

Now, note that when  $\lambda = 0$  the first two terms on the right-hand side of (25) cancel. Because  $v_1 + v_2 > 0$ , it follows that

$$q(0) = 3\mu \epsilon_* (v_1 + v_2)^2 > 0 \quad (26)$$

In addition, simple calculation shows that

$$q\left(\frac{1}{2}(v_1 + v_2)\right) = -\frac{1}{4}(v_1 + v_2)(v_1 - v_2)^2 \quad (27)$$

Finally, some much more tedious but still routine algebraic manipulation shows that

$$q(\epsilon_*) = (v_1 + v_2) \epsilon_* [(9 - 6\mu) \epsilon_* + (3\mu - 5)(v_1 + v_2)] + 2v_1 v_2 \epsilon_* \quad (28)$$

Because  $v_1 + v_2 > 0$  but  $w(\mathbf{v}) \leq 0$ , it follows from (3) that  $v_1$  and  $v_2$  have opposite signs. In particular  $v_1 \neq v_2$  and  $v_1 v_2 < 0$ . Then (27) implies that  $q(\frac{1}{2}(v_1 + v_2)) < 0$  whereas (28) implies that for any  $\mu$  satisfying  $9 < 6\mu < 10$  one has  $q(\epsilon_*) < 0$ . So choosing any fixed  $\mu \in (\frac{3}{2}, \frac{5}{3})$  guarantees that, by the intermediate value theorem, the quadratic equation  $q(\lambda) = 0$  has a unique root  $\lambda^\epsilon(\mathbf{v})$  in the open interval  $I^\epsilon(\mathbf{v}) = (0, \min\{\epsilon_*, \frac{1}{2}(v_1 + v_2)\})$  defined by (19).<sup>6</sup> In particular, for each  $\mathbf{v}$  satisfying (18), the root  $\lambda^\epsilon(\mathbf{v})$  of  $q(\lambda) = 0$  that we have found lies in  $(0, \epsilon_*)$ , as required.

---

<sup>6</sup>Because  $q(\lambda) \rightarrow +\infty$  as  $\lambda \rightarrow +\infty$ , the quadratic equation  $q(\lambda) = 0$  has a second real root that satisfies  $\lambda > \max\{\epsilon_*, \frac{1}{2}(v_1 + v_2)\}$ . But this root is irrelevant.

Finally, putting all three cases together gives  $W(\phi^\epsilon(\mathbf{v})) \equiv \psi^\epsilon(W(\mathbf{v}))$ , where  $\phi^\epsilon(\mathbf{v}) = \mathbf{v} - \lambda^\epsilon(\mathbf{v}) \mathbf{e}$ , and then

$$\psi^\epsilon(W) := \begin{cases} W - \frac{1}{2}(\epsilon_1 + \epsilon_2) < 0 & \text{if } W \leq 0; \\ W \exp(-\mu \epsilon_*) \in (0, 1] & \text{if } 0 < W \leq 1; \\ \max\{W - \frac{1}{2}\epsilon_*, \frac{1}{2}(W + 1)\} > 1 & \text{if } W > 1. \end{cases}$$

In particular,  $\psi^\epsilon$  is strictly increasing in  $W$  for each fixed  $\epsilon \gg 0$ .  $\square$

### 2.3 Diagnosis

Next, to see where his proof erred, we introduce a definition that incorporates some more notation from Roberts (1980, pp. 425–6).

**Definition 1.** *Given the SWFL  $f$ , the strict preference relation  $\succ$  on utility vectors in  $\mathbb{R}^N$  is defined so that  $\mathbf{a} \succ \mathbf{b}$  just in case there exist a utility function profile  $\mathbf{u}^N \in \mathcal{D}$  and two social states  $x, y \in X$  with  $x f(\mathbf{u}^N) y$  such that  $\mathbf{a} \gg \mathbf{u}(x)$  and  $\mathbf{u}(y) \gg \mathbf{b}$ . Then, for each  $\mathbf{v}^* \in \mathbb{R}^N$ , the three sets  $L(\mathbf{v}^*)$ ,  $M(\mathbf{v}^*)$  and  $N(\mathbf{v}^*)$  are defined respectively by*

$$L(\mathbf{v}^*) := \{\mathbf{v} \in \mathbb{R}^N \mid \mathbf{v}^* \succ \mathbf{v}\} \quad (29)$$

$$M(\mathbf{v}^*) := \{\mathbf{v} \in \mathbb{R}^N \mid \mathbf{v} \succ \mathbf{v}^*\} \quad (30)$$

$$N(\mathbf{v}^*) := \mathbb{R}^N \setminus [L(\mathbf{v}^*) \cup M(\mathbf{v}^*)] \quad (31)$$

Now, in the above example the set  $N(\mathbf{0})$  is equal to the middle region where  $W(\mathbf{v}) \in (0, 1]$ . Note too that, although  $\mathbf{v} \in N(\mathbf{0})$  whenever  $W(\mathbf{v}) \in (0, 1]$ , one will have  $\mathbf{v} - \boldsymbol{\eta} \succ -\boldsymbol{\eta}'$  whenever  $\boldsymbol{\eta}, \boldsymbol{\eta}' \gg \mathbf{0}$  with  $\boldsymbol{\eta}$  small enough so that  $W(\mathbf{v}) \geq 0$  because  $\eta_1 + \eta_2 \leq v_1 + v_2$ . This contradicts Roberts' claim, in the course of trying to prove Lemma 6, that: "...as  $\mathbf{v} + \boldsymbol{\gamma} \in N(\mathbf{v}^*)$ , [condition] (WC) ensures that  $\mathbf{v} + \boldsymbol{\gamma} - \boldsymbol{\eta}_3 \in N(\mathbf{v}^* - \boldsymbol{\eta}_4)$  for some  $\epsilon \gg \boldsymbol{\eta}_3, \boldsymbol{\eta}_4 \gg \mathbf{0}$  where  $\epsilon$  is subject to choice."

## 3 A New Sufficient Condition

### 3.1 Statement of Pairwise Continuity

Roberts (1983, p. 74) later introduced a *shift invariance* condition which can be slightly restated as follows:

**Condition (SI)** Given any  $\epsilon \in \mathbb{R}_{++}^N$ , for all profiles  $\mathbf{u}^N \in \mathcal{D}$ , there exists an  $\epsilon' \in \mathbb{R}_{++}^N$  and a profile  $\tilde{\mathbf{u}}^N \in \mathcal{D}$  such that  $f(\mathbf{u}^N) = f(\tilde{\mathbf{u}}^N)$  and, for all  $x \in X$ , one has  $\epsilon \gg \mathbf{u}^N(x) - \tilde{\mathbf{u}}^N(x) \gg \epsilon'$ .

As he states in a footnote: “Shift invariance is slightly stronger than ... (WC). ... The strengthening allows one to deal with problems that are akin to the existence of poles in a consumer’s indifference map ... .”<sup>7</sup> However, when proving his Lemma A.5, it seems that Roberts (1983, p. 90) in the end reverses the order of the quantified statements “for all profiles  $\mathbf{u}^N \in \mathcal{U}^N$ ” and “there exists an  $\boldsymbol{\epsilon}' \in \mathbb{R}_{++}^N$ ” and actually uses the following *uniform shift invariance* assumption:

**Condition (USI)** Given any  $\boldsymbol{\epsilon} \in \mathbb{R}_{++}^N$ , there exists an  $\boldsymbol{\epsilon}' \in \mathbb{R}_{++}^N$  for which, for all profiles  $\mathbf{u}^N \in \mathcal{D}$ , there exists a profile  $\tilde{\mathbf{u}}^N \in \mathcal{D}$  such that  $f(\mathbf{u}^N) = f(\tilde{\mathbf{u}}^N)$  and, for all  $x \in X$ , one has  $\boldsymbol{\epsilon} \gg \mathbf{u}^N(x) - \tilde{\mathbf{u}}^N(x) \gg \boldsymbol{\epsilon}'$ .

Instead of (WC) or (SI), I shall use the following *pairwise continuity* assumption which weakens (USI):

**Condition (PC)** Given any  $\boldsymbol{\epsilon} \in \mathbb{R}_{++}^N$ , there exists an  $\boldsymbol{\epsilon}' \in \mathbb{R}_{++}^N$  for which, for all profiles  $\mathbf{u}^N \in \mathcal{D}$ , given any pair  $x, y \in X$  satisfying  $x P(\mathbf{u}^N) y$ , there exists a profile  $\tilde{\mathbf{u}}^N \in \mathcal{D}$  with  $\tilde{\mathbf{u}}^N(x) \ll \mathbf{u}^N(x) - \boldsymbol{\epsilon}'$  and  $\tilde{\mathbf{u}}^N(y) \gg \mathbf{u}^N(y) - \boldsymbol{\epsilon}$  such that  $x P(\tilde{\mathbf{u}}^N) y$ .

Like shift invariance, condition (PC) strengthens weak continuity because the same strictly positive vector  $\boldsymbol{\epsilon}'$  must work *simultaneously* for all  $x, y \in X$ . Like uniform shift invariance, it also strengthens shift invariance because the same strictly positive vector  $\boldsymbol{\epsilon}'$  must also work for all profiles  $\mathbf{u}^N \in \mathcal{D}$ . On the other hand, condition (PC) weakens even condition (WC), as well as condition (USI), to the extent that the profile  $\tilde{\mathbf{u}}^N$  can depend on the pair  $x, y \in X$ , and also because it requires only one-way strict inequalities which, moreover, are different for  $x$  and  $y$ .

### 3.2 Sufficiency of Pairwise Continuity

With condition (PC) replacing (WC), Lemma 6 of Roberts (1980) will be proved via the following two separate lemmas that involve definitions (30) and (31):

**Lemma 1.** *If  $f$  satisfies (U), (I) and (P), then for all  $\mathbf{v}, \mathbf{v}', \boldsymbol{\eta}, \boldsymbol{\eta}' \in \mathbb{R}^N$  with  $\boldsymbol{\eta}, \boldsymbol{\eta}' \gg \mathbf{0}$ , one has  $\mathbf{v} \in N(\mathbf{v}') \implies \mathbf{v} + \boldsymbol{\eta} \in M(\mathbf{v}' - \boldsymbol{\eta}')$ .*<sup>8</sup>

<sup>7</sup>Indeed, it is this footnote that suggested to me how the above counter example might be constructed.

<sup>8</sup>This is the correct “preliminary result” in Roberts’ discussion of Lemma 6. However, it seemed that the proof provided there could benefit from more detail.

*Proof.* Suppose that  $x, y, z$  are three distinct elements of  $X$ . By condition (U), there exists a profile  $\mathbf{u}^N \in \mathcal{D}$  such that

$$\mathbf{v} + \boldsymbol{\eta} \gg \mathbf{u}^N(x) \gg \mathbf{u}^N(y) \gg \mathbf{v} \text{ and } \mathbf{v}' \gg \mathbf{u}^N(z) \gg \mathbf{v}' - \boldsymbol{\eta}' \quad (32)$$

By Definition 1, if  $z P(\mathbf{u}^N) y$  were true, it would imply that  $\mathbf{v}' \succ \mathbf{v}$ . So  $\mathbf{v} \in N(\mathbf{v}') \implies y R(\mathbf{u}^N) z$ . Then the Pareto condition (P) implies that  $x P(\mathbf{u}^N) y$ , and so  $\mathbf{v} \in N(\mathbf{v}') \implies x P(\mathbf{u}^N) z$  because  $R(\mathbf{u}^N)$  is transitive. From (32) it follows that  $\mathbf{v} \in N(\mathbf{v}') \implies \mathbf{v} + \boldsymbol{\eta} \succ \mathbf{v}' - \boldsymbol{\eta}'$ .  $\square$

Part (b) of the following Lemma is a minor restatement of the conclusion of Lemma 6 in Roberts (1980):

**Lemma 2.** *If  $f$  satisfies (U), (I), (P) and (PC), then:*

(a) *if  $\boldsymbol{\epsilon}, \mathbf{v}, \mathbf{v}'$  in  $\mathbb{R}^N$  satisfy  $\boldsymbol{\epsilon} \gg \mathbf{0}$  as well as  $\mathbf{v} + \boldsymbol{\eta} \in M(\mathbf{v}' + \boldsymbol{\epsilon})$  for all  $\boldsymbol{\eta} \gg \mathbf{0}$ , then  $\mathbf{v} \in M(\mathbf{v}')$ ;*

(b) *for all  $\mathbf{v}, \mathbf{v}^*$  in  $\mathbb{R}^N$  that satisfy  $\mathbf{v} \in N(\mathbf{v}^*)$ , there is no  $\boldsymbol{\gamma} \in \mathbb{R}_{++}^N$  such that  $\mathbf{v} + \boldsymbol{\gamma} \in N(\mathbf{v}^*)$ .*

*Proof.* (a) Given  $\boldsymbol{\epsilon} \gg \mathbf{0}$ , let  $\boldsymbol{\epsilon}' \gg \mathbf{0}$  be specified as in the statement of condition (PC). Choose  $\boldsymbol{\eta} \gg \mathbf{0}$  so that  $\boldsymbol{\eta} \ll \boldsymbol{\epsilon}'$ . Because  $\mathbf{v} + \boldsymbol{\eta} \in M(\mathbf{v}' + \boldsymbol{\epsilon})$ , condition (U) and Definition 1 imply that there exist  $\mathbf{u}^N \in \mathcal{D}$  and  $x, y \in X$  such that  $x P(\mathbf{u}^N) y$  while

$$\mathbf{v} + \boldsymbol{\eta} \gg \mathbf{u}^N(x) \text{ and } \mathbf{u}^N(y) \gg \mathbf{v}' + \boldsymbol{\epsilon} \quad (33)$$

By condition (PC), there exists  $\tilde{\mathbf{u}}^N \in \mathcal{D}$  such that  $x P(\tilde{\mathbf{u}}^N) y$  while

$$\tilde{\mathbf{u}}^N(x) \ll \mathbf{u}^N(x) - \boldsymbol{\epsilon}' \text{ and } \tilde{\mathbf{u}}^N(y) \gg \mathbf{u}^N(y) - \boldsymbol{\epsilon} \quad (34)$$

But then (33) and (34) together imply that  $\tilde{\mathbf{u}}^N(x) \ll \mathbf{v} + \boldsymbol{\eta} - \boldsymbol{\epsilon}' \ll \mathbf{v}$ , because of the choice of  $\boldsymbol{\eta}$ ; they also imply that  $\tilde{\mathbf{u}}^N(y) \gg \mathbf{v}'$ . Because  $x P(\tilde{\mathbf{u}}^N) y$ , it follows from Definition 1 that  $\mathbf{v} \succ \mathbf{v}'$ .

(b) Suppose that  $\mathbf{v} + \boldsymbol{\gamma} \in N(\mathbf{v}^*)$ . By definition (31), it follows that  $\mathbf{v}^* \in N(\mathbf{v} + \boldsymbol{\gamma})$ . Choose any  $\boldsymbol{\gamma}' \gg \mathbf{0}$  satisfying  $\boldsymbol{\gamma}' \ll \boldsymbol{\gamma}$ . Now Lemma 1 implies that  $\mathbf{v}^* + \boldsymbol{\eta} \in M(\mathbf{v} + \boldsymbol{\gamma}')$  for all  $\boldsymbol{\eta} \gg \mathbf{0}$ . So part (a) implies that  $\mathbf{v}^* \in M(\mathbf{v})$ . In particular,  $\mathbf{v} \notin N(\mathbf{v}^*)$ .  $\square$

### 3.3 Invariance Conditions and Pairwise Continuity<sup>9</sup>

Just as with Roberts' (WC) and (SI) conditions, condition (USI) and so (PC) is certainly satisfied if  $f$  is invariant under a set of transformations of individual utility functions large enough to include all shifts of the form  $\tilde{u}_i(x) \equiv \alpha + u_i(x)$  (for all  $i \in N$  and  $x \in X$ ) with  $\alpha \in \mathbb{R}$  independent of  $i$ . Of the six invariance classes  $\Phi$  presented on p. 423 of Roberts (1980), this is true of the first five, namely (ONC), (CNC), (OLC), (CUC), and (CFC).

It remains to discuss the sixth cardinal ratio-scale or (CRS) class, as well as the broader non-comparable ratio-scale or (NRS) invariance class that includes all transformations of the form  $\tilde{u}_i(x) \equiv \beta_i u_i(x)$  (for all  $i \in N$  and  $x \in X$ ) with  $\beta_i > 0$  for each individual  $i \in N$ . Indeed, since the unrestricted domain condition (U) allows each utility function to have both positive and negative values, there are difficulties with the key condition (PC) in this case.<sup>10</sup>

When Roberts (1980, p. 423) discusses the (CRS) class, he states that "For simplicity, it will be assumed that welfares are always strictly positive." This, of course, violates condition (U), but almost all the relevant results are easily restated for any restricted domain  $\mathcal{D} \subset \mathcal{U}^N$  large enough to include all profiles of utility functions whose values are strictly positive. Nevertheless, the key pairwise continuity condition (PC) is violated because, no matter how small  $\epsilon' \in \mathbb{R}_{++}^N$  may be, for any  $x \in X$  there will always be a strictly positive-valued profile  $\mathbf{u}^N \in \mathcal{D}$  such that  $\mathbf{u}^N(x) \ll \epsilon'$ ; this makes it impossible to find any profile  $\tilde{\mathbf{u}}^N$  of strictly positive-valued utility functions that satisfies  $\tilde{\mathbf{u}}^N(x) \ll \mathbf{u}^N(x) - \epsilon'$ .

When utilities are restricted to have strictly positive values throughout  $X$ , however, one can work instead with  $\ln u_i(x)$  as a transformed utility function. Then ratio-scale invariance for utilities is equivalent to invariance under additive shifts of their logarithms. Moreover, condition (PC) can be restated for these transformed utility functions. A similar trick works when utility functions are restricted to have strictly negative values; one works instead with  $-\ln[-u_i(x)]$  as a transformed utility function. The (CRS) and (NRS) invariance classes present problems only when utilities are allowed to change signs, or to become zero.

<sup>9</sup>This subsection in particular owes much to John Weymark and the anonymous referee.

<sup>10</sup>There are also difficulties with Lemma 8 of Roberts (1980). We note that Blackorby and Donaldson (1982) as well as Tsui and Weymark (1997), in their work on ratio-scale invariant social welfare functionals, imposed a more direct continuity condition on a social preference ordering which is defined on the space of utility vectors.

## 4 Conclusion

The weak neutrality or welfarism theorem due to Roberts (1980) is indeed “both important and useful” (p. 428). The minor errors in its statement and in the proof of the key Lemma 6 are not very difficult to correct by replacing the weak continuity condition (WC) with the new pairwise continuity condition (PC) stated here in Section 3.1.

An open question is whether the closely related Theorem 1 of Roberts (1983) holds under shift invariance (SI) instead of uniform shift invariance (USI), which is stronger than (PC). However, even (USI) is weak enough that having to impose it instead of (WC) or (SI) would do little to detract from the significance or wide applicability of Roberts’ theorem.

Only in the case of ratio-scale measurability of utilities that can change sign does Roberts’ theorem seem inapplicable. With zero as the utility of the infinite set of potential people who, in each possible social state, never come into existence, this happens to be exactly the setting that we consider in Chichilnisky et al. (2020). But in that paper we extend the kind of original position due to Vickrey (1945) and Harsanyi (1953, 1955). Then we use an “impartial benefactor” argument to derive a utilitarian SWFL more directly.

## Acknowledgements

An earlier version appeared as Stanford University Department of Economics Working Paper No. 99-021, which is available at <https://web.stanford.edu/~hammond/roberts.pdf>. Later, during John Weymark’s term as an editor of *Social Choice and Welfare*, he very kindly encouraged me to submit a revised version. This seems an excellent occasion on which to follow his suggestion. Finally, I am grateful for the most helpful comments of a diligent referee.

## References

- Arrow, K.J. (1963) *Social Choice and Individual Values*, 2nd edn. (New Haven: Yale University Press).
- Blackorby, C. and D. Donaldson (1982) “Ratio-scale and translation-scale full interpersonal comparability without domain restrictions: Admissible social-evaluation functions” *International Economic Review* 23: 249–268.
- Blackorby, C. D. Donaldson and J.A. Weymark (1984) “Social choice with interpersonal utility comparisons: A diagrammatic introduction” *International Economic Review* 25: 327–356.



- Bordes, G., P.J. Hammond and M. Le Breton (2005), “Social welfare functionals on restricted domains and in economic environments” *Journal of Public Economic Theory* 7: 1–25.
- Bordes, G. and M. Le Breton (1987), “Roberts’s theorem in economic environments: (2) The private goods case” preprint, Faculté des Sciences Economiques et de Gestion, Université de Bordeaux I.
- Bossert, W and J.A. Weymark (2004) “Utility in social choice,” in *Handbook of Utility Theory, Vol. II: Applications and Extensions* by S. Barberà, P.J. Hammond, and C. Seidl, eds. (Kluwer: Dordrecht).
- Chichilnisky, G., P.J. Hammond and N. Stern (2020) “Fundamental utilitarianism and intergenerational equity with extinction discounting” *Social Choice and Welfare* 54: 397–427.
- D’Aspremont, C. (1985) “Axioms for social welfare orderings” in *Social Goals and Social Organization: Essays in Memory of Elisha Pazner* L. Hurwicz, D. Schmeidler and H. Sonnenschein (eds.) (Cambridge University Press: Cambridge) ch. 1, pp. 19–76.
- D’Aspremont, C. and L. Gevers (1977) “Equity and the informational basis of collective choice” *Review of Economic Studies* 44: 199–209.
- Harsanyi, J.C. (1953) “Cardinal utility in welfare economics and in the theory of risk-taking” *Journal of Political Economy* 61: 434–435.
- Harsanyi, J.C. (1955) “Cardinal welfare, individualistic ethics, and interpersonal comparisons of utility” *Journal of Political Economy* 63: 309–321.
- Lavi, R., A. Mu’alem and N. Nisan (2009) “Two simplified proofs for Roberts’ theorem” *Social Choice and Welfare* 32: 407–423.
- Le Breton, M. (1987) “Roberts’ theorem in economic environments: 1. — The public goods case” preprint, Faculté des Sciences Economiques et de Gestion, Université de Rennes.
- Mishra, D. and A. Sen (2012) “Roberts’ theorem with neutrality: A social welfare ordering approach” *Games and Economic Behavior* 75: 283–298.
- Mongin, P. and C. d’Aspremont (1998) “Utility theory and ethics” in *Handbook of Utility Theory, Vol. 1: Principles* S. Barberà, P.J. Hammond, and C. Seidl (eds.) (Kluwer: Dordrecht) ch. 10, pp. 371–481.
- Roberts, K.W.S. (1979) “The characterization of implementable choice rules” in *Aggregation and Revelation of Preferences* J.-J. Laffont (ed.) (Amsterdam: North-Holland) 321–349.
- Roberts, K.W.S. (1980) “Interpersonal comparability and social choice theory” *Review of Economic Studies* 47: 421–439.
- Roberts, K.W.S. (1983) “Social choice rules and real valued representations” *Journal of Economic Theory* 29: 72–94.

- Sen, A.K. (1970) *Collective Choice and Social Welfare* (San Francisco: Holden-Day).
- Sen, A.K. (1977) "On weights and measures: Informational constraints in social welfare analysis" *Econometrica* 45: 1539–72.
- Sen, A.K. (1984) "Social choice theory" in *Handbook of Mathematical Economics, Vol. 3* K.J. Arrow and M.D. Intriligator (eds.) (North-Holland: Amsterdam) ch. 22, pp. 1073–1181.
- Tsui, K.-Y. and J.A. Weymark (1997) "Social welfare orderings for ratio-scale measurable utilities" *Economic Theory* 10: 241–256.
- Vickrey, W.S. (1945) "Measuring marginal utility by reactions to risk" *Econometrica* 13: 319–333.